

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
24 December 2003 (24.12.2003)

PCT

(10) International Publication Number  
**WO 03/105855 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 31/535**,  
31/40, C07D 265/30, 207/02, A61P 43/00

William, F. [US/US]; 126 East Lincoln Avenue, Rahway,  
NJ 07065-0907 (US). **TASBER, Edward, S.** [US/US];  
126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(21) International Application Number: PCT/US03/18482

(74) Common Representative: **MERCK & CO., INC.**; 126  
East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(22) International Filing Date: 12 June 2003 (12.06.2003)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,  
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VC, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:  
60/388,621 14 June 2002 (14.06.2002) US  
60/403,830 15 August 2002 (15.08.2002) US  
60/426,940 15 November 2002 (15.11.2002) US  
60/458,318 28 March 2003 (28.03.2003) US

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **MERCK  
& CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway,  
NJ 07065-0907 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ARRINGTON,  
Kenneth, L.** [US/US]; 126 East Lincoln Avenue, Rahway,  
NJ 07065-0907 (US). **COLEMAN, Paul, J.** [US/US];  
126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).  
**COX, Christopher, D.** [US/US]; 126 East Lincoln Ave-  
nue, Rahway, NJ 07065-0907 (US). **FRALEY, Mark,  
E.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ  
07065-0907 (US). **GARBACCIO, Robert, M.** [US/US];  
126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).  
**HARTMAN, George, D.** [US/US]; 126 East Lincoln  
Avenue, Rahway, NJ 07065-0907 (US). **HOFFMAN,**

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: MITOTIC KINESIN INHIBITORS

(57) Abstract: The present invention relates to dihydropyrrole compounds that are useful for treating cellular proliferative diseases, for treating disorders associated with KSP kinesin activity, and for inhibiting KSP kinesin. The invention also related to compositions which comprise these compounds, and methods of using them to treat cancer in mammals.



**WO 03/105855 A1**

TITLE OF THE INVENTION  
MITOTIC KINESIN INHIBITORS

BACKGROUND OF THE INVENTION

5                   This invention relates to dihydropyrrole derivatives that are inhibitors of mitotic kinesins, in particular the mitotic kinesin KSP, and are useful in the treatment of cellular proliferative diseases, for example cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders and inflammation.

                  Among the therapeutic agents used to treat cancer are the taxanes and  
10   vinca alkaloids. Taxanes and vinca alkaloids act on microtubules, which are present in a variety of cellular structures. Microtubules are the primary structural element of the mitotic spindle. The mitotic spindle is responsible for distribution of replicate copies of the genome to each of the two daughter cells that result from cell division. It is presumed that disruption of the mitotic spindle by these drugs results in inhibition of  
15   cancer cell division, and induction of cancer cell death. However, microtubules form other types of cellular structures, including tracks for intracellular transport in nerve processes. Because these agents do not specifically target mitotic spindles, they have side effects that limit their usefulness.

                  Improvements in the specificity of agents used to treat cancer is of  
20   considerable interest because of the therapeutic benefits which would be realized if the side effects associated with the administration of these agents could be reduced. Traditionally, dramatic improvements in the treatment of cancer are associated with identification of therapeutic agents acting through novel mechanisms. Examples of this include not only the taxanes, but also the camptothecin class of topoisomerase I  
25   inhibitors. From both of these perspectives, mitotic kinesins are attractive targets for new anti-cancer agents.

                  Mitotic kinesins are enzymes essential for assembly and function of the mitotic spindle, but are not generally part of other microtubule structures, such as in nerve processes. Mitotic kinesins play essential roles during all phases of mitosis.  
30   These enzymes are "molecular motors" that transform energy released by hydrolysis of ATP into mechanical force which drives the directional movement of cellular cargoes along microtubules. The catalytic domain sufficient for this task is a compact structure of approximately 340 amino acids. During mitosis, kinesins organize microtubules into the bipolar structure that is the mitotic spindle. Kinesins mediate  
35   movement of chromosomes along spindle microtubules, as well as structural changes

in the mitotic spindle associated with specific phases of mitosis. Experimental perturbation of mitotic kinesin function causes malformation or dysfunction of the mitotic spindle, frequently resulting in cell cycle arrest and cell death.

Among the mitotic kinesins which have been identified is KSP. KSP  
5 belongs to an evolutionarily conserved kinesin subfamily of plus end-directed microtubule motors that assemble into bipolar homotetramers consisting of antiparallel homodimers. During mitosis KSP associates with microtubules of the mitotic spindle. Microinjection of antibodies directed against KSP into human cells prevents spindle pole separation during prometaphase, giving rise to monopolar  
10 spindles and causing mitotic arrest and induction of programmed cell death. KSP and related kinesins in other, non-human, organisms, bundle antiparallel microtubules and slide them relative to one another, thus forcing the two spindle poles apart. KSP may also mediate in anaphase B spindle elongation and focussing of microtubules at the spindle pole.

15 Human KSP (also termed HsEg5) has been described [Blangy, et al., Cell, 83:1159-69 (1995); Whitehead, et al., Arthritis Rheum., 39:1635-42 (1996); Galgio et al., J. Cell Biol., 135:339-414 (1996); Blangy, et al., J Biol. Chem., 272:19418-24 (1997); Blangy, et al., Cell Motil Cytoskeleton, 40:174-82 (1998); Whitehead and Rattner, J. Cell Sci., 111:2551-61 (1998); Kaiser, et al., JBC  
20 274:18925-31 (1999); GenBank accession numbers: X85137, NM004523 and U37426] , and a fragment of the KSP gene (TRIP5) has been described [Lee, et al., Mol Endocrinol., 9:243-54 (1995); GenBank accession number L40372]. Xenopus KSP homologs (Eg5), as well as Drosophila K-LP61 F/KRP 130 have been reported.


Certain quinazolinones have recently been described as being  
25 inhibitors of KSP (PCT Publ. WO 01/30768, May 3, 2001).

Mitotic kinesins are attractive targets for the discovery and development of novel mitotic chemotherapeutics. Accordingly, it is an object of the present invention to provide compounds, methods and compositions useful in the inhibition of KSP, a mitotic kinesin.

30


#### SUMMARY OF THE INVENTION

The present invention relates to dihydropyrrole derivatives, that are useful for treating cellular proliferative diseases, for treating disorders associated with KSP kinesin activity, and for inhibiting KSP kinesin. The compounds of the  
35 invention may be illustrated by the Formula I:



### DETAILED DESCRIPTION OF THE INVENTION

The compounds of this invention are useful in the inhibition of mitotic  
 5 kinesins and are illustrated by a compound of Formula I:



or a pharmaceutically acceptable salt or stereoisomer thereof, wherein  
 10

- a is 0 or 1;
- b is 0 or 1;
- m is 0, 1, or 2;
- n is 0 or 1;
- 15 r is 0 or 1;
- s is 0 or 1;
- u is 2, 3, 4 or 5;

a dashed line represents an optional double bond, provided that one and only one  
 20 double bond is present in the ring;



R<sup>1</sup> is selected from:

- 1) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 2) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)aryl,
  - 3) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>2</sub>-C<sub>10</sub> alkenyl,
  - 5 4) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>2</sub>-C<sub>10</sub> alkynyl,
  - 5) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
  - 6) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)heterocyclyl,
  - 7) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)NR<sup>c</sup>R<sup>c'</sup>,
  - 8) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>,
  - 10 9) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkenyl,
  - 11) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkynyl,
  - 12) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-aryl,
  - 13) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-heterocyclyl,
  - 15 14) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
  - 15) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>P(=O)R<sup>d</sup>R<sup>d'</sup>,
  - 16) aryl;
  - 17) heterocyclyl; and
  - 18) C<sub>1</sub>-C<sub>10</sub> alkyl;
- 20 said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, alkylene, heteroaryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>2</sup> and R<sup>6</sup> are independently selected from:

- 1) aryl,
- 25 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 4) heterocyclyl,

said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

30

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>8</sup>, and R<sup>9</sup> are independently selected from:

- 1) H,
- 2) C<sub>1</sub>-C<sub>10</sub> alkyl,
- 3) aryl,

- 4) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 5) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 6) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 7) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 5 8) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 9) heterocyclyl,

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>; or

- 10 R<sup>4</sup> and R<sup>5</sup>, or R<sup>8</sup> and R<sup>9</sup>, attached to the same carbon atom are combined to form -(CH<sub>2</sub>)<sub>u</sub>- wherein one of the carbon atoms is optionally replaced by a moiety selected from O, S(O)<sub>m</sub>, -N(R<sup>a</sup>)C(O)-, -N(R<sup>b</sup>)- and -N(COR<sup>a</sup>)-;

R<sup>10</sup> is independently selected from:

- 15 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 20 6) CO<sub>2</sub>H,
- 7) halo,
- 8) CN,
- 9) OH,
- 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 25 11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
- 12) S(O)<sub>m</sub>R<sup>a</sup>,
- 13) S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
- 14) oxo,
- 15) CHO,
- 30 16) (N=O)R<sup>12</sup>R<sup>13</sup>, or
- 17) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R<sup>11</sup>;

R<sup>11</sup> is selected from:

- 1)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_1\text{-C}_{10})\text{alkyl}$ ,
- 2)  $\text{O}_r(\text{C}_1\text{-C}_3)\text{perfluoroalkyl}$ ,
- 3)  $(\text{C}_0\text{-C}_6)\text{alkylene-S}(\text{O})_m\text{R}^a$ ,
- 4) oxo,
- 5) OH,
- 6) halo,
- 7) CN,
- 8)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_2\text{-C}_{10})\text{alkenyl}$ ,
- 9)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_2\text{-C}_{10})\text{alkynyl}$ ,
- 10)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_3\text{-C}_6)\text{cycloalkyl}$ ,
- 11)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-aryl}$ ,
- 12)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-heterocyclyl}$ ,
- 13)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-N}(\text{R}^b)_2$ ,
- 14)  $\text{C}(\text{O})\text{R}^a$ ,
- 15)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{R}^a$ ,
- 16)  $\text{C}(\text{O})\text{H}$ ,
- 17)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{H}$ ,
- 18)  $\text{C}(\text{O})\text{N}(\text{R}^b)_2$ ,
- 19)  $\text{S}(\text{O})_m\text{R}^a$ , and
- 20)  $\text{S}(\text{O})_2\text{N}(\text{R}^b)_2$

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $\text{R}^b$ , OH,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ , halogen,  $\text{CO}_2\text{H}$ , CN,  $\text{O}(\text{C}=\text{O})\text{C}_1\text{-C}_6\text{ alkyl}$ , oxo, and  $\text{N}(\text{R}^b)_2$ ;

25  $\text{R}^{12}$  and  $\text{R}^{13}$  are independently selected from:

- 1) H,
- 2)  $(\text{C}=\text{O})\text{O}_b\text{C}_1\text{-C}_{10}\text{ alkyl}$ ,
- 3)  $(\text{C}=\text{O})\text{O}_b\text{C}_3\text{-C}_8\text{ cycloalkyl}$ ,
- 4)  $(\text{C}=\text{O})\text{O}_b\text{aryl}$ ,
- 30) 5)  $(\text{C}=\text{O})\text{O}_b\text{heterocyclyl}$ ,
- 6)  $\text{C}_1\text{-C}_{10}\text{ alkyl}$ ,
- 7) aryl,
- 8)  $\text{C}_2\text{-C}_{10}\text{ alkenyl}$ ,
- 9)  $\text{C}_2\text{-C}_{10}\text{ alkynyl}$ ,

- 10) heterocyclyl,
  - 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
  - 12) SO<sub>2</sub>R<sup>a</sup>, and
  - 13) (C=O)NR<sup>b</sup><sub>2</sub>,
- 5 said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from R<sup>11</sup>, or

R<sup>12</sup> and R<sup>13</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and  
 10 optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from R<sup>11</sup>;

R<sup>14</sup> is independently selected from:

- 15 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 20 6) CO<sub>2</sub>H,
- 7) halo,
- 8) CN,
- 9) OH,
- 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 25 11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
- 12) S(O)<sub>m</sub>R<sup>a</sup>,
- 13) S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
- 14) oxo,
- 15) CHO,
- 30 16) (N=O)R<sup>12</sup>R<sup>13</sup>, and
- 17) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R<sup>11</sup>;

R<sup>a</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one to three substituents selected from R<sup>14</sup>;

R<sup>b</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>, optionally substituted with one to three substituents selected from R<sup>14</sup>;

R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>10</sup>, or

R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;


R<sup>d</sup> and R<sup>d'</sup> are independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy and NR<sup>b</sup><sub>2</sub>, or

R<sup>d</sup> and R<sup>d'</sup> can be taken together with the phosphorous to which they are attached to form a monocyclic heterocycle with 5-7 members the ring and optionally containing, in addition to the phosphorous, one or two additional heteroatoms selected from NR<sup>e</sup>, O and S, said monocyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

R<sup>e</sup> is selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl; and

X is selected from O, NR<sup>e</sup> and S.

In a second embodiment, the compounds of this invention are illustrated by a compound of Formula I:



or a pharmaceutically acceptable salt or stereoisomer thereof, wherein

- 5    a is    0 or 1;  
      b is    0 or 1;  
      m is    0, 1, or 2;  
      n is    0 or 1;  
      r is    0 or 1;  
 10    s is    0 or 1;  
      u is    2, 3, 4 or 5;

a dashed line represents an optional double bond, provided that one and only one double bond is present in the ring;

15

R<sup>1</sup> is selected from:

- 1) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)aryl,
- 3) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 20 4) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 6) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)heterocyclyl,
- 7) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)NR<sup>c</sup>R<sup>c'</sup>,
- 8) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>,
- 25 9) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 11) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 12) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-aryl,

- 13) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-heterocyclyl,
- 14) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 15) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>P(=O)R<sup>d</sup>R<sup>d'</sup>,
- 16) aryl;
- 5 17) heterocyclyl; and
- 18) C<sub>1</sub>-C<sub>10</sub> alkyl;

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, alkylene, heteroaryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

10 R<sup>2</sup> and R<sup>6</sup> are independently selected from:

- 1) aryl,
- 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 4) heterocyclyl,

15 said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>8</sup>, and R<sup>9</sup> are independently selected from:

- 1) H,
- 20 2) C<sub>1</sub>-C<sub>10</sub> alkyl,
- 3) aryl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 5) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 6) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 25 7) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 8) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 9) heterocyclyl,

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>; or

30

R<sup>4</sup> and R<sup>5</sup>, or R<sup>8</sup> and R<sup>9</sup>, attached to the same carbon atom are combined to form -(CH<sub>2</sub>)<sub>u</sub>- wherein one of the carbon atoms is optionally replaced by a moiety selected from O, S(O)<sub>m</sub>, -N(R<sup>a</sup>)C(O)-, -N(R<sup>b</sup>)- and -N(COR<sup>a</sup>)-;

R<sup>10</sup> is independently selected from:

- 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 5 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 6) CO<sub>2</sub>H,
- 7) halo,
- 8) CN,
- 10 9) OH,
- 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
- 12) S(O)<sub>m</sub>R<sup>a</sup>,
- 13) S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
- 15 14) oxo,
- 15) CHO,
- 16) (N=O)R<sup>12</sup>R<sup>13</sup>, and
- 17) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted  
20 with one or more substituents selected from R<sup>11</sup>;

R<sup>11</sup> is selected from:

- 1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl,
- 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl,
- 25 3) (C<sub>0</sub>-C<sub>6</sub>)alkylene-S(O)<sub>m</sub>R<sup>a</sup>,
- 4) oxo,
- 5) OH,
- 6) halo,
- 7) CN,
- 30 8) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>2</sub>-C<sub>10</sub>)alkenyl,
- 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>2</sub>-C<sub>10</sub>)alkynyl,
- 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,
- 11) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,
- 12) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-heterocyclyl,



- 13)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-N}(\text{R}^b)_2$ ,  
 14)  $\text{C}(\text{O})\text{R}^a$ ,  
 15)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{R}^a$ ,  
 16)  $\text{C}(\text{O})\text{H}$ ,  
 5 17)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{H}$ ,  
 18)  $\text{C}(\text{O})\text{N}(\text{R}^b)_2$ ,  
 19)  $\text{S}(\text{O})_m\text{R}^a$ , and  
 20)  $\text{S}(\text{O})_2\text{N}(\text{R}^b)_2$

10 said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $\text{R}^b$ , OH,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ , halogen,  $\text{CO}_2\text{H}$ , CN,  $\text{O}(\text{C}=\text{O})\text{C}_1\text{-C}_6$  alkyl, oxo, and  $\text{N}(\text{R}^b)_2$ ;

$\text{R}^{12}$  and  $\text{R}^{13}$  are independently selected from:

- 15 1) H,  
 2)  $(\text{C}=\text{O})\text{O}_b\text{C}_1\text{-C}_{10}$  alkyl,  
 3)  $(\text{C}=\text{O})\text{O}_b\text{C}_3\text{-C}_8$  cycloalkyl,  
 4)  $(\text{C}=\text{O})\text{O}_b\text{aryl}$ ,  
 5)  $(\text{C}=\text{O})\text{O}_b\text{heterocyclyl}$ ,  
 6)  $\text{C}_1\text{-C}_{10}$  alkyl,  
 20 7) aryl,  
 8)  $\text{C}_2\text{-C}_{10}$  alkenyl,  
 9)  $\text{C}_2\text{-C}_{10}$  alkynyl,  
 10) heterocyclyl,  
 11)  $\text{C}_3\text{-C}_8$  cycloalkyl,  
 25 12)  $\text{SO}_2\text{R}^a$ , and  
 13)  $(\text{C}=\text{O})\text{NR}^b_2$ ,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from  $\text{R}^{11}$ , or

- 30  $\text{R}^{12}$  and  $\text{R}^{13}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from  $\text{R}^{11}$ ;

$R^a$  is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl;

$R^b$  is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl,  
5 (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>;

$R^c$  and  $R^{c'}$  are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and  
(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected  
from R<sup>10</sup>, or

10

$R^c$  and  $R^{c'}$  can be taken together with the nitrogen to which they are attached to form  
a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally  
containing, in addition to the nitrogen, one or two additional heteroatoms selected  
from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with  
15 one, two or three substituents selected from R<sup>11</sup>;

$R^d$  and  $R^{d'}$  are independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy and  
NR<sup>b</sup><sub>2</sub>, or

20  $R^d$  and  $R^{d'}$  can be taken together with the phosphorous to which they are attached to  
form a monocyclic heterocycle with 5-7 members the ring and optionally containing,  
in addition to the phosphorous, one or two additional heteroatoms selected from NR<sup>e</sup>,  
O and S, said monocyclic heterocycle optionally substituted with one, two or three  
substituents selected from R<sup>11</sup>;


25

$R^e$  is selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl; and

X is selected from O, NR<sup>e</sup> and S.

30

In another embodiment of the instant invention, the compounds are  
illustrated by a compound of Formula I:



or a pharmaceutically acceptable salt or stereoisomer thereof, wherein

- 5    a is    0 or 1;  
      b is    0 or 1;  
      m is    0, 1, or 2;  
      n is    0 or 1;  
      r is    0 or 1;  
 10    s is    0 or 1;  
      u is    2, 3, 4 or 5;

a dashed line represents an optional double bond, provided that one and only one double bond is present in the ring;

15

R¹ is selected from:

- 1) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)aryl,
- 3) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 20 4) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 6) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)heterocyclyl,
- 7) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)NR<sup>c</sup>R<sup>c'</sup>,
- 8) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>,
- 25 9) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 11) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 12) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-aryl,

- 13) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-heterocyclyl,
- 14) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 15) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>P(=O)R<sup>d</sup>R<sup>d'</sup>,
- 16) aryl;
- 5 17) heterocyclyl; and
- 18) C<sub>1</sub>-C<sub>10</sub> alkyl;

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, alkylene, heteroaryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

10 R<sup>2</sup> and R<sup>6</sup> are independently selected from:

- 1) aryl,
- 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 4) heterocyclyl,

15 said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>8</sup>, and R<sup>9</sup> are independently selected from:

- 1) H,
- 20 2) C<sub>1</sub>-C<sub>10</sub> alkyl,
- 3) aryl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 5) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 6) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 25 7) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 8) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 9) heterocyclyl,

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>; or

30

R<sup>4</sup> and R<sup>5</sup>, or R<sup>8</sup> and R<sup>9</sup>, attached to the same carbon atom are combined to form -(CH<sub>2</sub>)<sub>u</sub>- wherein one of the carbon atoms is optionally replaced by a moiety selected from O, S(O)<sub>m</sub>, -N(R<sup>a</sup>)C(O)-, -N(R<sup>b</sup>)- and -N(COR<sup>a</sup>)-;

R<sup>10</sup> is independently selected from:

- 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 5 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 6) CO<sub>2</sub>H,
- 7) halo,
- 8) CN,
- 10 9) OH,
- 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
- 12) S(O)<sub>m</sub>R<sup>a</sup>,
- 13) S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
- 15 14) oxo,
- 15) CHO,
- 16) (N=O)R<sup>12</sup>R<sup>13</sup>, and
- 17) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,

20 said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R<sup>11</sup>;

R<sup>11</sup> is selected from:

- 1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl,
- 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl,
- 25 3) (C<sub>0</sub>-C<sub>6</sub>)alkylene-S(O)<sub>m</sub>R<sup>a</sup>,
- 4) oxo,
- 5) OH,
- 6) halo,
- 7) CN,
- 30 8) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>2</sub>-C<sub>10</sub>)alkenyl,
- 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>2</sub>-C<sub>10</sub>)alkynyl,
- 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,
- 11) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,
- 12) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-heterocyclyl,

- 13)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-N(R}^b)_2$ ,
- 14)  $\text{C(O)R}^a$ ,
- 15)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{R}^a$ ,
- 16)  $\text{C(O)H}$ ,
- 5 17)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{H}$ ,
- 18)  $\text{C(O)N(R}^b)_2$ ,
- 19)  $\text{S(O)}_m\text{R}^a$ , and
- 20)  $\text{S(O)}_2\text{N(R}^b)_2$

10 said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $\text{R}^b$ , OH,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ , halogen,  $\text{CO}_2\text{H}$ , CN,  $\text{O}(\text{C}=\text{O})\text{C}_1\text{-C}_6$  alkyl, oxo, and  $\text{N(R}^b)_2$ ;

$\text{R}^{12}$  and  $\text{R}^{13}$  are independently selected from:

- 1) H,
- 15 2)  $(\text{C}=\text{O})\text{O}_b\text{C}_1\text{-C}_{10}$  alkyl,
- 3)  $(\text{C}=\text{O})\text{O}_b\text{C}_3\text{-C}_8$  cycloalkyl,
- 4)  $(\text{C}=\text{O})\text{O}_b\text{aryl}$ ,
- 5)  $(\text{C}=\text{O})\text{O}_b\text{heterocyclyl}$ ,
- 6)  $\text{C}_1\text{-C}_{10}$  alkyl,
- 20 7) aryl,
- 8)  $\text{C}_2\text{-C}_{10}$  alkenyl,
- 9)  $\text{C}_2\text{-C}_{10}$  alkynyl,
- 10) heterocyclyl,
- 11)  $\text{C}_3\text{-C}_8$  cycloalkyl,
- 25 12)  $\text{SO}_2\text{R}^a$ , and
- 13)  $(\text{C}=\text{O})\text{NR}^b_2$ ,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from  $\text{R}^{11}$ , or

30  $\text{R}^{12}$  and  $\text{R}^{13}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from  $\text{R}^{11}$ ;

$R^a$  is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl;

$R^b$  is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub> $R^a$ ;

5

$R^c$  and  $R^{c'}$  are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl or

10  $R^c$  and  $R^{c'}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^{11}$ ;

15  $R^d$  and  $R^{d'}$  are independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy and  $NR^{b2}$ , or


20  $R^d$  and  $R^{d'}$  can be taken together with the phosphorous to which they are attached to form a monocyclic heterocycle with 5-7 members the ring and optionally containing, in addition to the phosphorous, one or two additional heteroatoms selected from  $NR^e$ , O and S, said monocyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^{11}$ ;

$R^e$  is selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl; and

25

X is selected from O,  $NR^e$  and S.

A further embodiment of the present invention is illustrated by a compound of Formula II:



30

or a pharmaceutically acceptable salt or stereoisomer thereof,  
wherein:

- 5    a is    0 or 1;  
     b is    0 or 1;  
     m is    0, 1, or 2;  
     n is    0 or 1;  
     r is    0 or 1;  
10    s is    0 or 1;

a dashed line represents an optional double bond, provided that one and only one double bond is present in the ring;

15    R<sup>1</sup> is selected from:

- 1) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=O)C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=O)aryl,
- 3) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=O)C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 4) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=O)C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 20    5) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=O)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 6) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=O)heterocyclyl,
- 7) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=O)NR<sup>c</sup>R<sup>c'</sup>,
- 8) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>,
- 9) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 25    10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-aryl,
- 11) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-heterocyclyl,
- 12) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 13) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>P(=O)R<sup>d</sup>R<sup>d'</sup>,
- 14) aryl;
- 30    15) heterocyclyl; and
- 16) C<sub>1</sub>-C<sub>10</sub> alkyl;

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, alkylene, heteroaryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

35    R<sup>2</sup> and R<sup>6</sup> are independently selected from:



- 1) aryl,
- 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 4) heterocyclyl,

5 said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>3</sup>, R<sup>4</sup> and R<sup>8</sup> are independently selected from:

- 1) H,
- 10 2) C<sub>1</sub>-C<sub>10</sub> alkyl,
- 3) aryl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 5) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 6) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 15 7) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 8) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 9) heterocyclyl,

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

20

R<sup>10</sup> is independently selected from:

- 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 25 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 6) CO<sub>2</sub>H,
- 7) halo,
- 8) CN,
- 30 9) OH,
- 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
- 12) S(O)<sub>m</sub>R<sup>a</sup>,
- 13) S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
- 35 14) oxo,

- 15) CHO,
- 16) (N=O)R<sup>12</sup>R<sup>13</sup>, and
- 17) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted

5 with one, two or three substituents selected from R<sup>11</sup>;

R<sup>11</sup> is selected from:

- 1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl,
- 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl,
- 10 3) oxo,
- 4) OH,
- 5) halo,
- 6) CN,
- 7) (C<sub>2</sub>-C<sub>10</sub>)alkenyl,
- 15 8) (C<sub>2</sub>-C<sub>10</sub>)alkynyl,
- 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,
- 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,
- 11) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-heterocyclyl,
- 12) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-N(R<sup>b</sup>)<sub>2</sub>,
- 20 13) C(O)R<sup>a</sup>,
- 14) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>R<sup>a</sup>,
- 15) C(O)H,
- 16) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>H,
- 17) C(O)N(R<sup>b</sup>)<sub>2</sub>,
- 25 18) S(O)<sub>m</sub>R<sup>a</sup>, and
- 19) S(O)<sub>2</sub>N(R<sup>b</sup>)<sub>2</sub>;

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from R<sup>b</sup>, OH, (C<sub>1</sub>-C<sub>6</sub>)alkoxy,

30 halogen, CO<sub>2</sub>H, CN, O(C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, oxo, and N(R<sup>b</sup>)<sub>2</sub>;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

- 1) H,
- 2) (C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,

- 3) (C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
 4) (C=O)O<sub>b</sub>aryl,  
 5) (C=O)O<sub>b</sub>heterocyclyl,  
 6) C<sub>1</sub>-C<sub>10</sub> alkyl,  
 5 7) aryl,  
 8) C<sub>2</sub>-C<sub>10</sub> alkenyl,  
 9) C<sub>2</sub>-C<sub>10</sub> alkynyl,  
 10) heterocyclyl,  
 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
 10 12) SO<sub>2</sub>R<sup>a</sup>, and  
 13) (C=O)NR<sup>b</sup><sub>2</sub>,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R<sup>11</sup>, or

- 15 R<sup>12</sup> and R<sup>13</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

20 R<sup>a</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl;

R<sup>b</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>;


25 R<sup>c</sup> and R<sup>c'</sup> are independently selected from:  
 H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or

- 30 R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

$R^d$  and  $R^{d'}$  are independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy and  $NR^{b2}$ , or

- $R^d$  and  $R^{d'}$  can be taken together with the phosphorous to which they are attached to form a monocyclic heterocycle with 5-7 members the ring and optionally containing, in addition to the phosphorous, one or two additional heteroatoms selected from N $R^e$ , O and S, said monocyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^{11}$ ; and
- 10  $R^e$  is selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

A further embodiment of the present invention is illustrated by a compound of Formula III:



- 15 or a pharmaceutically acceptable salt or stereoisomer thereof, wherein

- a is 0 or 1;  
 b is 0 or 1;  
 20 m is 0, 1, or 2;  
 r is 0 or 1;  
 s is 0 or 1;

- a dashed line represents an optional double bond, provided that one and only one  
 25 double bond is present in the ring;

$R^1$  is selected from:

- 1) (C=O)C<sub>1</sub>-C<sub>10</sub> alkyl,

- 2) (C=O)aryl,
- 3) (C=O)C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 4) (C=O)C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C=O)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 5 6) (C=O)heterocyclyl,
- 7) (C=O)NR<sup>c</sup>R<sup>c'</sup>,
- 8) SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>,
- 9) SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 10) SO<sub>2</sub>-aryl,
- 10 11) SO<sub>2</sub>-heterocyclyl,
- 12) SO<sub>2</sub>-C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 13) P(=O)R<sup>d</sup>R<sup>d'</sup>,

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, heteroaryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

15

R<sup>2</sup> is selected from:

- 1) aryl,
- 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 20 4) heterocyclyl,

said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>3</sup>, R<sup>4</sup> and R<sup>8</sup> are independently selected from:

- 25 1) H,
- 2) C<sub>1</sub>-C<sub>10</sub> alkyl,
- 3) aryl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 5) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 30 6) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 7) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 8) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 9) heterocyclyl,

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

35

R<sup>10</sup> is independently selected from:

- 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
  - 5 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
  - 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
  - 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
  - 6) CO<sub>2</sub>H,
  - 7) halo,
  - 10 8) CN,
  - 9) OH,
  - 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
  - 11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
  - 12) S(O)<sub>m</sub>R<sup>a</sup>,
  - 15 13) S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
  - 14) oxo,
  - 15) CHO,
  - 16) (N=O)R<sup>12</sup>R<sup>13</sup>, and
  - 17) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 20 said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

R<sup>10'</sup> is halogen;

25 R<sup>11</sup> is selected from:

- 1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl,
- 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl,
- 3) oxo,
- 4) OH,
- 30 5) halo,
- 6) CN,
- 7) (C<sub>2</sub>-C<sub>10</sub>)alkenyl,
- 8) (C<sub>2</sub>-C<sub>10</sub>)alkynyl,
- 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,
- 35 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,

- 11)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-heterocyclyl}$ ,
  - 12)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-N}(\text{R}^b)_2$ ,
  - 13)  $\text{C}(\text{O})\text{R}^a$ ,
  - 14)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{R}^a$ ,
  - 5 15)  $\text{C}(\text{O})\text{H}$ ,
  - 16)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{H}$ ,
  - 17)  $\text{C}(\text{O})\text{N}(\text{R}^b)_2$ ,
  - 18)  $\text{S}(\text{O})_m\text{R}^a$ , and
  - 19)  $\text{S}(\text{O})_2\text{N}(\text{R}^b)_2$ ;
- 10 said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heterocyclyl is optionally substituted with up to three substituents selected from  $\text{R}^b$ , OH,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ , halogen,  $\text{CO}_2\text{H}$ , CN,  $\text{O}(\text{C}=\text{O})\text{C}_1\text{-C}_6$  alkyl, oxo, and  $\text{N}(\text{R}^b)_2$ ;

$\text{R}^{12}$  and  $\text{R}^{13}$  are independently selected from:

- 15 1) H,
- 2)  $(\text{C}=\text{O})\text{O}_b\text{C}_1\text{-C}_{10}$  alkyl,
- 3)  $(\text{C}=\text{O})\text{O}_b\text{C}_3\text{-C}_8$  cycloalkyl,
- 4)  $(\text{C}=\text{O})\text{O}_b\text{aryl}$ ,
- 5)  $(\text{C}=\text{O})\text{O}_b\text{heterocyclyl}$ ,
- 20 6)  $\text{C}_1\text{-C}_{10}$  alkyl,
- 7) aryl,
- 8)  $\text{C}_2\text{-C}_{10}$  alkenyl,
- 9)  $\text{C}_2\text{-C}_{10}$  alkynyl,
- 10) heterocyclyl,
- 25 11)  $\text{C}_3\text{-C}_8$  cycloalkyl,
- 12)  $\text{SO}_2\text{R}^a$ , and
- 13)  $(\text{C}=\text{O})\text{NR}^b_2$ ,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from  $\text{R}^{11}$ , or

30

$\text{R}^{12}$  and  $\text{R}^{13}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms

selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

R<sup>a</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl;

5

R<sup>b</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>;

R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and  
10 (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or

R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected  
15 from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

R<sup>d</sup> and R<sup>d'</sup> are independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy and NR<sup>b</sup><sub>2</sub>, or


20

R<sup>d</sup> and R<sup>d'</sup> can be taken together with the phosphorous to which they are attached to form a monocyclic heterocycle with 5-7 members the ring and optionally containing, in addition to the phosphorous, one or two additional heteroatoms selected from NR<sup>e</sup>, O and S, said monocyclic heterocycle optionally substituted with one, two or three  
25 substituents selected from R<sup>11</sup>; and

R<sup>e</sup> is selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

A further embodiment of the present invention is illustrated by a  
30 compound of Formula IV:





or a pharmaceutically acceptable salt or stereoisomer thereof, wherein

- 5    a is    0 or 1;  
      b is    0 or 1;  
      m is    0, 1, or 2;  
      r is    0 or 1;  
      s is    0 or 1;

10

R<sup>1</sup> is selected from:

- 1)    (C=O)C<sub>1</sub>-C<sub>10</sub> alkyl,  
      2)    (C=O)aryl,  
      3)    (C=O)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
      4)    (C=O)heterocyclyl,  
      5)    (C=O)NR<sup>c</sup>R<sup>c'</sup>,  
      6)    (C=S)NR<sup>c</sup>R<sup>c'</sup>,  
      7)    SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>,  
      8)    SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,  
      9)    SO<sub>2</sub>-aryl, and  
      10)   SO<sub>2</sub>-heterocyclyl,

15

20

said alkyl, aryl, cycloalkyl, and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>; or

25    R<sup>2</sup> is selected from:

- 1)    aryl,  
      2)    C<sub>1</sub>-C<sub>6</sub> aralkyl,  
      3)    C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and  
      4)    heterocyclyl,

said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>3</sup>, R<sup>4</sup> and R<sup>8</sup> are independently selected from:

- 5           1)     H,
- 2)     C<sub>1</sub>-C<sub>10</sub> alkyl, and
- 3)     C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

said alkyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

10   R<sup>6</sup> is selected from:

- 1)     aryl,
- 2)     C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 3)     C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 4)     heterocyclyl,

15   said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>10</sup> is independently selected from:

- 1)     (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 20       2)     (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3)     C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 4)     C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5)     (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 6)     CO<sub>2</sub>H,
- 25       7)     halo,
- 8)     CN,
- 9)     OH,
- 10)    O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 11)    O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
- 30       12)    S(O)<sub>m</sub>R<sup>a</sup>,
- 13)    S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
- 14)    oxo,
- 15)    CHO,
- 16)    (N=O)R<sup>12</sup>R<sup>13</sup>, and
- 35       17)    (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

R<sup>11</sup> is selected from:

- 5           1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl,
- 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl,
- 3) oxo,
- 4) OH,
- 5) halo,
- 10          6) CN,
- 7) (C<sub>2</sub>-C<sub>10</sub>)alkenyl,
- 8) (C<sub>2</sub>-C<sub>10</sub>)alkynyl,
- 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,
- 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,
- 15          11) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-heterocyclyl,
- 12) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-N(R<sup>b</sup>)<sub>2</sub>,
- 13) C(O)R<sup>a</sup>,
- 14) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>R<sup>a</sup>,
- 15) C(O)H,
- 20          16) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>H,
- 17) C(O)N(R<sup>b</sup>)<sub>2</sub>,
- 18) S(O)<sub>m</sub>R<sup>a</sup>, and
- 19) S(O)<sub>2</sub>N(R<sup>b</sup>)<sub>2</sub>;

25          said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heterocyclyl is optionally substituted with up to three substituents selected from R<sup>b</sup>, OH, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halogen, CO<sub>2</sub>H, CN, O(C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, oxo, and N(R<sup>b</sup>)<sub>2</sub>;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

- 30          1) H,
- 2) (C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 3) (C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 4) (C=O)O<sub>b</sub>aryl,
- 5) (C=O)O<sub>b</sub>heterocyclyl,
- 6) C<sub>1</sub>-C<sub>10</sub> alkyl,

- 7) aryl,  
8) C<sub>2</sub>-C<sub>10</sub> alkenyl,  
9) C<sub>2</sub>-C<sub>10</sub> alkynyl,  
10) heterocyclyl,  
5 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
12) SO<sub>2</sub>R<sup>a</sup>, and  
13) (C=O)NR<sup>b</sup><sub>2</sub>,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R<sup>11</sup>, or

10

R<sup>12</sup> and R<sup>13</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

15

R<sup>a</sup> is independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, and heterocyclyl;

20

R<sup>b</sup> is independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>; and

R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl or

25

R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>.

30

Another embodiment is the compound of the Formula IV described immediately above, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein:

R<sup>1</sup> is selected from:

- 1) (C=O)NR<sup>c</sup>R<sup>c'</sup>,
- 2) SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>, and
- 5 3) SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,

said alkyl, is optionally substituted with one, two or three substituents selected from R<sup>10</sup>;

R<sup>2</sup> is selected from:

- 10 1) aryl, and
- 2) heteroaryl,

said aryl and heteroaryl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

15 R<sup>3</sup>, R<sup>4</sup> and R<sup>8</sup> are independently selected from:

- 1) H, and
- 2) C<sub>1</sub>-C<sub>10</sub> alkyl,

said alkyl is optionally substituted with one or more substituents selected from R<sup>10</sup>; and

20.

R<sup>6</sup> is selected from:

- 1) aryl, and
- 2) heterocyclyl,

said alkyl, aryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>; and

25

R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>a</sup>, R<sup>b</sup>, R<sup>c</sup> and R<sup>c'</sup> are as described immediately above.


In another embodiment of the compounds of Formula IV hereinabove,

30 R<sup>2</sup> and R<sup>6</sup> are independently selected from phenyl or pyridyl, optionally substituted with one or two substituents selected from R<sup>10</sup>.

Another embodiment is the compound of the Formula IV described immediately above, or a pharmaceutically acceptable salt or stereoisomer thereof, wherein R<sup>2</sup> is phenyl, optionally substituted with one or two substituents selected from R<sup>10</sup>.

35

A further embodiment of the present invention is illustrated by a compound of Formula V, or a pharmaceutically acceptable salt or stereoisomer;



5

wherein:

a is 0 or 1;

b is 0 or 1;

m is 0, 1, or 2;

10 

r is 0 or 1;

s is 0 or 1;

R<sup>1</sup> is selected from:

- 1) (C=O)C<sub>1</sub>-C<sub>10</sub> alkyl,
- 15 2) (C=O)aryl,
- 3) (C=O)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 4) (C=O)heterocyclyl,
- 5) (C=O)NR<sup>c</sup>R<sup>c'</sup>,
- 6) (C=S)NR<sup>c</sup>R<sup>c'</sup>,
- 20 7) SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>,
- 8) SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 9) SO<sub>2</sub>-aryl, and
- 10) SO<sub>2</sub>-heterocyclyl,

said alkyl, aryl, cycloalkyl, and heterocyclyl is optionally substituted with one or more  
25 substituents selected from R<sup>10</sup>;

R<sup>2</sup> is selected from:

- 1) aryl,
- 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,

- 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 4) heterocyclyl,

said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

5 R<sup>3</sup>, R<sup>4</sup> and R<sup>8</sup> are independently selected from:

- 1) H,
- 2) C<sub>1</sub>-C<sub>10</sub> alkyl, and
- 3) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

said alkyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

10

R<sup>10</sup> is independently selected from:

- 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 15 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 6) CO<sub>2</sub>H,
- 7) halo,
- 8) CN,
- 20 9) OH,
- 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
- 12) S(O)<sub>m</sub>R<sup>a</sup>,
- 13) S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
- 25 14) oxo,
- 15) CHO,
- 16) (N=O)R<sup>12</sup>R<sup>13</sup>, and
- 17) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,

30 said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

R<sup>10'</sup> is halogen;

R<sup>11</sup> is selected from:

- 1)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_1\text{-C}_{10})\text{alkyl}$ ,
  - 2)  $\text{O}_r(\text{C}_1\text{-C}_3)\text{perfluoroalkyl}$ ,
  - 3) oxo,
  - 4) OH,
  - 5) halo,
  - 6) CN,
  - 7)  $(\text{C}_2\text{-C}_{10})\text{alkenyl}$ ,
  - 8)  $(\text{C}_2\text{-C}_{10})\text{alkynyl}$ ,
  - 9)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_3\text{-C}_6)\text{cycloalkyl}$ ,
  - 10)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-aryl}$ ,
  - 11)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-heterocyclyl}$ ,
  - 12)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-N(R}^b)_2$ ,
  - 13)  $\text{C(O)R}^a$ ,
  - 14)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{R}^a$ ,
  - 15)  $\text{C(O)H}$ ,
  - 16)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{H}$ , and
  - 17)  $\text{C(O)N(R}^b)_2$ ,
  - 18)  $\text{S(O)}_m\text{R}^a$ , and
  - 19)  $\text{S(O)}_2\text{N(R}^b)_2$ ;
- 20 said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $\text{R}^b$ , OH,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ , halogen,  $\text{CO}_2\text{H}$ , CN,  $\text{O}(\text{C}=\text{O})\text{C}_1\text{-C}_6$  alkyl, oxo, and  $\text{N(R}^b)_2$ ;

$\text{R}^{12}$  and  $\text{R}^{13}$  are independently selected from:

- 25) 1) H,
- 2)  $(\text{C}=\text{O})\text{O}_b\text{C}_1\text{-C}_{10}$  alkyl,
- 3)  $(\text{C}=\text{O})\text{O}_b\text{C}_3\text{-C}_8$  cycloalkyl,
- 4)  $(\text{C}=\text{O})\text{O}_b\text{aryl}$ ,
- 5)  $(\text{C}=\text{O})\text{O}_b\text{heterocyclyl}$ ,
- 30) 6)  $\text{C}_1\text{-C}_{10}$  alkyl,
- 7) aryl,
- 8)  $\text{C}_2\text{-C}_{10}$  alkenyl,
- 9)  $\text{C}_2\text{-C}_{10}$  alkynyl,
- 10) heterocyclyl,



- 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 12) SO<sub>2</sub>R<sup>a</sup>, and
- 13) (C=O)NR<sup>b</sup><sub>2</sub>,

5 said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R<sup>11</sup>, or

10 R<sup>12</sup> and R<sup>13</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

15 R<sup>a</sup> is independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, and heterocyclyl;

R<sup>b</sup> is independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>; and

20 R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl or

25 R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>.

Specific examples of the compounds of the instant invention include:

30 4-(2-chloro-5-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

(+)-4-(2,5-difluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

- (-)-4-(2,5-difluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 4-(5-chloro-2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 4-(2-fluoro-5-methylphenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(5-bromo-2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 4-{{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}morpholine};
- 4-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}morpholine};
- 20 N,N-dimethyl-2,4-diphenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 3-[2-fluoro-5-(trifluoromethyl)phenyl]-N,N-dimethyl-5-phenyl-2,3-dihydro-1H-pyrrole-1-carboxamide;
- 25 2-(3-fluorophenyl)-4-(2,5-difluorophenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(4-fluorophenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 4-(2,5-difluorophenyl)-2-(2-fluorophenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 35

- 2-(3-bromophenyl)-4-(2,5-difluorophenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 2-(3-aminophenyl)-4-(2,5-difluorophenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-methylphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 (2S)-4-(5-chloro-2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-1-(methylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 15 4-(2,5-difluorophenyl)-1-(ethylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-2-phenyl-1-(propylsulfonyl)-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-1-(isopropylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 20 4-(5-chloro-2-fluorophenyl)-1-(methylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-(5-chloro-2-fluorophenyl)-1-(isopropylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 25 4-(2-fluoro-5-methylphenyl)-1-(isopropylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 2- {[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]sulfonyl} ethanamine;
- 30 2- {[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]sulfonyl}-N,N-dimethylethanamine;
- 1-acetyl-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 35 4-(2-chloro-5-fluorophenyl)-1-pivaloyl-2-phenyl-2,5-dihydro-1H-pyrrole;

- 4-(2-chloro-5-fluorophenyl)-1-isobutyryl-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-1-(2,2-dimethylpropanoyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 5 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-ol;
- 1-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-ol;
- 10 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine;
- 15 4-(2-fluoro-5-isocyanophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-phenyl-1-(trifluoroacetyl)-2,5-dihydro-1H-pyrrole;
- 20 4-(5-chloro-2-fluorophenyl)-2-phenyl-1-(trifluoroacetyl)-2,5-dihydro-1H-pyrrole;
- (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylpropylamine};
- 25 (1R)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylpropylamine};
- 4-(2,5-difluorophenyl)-2-phenyl-1-L-prolyl-2,5-dihydro-1H-pyrrole;
- 30 4-(2,5-difluorophenyl)-2-phenyl-1-D-prolyl-2,5-dihydro-1H-pyrrole;
- (4R)-4-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-1,3-thiazolidine};

- methyl (3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutanoate;
- (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanamide;
- (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-(methylthio)propylamine};
- 10 (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-(methylsulfonyl)propylamine};
- (2S)-2-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}piperidine};
- 15 (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}pentylamine};
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(thien-2-ylmethyl)ethylamine;
- 20 4-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-1,1-dioxidotetrahydro-2H-thiopyran-4-ylamine};
- (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methyl-1-oxopropan-2-amine;
- (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}propylamine};
- 30 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-phenylethanamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-phenylethanamine;
- 35

- (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanamide;
- 5 3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-amine;
- (1S,2S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylbutylamine;
- 10 (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}butylamine;
- (1S)-1-cyclopropyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 15 1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopropanamine;
- 20 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-oxopropan-2-amine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-methyl-2-oxoethylamine;
- 25 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(pyridin-2-ylmethyl)ethylamine;
- (1S)-1-cyclohexyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 30 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(4-iodobenzyl)-2-oxoethylamine;

- (1S)-1-benzyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine;
- 4-{(2S)-2-amino-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropyl}phenol;
- (3S)-3-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-1,2,3,4-tetrahydroisoquinoline;
- (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-phenylpropylamine;
- (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-methylbutylamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(pyridin-3-ylmethyl)ethylamine;
- 1-[(2S)-azetidin-2-ylcarbonyl]-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- (3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutanamide;
- 4-(2,5-difluorophenyl)-1-[(2S)-2,5-dihydro-1H-pyrrol-2-ylcarbonyl]-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-1-[(2-methylazetidin-2-yl)carbonyl]-2-phenyl-2,5-dihydro-1H-pyrrole;
- (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylamine;
- methyl (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanoate;

- 4-(2,5-difluorophenyl)-2-phenyl-1-[[ (2S,3S)-2-phenylpyrrolidin-3-yl]carbonyl]-2,5-dihydro-1H-pyrrole;
- 5 4-(2,5-difluorophenyl)-2-phenyl-1-[(5-phenylpyrrolidin-3-yl)carbonyl]-2,5-dihydro-1H-pyrrole;
- (2S)-2-amino-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-ol;
- 10 (2R,3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutan-2-ol;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-
- 15 (methoxymethyl)-2-oxoethylamine;
- 4-(2,5-difluorophenyl)-2-phenyl-1-(pyrrolidin-3-ylcarbonyl)-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-2-phenyl-1-[(3-phenylpyrrolidin-3-yl)acetyl]-2,5-dihydro-1H-
- 20 pyrrole;
- (1S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-3,3-difluoropropylamine;
- 25 (1S)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxo-1-phenylpropan-1-amine;
- 4-(2,5-difluorophenyl)-2-phenyl-1-[(4S)-4-phenyl-L-prolyl]-2,5-dihydro-1H-pyrrole;
- 30 1-[2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]cyclohexanamine;
- 2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;



- 4-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}piperidin-4-amine;
- (1S,3R)-3-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopentanamine;
- 5 (1R,4S)-4-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-2-en-1-amine;
- 10 (1S,4R)-4-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-2-en-1-amine;
- (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-ynylamine;
- 15 (1R)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxo-1-phenylpropan-1-amine;
- 3-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-phenylpiperidine;
- 20 (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-enylamine;
- 25 (2S)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-(methylamino)-3-oxopropan-1-ol;
- (3R,5S)-5-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}pyrrolidin-3-ol;
- 30 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(1,3-thiazol-4-ylmethyl)ethylamine;
- (1R)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-enylamine;
- 35

- (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,3-dimethyl-1-oxobutan-2-amine;
- 5 (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,4-dimethyl-1-oxopentan-2-amine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-[(1-methyl-1H-imidazol-4-yl)methyl]-2-oxoethylamine;
- 10 4-(2,5-difluorophenyl)-1-(N-6--formyl-L-lysyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- (2S,3S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,3-dimethyl-1-oxopentan-2-amine;
- 15 (1S)-1-(cyclohexylmethyl)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxoethylamine;
- 20 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(isocyanomethyl)-2-oxoethylamine;
- (1S)-1-1-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-3,3-dimethylbutylamine;
- 25 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,3-dimethyl-1-oxobutan-2-amine;
- 30 1-1-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]cyclohexanamine;
- 1-1-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]cyclopentanamine;
- 35

- (1S)-3-(benzyloxy)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}propylamine;
- 5 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,3-dimethyl-1-oxobutan-2-amine;
- 1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-3-en-1-amine;
- 10 (1S)-1-cyclopentyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 4-(2,5-difluorophenyl)-1-(2-methylpropyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 15 1-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine;
- (1S)-1-{{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylpropylamine;
- 20 (1S)-2-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-cyclopropyl-2-oxoethanamine;
- 25 (1S,2S)-1-{{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylbutylamine;
- (1S)-1-{{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}pentylamine;
- 30 (1S)-1-{{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3,3-dimethylbutylamine;
- (1S)-1-{{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylamine;
- 35

- (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylpropylamine;
- 5 (1S)-1-cyclohexyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-enylamine;
- 10 (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-ynylamine;
- (1S)-1-benzyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine;
- 15 (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 20 1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine;
- (1S)-1-{[(2S)-4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylamine;
- 25 (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}pentylamine;
- (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-methylbutylamine;
- 30 (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3,3-dimethylbutylamine;

- 1-cyclopropyl-3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-amine;
- (1S)-2-[(2S)-4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-cyclopropyl-2-oxoethanamine;
- 5 (1S)-1-[(2S)-4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl-2-methylpropylamine;
- 10 (1S,2S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl-2-methylbutylamine;
- 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole;
- 15 (2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole;
- 20 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole;
- (2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole;
- 25 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole;
- 3-1-[(2S)-2-amino-2-cyclopropylethanoyl]-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 30 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-isoleucyl-2,5-dihydro-1H-pyrrole;

- 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-norleucyl-2,5-dihydro-1H-pyrrole;
- 5 (2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(3-methyl-L-valyl)-2,5-dihydro-1H-pyrrole;
- (2S)-4-(2,5-Difluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 (2S)-4-(2,5-Difluorophenyl)-N-methyl-2-phenyl-N-(piperidin-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(5-Chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-[(3R)-pyrrolidin-3-yl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 (2S)-4-(5-chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(5-chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-[(3S)-pyrrolidin-3-yl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 (2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-pyrrolidin-3-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(2,5-difluorophenyl)-N-methyl-N-[(3S)-1-methylpyrrolidin-3-yl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 (2S)-4-(2,5-difluorophenyl)-N-methyl-N-[(3R)-1-methylpyrrolidin-3-yl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 (2S)-4-(2,5-difluorophenyl)-N-methyl-N-[(1-methyl-5-oxopyrrolidin-2-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(2,5-difluorophenyl)-N-(1,3-dioxolan-2-ylmethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 35

- (2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-tetrahydrofuran-3-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 (2S)-N-(1-allylpiperidin-4-yl)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- allyl 4-[[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl] (methyl)amino]piperidine-1-carboxylate;
- 10 allyl 4-[[[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl] carbonyl](methyl)amino]methyl]piperidine-1-carboxylate;
- (2S)-4-(2,5-difluorophenyl)-N-methyl-N-(1-methylpiperidin-4-yl)-2-phenyl-2,5-
- 15 dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-Difluorophenyl)-N-methyl-N-[(1-methylpiperidin-3-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(pyridin-3-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- N-benzyl-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 4-(2,5-difluorophenyl)-N-methyl-N-[(1-methyl-1H-pyrazol-4-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-[2-(dimethylamino)ethyl]-N-methyl-2-phenyl-2,5-dihydro-
- 30 1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-(2-hydroxyethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

- 4-(2,5-difluorophenyl)-N-isobutyl-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(2-pyridin-2-ylethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-(2-methoxyethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 4-(2,5-difluorophenyl)-N-(2,3-dihydroxypropyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(2-phenylethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 4-(2,5-difluorophenyl)-2-phenyl-N-propyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 1-Acetyl-4-(2,5-difluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole;
- 25 (2S)-1-[4-(2,5-difluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methyl-1-oxobutan-2-amine;
- (2S)-4-(2,5-difluorophenyl)-N,N,2-trimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 (2S)-4-(5-chloro-2-fluorophenyl)-N,N,2-trimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;



- (2S)-4-(5-chloro-2-fluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 (2S)-4-(5-chloro-2-fluorophenyl)-N-ethyl-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3,3-dimethyl-1-oxobutan-2-ol;
- 10 (2S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methyl-1-oxobutan-2-ol;
- (2S,3S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methyl-1-oxopentan-2-ol;
- 15 1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-oxohexan-2-ol;
- (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-oxo-3-phenylpropan-2-ol;
- 20 (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-methyl-1-oxopentan-2-ol;
- 25 (1S)-1-cyclohexyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanol;
- (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3,3-dimethyl-1-oxobutan-2-ol;
- 30 N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-N-2,N-2-dimethylglycinamide;
- N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-N-2-methylglycinamide;
- 35

- N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)glycinamide;
- 5 N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-methylalaninamide;
- N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)glycinamide;
- 10 N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2,N-2-dimethylglycinamide;
- N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2,N-2-dimethylglycinamide, N-oxide;
- 15 N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-methylalaninamide;
- N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2,N-2-dimethylglycinamide n-oxide;
- 20 N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-pyrrolidin-1-ylacetamide;
- 25 2-azetidin-1-yl-N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)acetamide;
- N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-morpholin-4-ylacetamide;
- 30 N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-piperazin-1-ylacetamide;

- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(4-methylpiperazin-1-yl)acetamide;
- 2-azetidin-1-yl-N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}acetamide;
- 5 N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-pyrrolidin-1-ylacetamide;
- 10 N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-piperidin-1-ylacetamide;
- N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-morpholin-4-ylacetamide;
- 15 N-1-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N-2-(2-hydroxyethyl)glycinamide;
- N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(4-methylpiperazin-1-yl)acetamide;
- 20 N-1-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N-2-isopropylglycinamide;
- 25 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}acetamide;
- N-1-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N-2-ethylglycinamide;
- 30 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-hydroxyacetamide;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}piperazine-1-carboxamide;
- 35

N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-piperidin-4-ylurea;

- 5 4-amino-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}piperidine-1-carboxamide;

N-(2-aminoethyl)-N'-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}urea;

10

N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-(3-morpholin-4-ylpropyl)urea;

- 15 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-[2-(dimethylamino)ethyl]urea;

2-azetidin-1-yl-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}ethanesulfonamide

- 20 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(isopropylamino)ethanesulfonamide;

N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-pyrrolidin-1-ylethanesulfonamide;

25

N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-morpholin-4-ylethanesulfonamide;

- 30 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-piperazin-1-ylethanesulfonamide;

N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(4-methylpiperazin-1-yl)ethanesulfonamide;

- N-(tert-butyl)-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetamide;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-  
5 isopropylacetamide;
- (2S)-1-(2-azetidin-1-yl-2-oxoethyl)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 10 (2S)-4-(2,5-difluorophenyl)-1-(2-oxo-2-pyrrolidin-1-ylethyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl}  
morpholine;  
15 1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl}  
piperazine;
- 1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl}-4-  
20 methylpiperazine;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-  
isopropylbutanamide;
- 25 4-{2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]butanoyl}  
morpholine;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-  
ethylacetamide;  
30 N-cyclobutyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetamide;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-  
35 ethylpropanamide;

- N-cyclobutyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanamide;
- 5 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methylpropanamide;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylpropanamide;
- 10 N-(tert-butyl)-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanamide;
- 4-{2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoyl} morpholine;
- 15 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethyl-2,2-dimethyl-4-oxobutanamide;
- 20 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxo-N-piperidin-4-ylbutanamide;
- (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid;
- 25 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2,2-tetramethyl-4-oxobutanamide;
- (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethyl-3-oxo-3-piperazin-1-ylpropylamine;
- 30 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropyl-2,2-dimethyl-4-oxobutanamide;

- (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,2,2-trimethyl-4-oxobutanamide;
- (3R)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2,2-tetramethyl-4-oxobutanamide;
- (3R)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid;
- (1R)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl-2,2-dimethyl-3-oxo-3-piperazin-1-ylpropylamine;
- 2-[(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]amino)-N-ethylacetamide;
- 2-[(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]amino)-N-methylacetamide;
- 2-[(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]amino)-N,N-dimethylacetamide;
- 2-[(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]amino)-N-methyl-N-ethylacetamide;
- 2-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]amino)-N-methylacetamide;
- 2-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]amino)-N-ethylacetamide;
- 2-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]amino)-N,N-dimethylacetamide;
- 2-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]amino)-N-isopropylacetamide;

- 2-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-ethyl-N-methylacetamide;
- 5 2-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N,N-diethylacetamide;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-N-(2-oxo-2-pyrrolidin-1-ylethyl)ethanamine;
- 10 (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-(2-morpholin-4-yl-2-oxoethyl)-2-oxoethanamine;
- 1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)acetyl]piperidin-4-ol;
- 15 (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]-2-oxoethanamine;
- (1S)-N-(2-azetidin-1-yl-2-oxoethyl)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 20 (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-[2-(1,1-dioxidothiomorpholin-4-yl)-2-oxoethyl]-2-oxoethanamine;
- 25 (1S)-N-[2-(4-acetylpiperazin-1-yl)-2-oxoethyl]-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- (1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-(2-morpholin-4-yl-2-oxoethyl)-2-oxoethanamine;
- 30 (1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-N-(2-oxo-2-pyrrolidin-1-ylethyl)ethanamine;



- 2-(((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)-N-isopropylacetamide;
- 2-(dimethylamino)ethyl (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate;
- 1-methylpiperidin-4-yl (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate;
- (2S)-4-cyclopropyl-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-cyclopentyl-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 4-methylpiperazine-1-carboxylate;
- 1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 2-morpholin-4-ylethylcarbamate;
- N-(((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)oxy)-carbonyl]glycine;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 1-methylpiperidin-4-ylcarbamate;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylmethyl(1-methylpiperidin-4-yl)carbamate;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 4-dimethylamino)piperidine-1-carboxylate;
- tert-butyl (2S)-4-(2-chloro-5-fluoropyrimidin-4-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate;

- (2S)-4-(5-fluoro-2-methylpyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 (2S)-4-(2-chloro-5-fluoropyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(4-chloro-5-methylpyrimidin-2-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 (2S)-4-(6-chloropyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(2-chloropyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 (2S)-N,N-dimethyl-4-(4-methylpyridin-3-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-N,N-dimethyl-2-phenyl-4-(1,3-thiazol-2-yl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 (2S)-N,N-dimethyl-2-phenyl-4-(1,3-thiazol-4-yl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 (2S)-N,N-dimethyl-2-phenyl-4-(1,2-thiazol-5-yl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 (2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(2,5-difluorophenyl)-N-(2-hydroxyethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 35

- N-{[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-N-methyl-beta-alanine;
- 5 methyl N-{[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-N-methyl-beta-alaninate;
- 4-{[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]acetyl}morpholin-4-ium;
- 10 3-[(2S)-4-(2,5-difluorophenyl)-1-(2-hydroxy-2-methylpropanoyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-sulfonamide;
- 15 3-[4-(2,5-difluorophenyl)-1-(methylsulfonyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 3-[4-(2,5-difluorophenyl)-1-(morpholin-4-ylcarbonyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 20 3-[4-(2,5-difluorophenyl)-1-(2,2-dimethylpropanoyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 25 (2S)-4-(2,5-difluorophenyl)-1-[(methylsulfonyl)acetyl]-2-phenyl-2,5-dihydro-1H-pyrrole;
- (2S)-4-(2,5-difluorophenyl)-2-phenyl-1-[(phenylsulfonyl)acetyl]-2,5-dihydro-1H-pyrrole;
- 30 3-[(2S)-1-[(2S)-2-cyclopropyl-2-hydroxyethanoyl]-4-(2,5-difluorophenyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 35 3-{(2S)-4-(2,5-difluorophenyl)-1-[(2S)-2-hydroxy-3,3-dimethylbutanoyl]-2,5-dihydro-1H-pyrrol-2-yl}phenol;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanol;

- (2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-tetrahydrofuran-3-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 (2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(2-methoxyethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-tetrahydro-2H-pyran-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 3-[(2S)-4-(2,5-difluorophenyl)-1-(piperidin-1-ylcarbonyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 4-(2,5-difluorophenyl)-N-[1-(2-fluoroethyl)piperidin-4-yl]-2-(3-hydroxyphenyl)-N-
- 15 methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]piperidinium trifluoroacetate;
- 20 2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethyl 4-methylpiperazine-1-carboxylate;
- 3-{4-(2,5-difluorophenyl)-1-[(4-methylpiperazin-1-yl)carbonyl]-2,5-dihydro-1H-pyrrol-2-yl}phenol;
- 25 2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethyl morpholine-4-carboxylate;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(1-methyl-1H-pyrazol-4-
- 30 yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(isoxazol-5-ylmethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

- 2-[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethyl dimethylaminocarboxylate;
- 2-[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethyl piperidine-1-carboxylate;
- 5 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(5-oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(tetrahydro-2H-pyran-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-{[5-(methoxymethyl)-1H-pyrazol-3-yl]methyl}-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(4-methyl-1,2,5-oxadiazol-3-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-2-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(isoxazol-3-ylmethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-1,2,4-triazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-pyrazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 35

- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(1-methyl-5-oxopyrrolidin-2-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(1-isoxazol-3-ylethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-(1,3-dioxolan-2-ylmethyl)-2-(3-hydroxyphenyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 4-(2,5-difluorophenyl)-N-(1,4-dioxan-2-ylmethyl)-2-(3-hydroxyphenyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(5-methyl-1,3,4-oxadiazol-2-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 2-[(4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl(methylamino)ethanesulfonic acid;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(methylsulfonyl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 2-[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl(methylamino)ethanesulfonic acid];
- 2-hydroxyethyl (1S)-1-[[2S]-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2,2-dimethylpropylcarbamate;
- 25 3-hydroxypropyl (1S)-1-[[2S]-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2,2-dimethylpropylcarbamate;
- 2-hydroxyethyl {(1S)-1-isopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}carbamate;
- 30 2-hydroxyethyl {(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}carbamate;

- 4-hydroxybutyl (1*S*)-1-{[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylcarbamate;
- (2*S*)-4-(2,5-difluorophenyl)-1-[2-(methylsulfonyl)ethyl]-2-phenyl-2,5-dihydro-1*H*-pyrrole;
- (2*S*)-4-(2,5-difluorophenyl)-1-[2-(ethylsulfonyl)ethyl]-2-phenyl-2,5-dihydro-1*H*-pyrrole;
- 1-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]pentan-3-one;
- 4-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]butan-2-one;
- 4-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-3-methylbutan-2-one;
- 2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-*N,N*-dimethylethanesulfonamide;
- 3-[(2*S*)-4-(2,5-difluorophenyl)-1-[2-(methylsulfonyl)ethyl]-2,5-dihydro-1*H*-pyrrol-2-yl]phenol;
- (2*S*)-4-(2,5-difluorophenyl)-1-[2-(ethylsulfonyl)propyl]-2-phenyl-2,5-dihydro-1*H*-pyrrole;
- 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-*N*-methylpropanamide;
- 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-*N,N*-dimethylpropanamide;
- 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-*N,N*,2-trimethylpropanamide;
- 4-{3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoyl}morpholine;
- 1-{3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoyl}-4-(methylsulfonyl)piperazine;
- 1-{3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoyl}piperidin-4-ol;

- methyl 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoate;
- 2-(((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl)oxy)-*N*-ethylacetamide ;
- 4-(((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy)acetyl)morpholine;
- 2-((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy)-*N*-(2-hydroxyethyl)acetamide;
- 1-(((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy)acetyl)-4-methylpiperazine;
- 1-(((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy)acetyl)piperazine;
- 2-((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy)-*N*-piperidin-4-ylacetamide;
- 1-(((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy)acetyl)piperidin-4-amine;
- N*-((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl)-3-morpholin-4-yl-3-oxopropan-1-amine;
- N*<sup>3</sup>-((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl)-*N*<sup>1</sup>,*N*<sup>1</sup>-dimethyl-β-alaninamide;
- ((1*S*)-1-([(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]carbonyl)-2,2-dimethylpropyl)(3-morpholin-4-yl-3-oxopropyl)amine;
- or a pharmaceutically acceptable salt or stereoisomer thereof.

35

Particular examples of the compounds of the instant invention are:



(-)-4-(2,5-difluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

4-(5-chloro-2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

4-(2,5-difluorophenyl)-1-(isopropylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;

1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-ol;

1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine;

(1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylamine;

(2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;

or a pharmaceutically acceptable salt or stereoisomer thereof.

In another embodiment, particular examples of the compounds of the instant invention are:

(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;

(2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;

(1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylamine;

2-({(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-ethylacetamide;

(2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

(2S)-4-(2,5-difluorophenyl)-N-methyl-N-(1-methylpiperidin-4-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide; and

(2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole

or a pharmaceutically acceptable salt or stereoisomer thereof.

Further examples of the compounds of the instant invention are the  
TFA salts of:

2-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]sulfonyl)-N,N-dimethylethanamine;

1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine;

4-(5-chloro-2-fluorophenyl)-2-phenyl-1-(trifluoroacetyl)-2,5-dihydro-1H-pyrrole;

(1S)-1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2-methylpropylamine;

(1R)-1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2-methylpropylamine;

4-(2,5-difluorophenyl)-2-phenyl-1-L-prolyl-2,5-dihydro-1H-pyrrole;

4-(2,5-difluorophenyl)-2-phenyl-1-D-prolyl-2,5-dihydro-1H-pyrrole;

- (4R)-4-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-1,3-thiazolidine;
- 5 methyl (3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutanoate;
- (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanamide;
- 10 (1S)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-3-(methylthio)propylamine;
- (1S)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-3-(methylsulfonyl)propylamine;
- 15 (2S)-2-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}piperidine;
- (1S)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}pentylamine;
- 20 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(thien-2-ylmethyl)ethylamine;
- 25 4-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-1,1-dioxidotetrahydro-2H-thiopyran-4-ylamine;
- (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methyl-1-oxopropan-2-amine;
- 30 (1S)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}propylamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-phenylethanamine;
- 35

- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-phenylethanamine;
- 5 (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanamide
- 3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-amine;
- 10 (1S,2S)-1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2-methylbutylamine;
- (1S)-1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)
- 15 butylamine;
- (1S)-1-cyclopropyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 20 1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)cyclopropanamine;
- 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-oxopropan-2-amine;
- 25 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-methyl-2-oxoethylamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-
- 30 (pyridin-2-ylmethyl)ethylamine;
- (1S)-1-cyclohexyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;

- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(4-iodobenzyl)-2-oxoethylamine;
- 5 (1S)-1-benzyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine;
- 4-[(2S)-2-amino-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropyl]phenol;
- 10 (3S)-3-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-1,2,3,4-tetrahydroisoquinoline;
- (1S)-1-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-3-phenylpropylamine;
- 15 (1S)-1-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-3-methylbutylamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(pyridin-3-ylmethyl)ethylamine;
- 20 1-[(2S)-azetidin-2-ylcarbonyl]-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 25 (3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutanamide;
- 4-(2,5-difluorophenyl)-1-[(2-methylazetidin-2-yl)carbonyl]-2-phenyl-2,5-dihydro-1H-pyrrole;
- 30 (1S)-1-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-2,2-dimethylpropylamine;
- methyl (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanoate;
- 35

- 4-(2,5-difluorophenyl)-2-phenyl-1-([(2S,3S)-2-phenylpyrrolidin-3-yl]carbonyl)-2,5-dihydro-1H-pyrrole;
- 5 4-(2,5-difluorophenyl)-2-phenyl-1-[(5-phenylpyrrolidin-3-yl)carbonyl]-2,5-dihydro-1H-pyrrole;
- (2S)-2-amino-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-ol;
- 10 (2R,3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutan-2-ol;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(methoxymethyl)-2-oxoethylamine;
- 15 4-(2,5-difluorophenyl)-2-phenyl-1-(pyrrolidin-3-ylcarbonyl)-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-2-phenyl-1-[(3-phenylpyrrolidin-3-yl)acetyl]-2,5-dihydro-1H-pyrrole;
- 20 (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3,3-difluoropropylamine;
- (1S)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxo-1-phenylpropan-1-amine;
- 25 4-(2,5-difluorophenyl)-2-phenyl-1-[(4S)-4-phenyl-L-prolyl]-2,5-dihydro-1H-pyrrole;
- 1-{2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}cyclohexanamine;
- 30 2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;

- 4-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}piperidin-4-amine;
- (1S,3R)-3-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}  
5 cyclopentanamine;
- (1R,4S)-4-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-2-en-1-amine;
- 10 (1S,4R)-4-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-2-en-1-amine;
- (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-ynylamine;
- 15 (1R)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxo-1-phenylpropan-1-amine;
- 3-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-phenylpiperidine;
- 20 (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-enylamine;
- (2S)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-(methylamino)-3-oxopropan-1-ol;
- 25 (3R,5S)-5-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}pyrrolidin-3-ol;
- 30 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(1,3-thiazol-4-ylmethyl)ethylamine;
- (1R)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-enylamine;
- 35

- (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,3-dimethyl-1-oxobutan-2-amine;
- 5 (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,4-dimethyl-1-oxopentan-2-amine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-[(1-methyl-1H-imidazol-4-yl)methyl]-2-oxoethylamine;
- 10 4-(2,5-difluorophenyl)-1-(N-6-formyl-L-lysyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- (2S,3S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,3-dimethyl-1-oxopentan-2-amine;
- 15 (1S)-1-(cyclohexylmethyl)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxoethylamine;
- 20 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(isocyanomethyl)-2-oxoethylamine;
- (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3,3-dimethylbutylamine;
- 25 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,3-dimethyl-1-oxobutan-2-amine;
- 30 1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclohexanamine;
- 1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopentanamine;
- 35



- (1S)-3-(benzyloxy)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}propylamine;
- 5 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,3-dimethyl-1-oxobutan-2-amine;
- 1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-3-en-1-amine;
- 10 (1S)-1-cyclopentyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 4-(2,5-difluorophenyl)-1-(2-methylpropyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 15 (1S)-2-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-cyclopropyl-2-oxoethanamine;
- (1S,2S)-1-{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylbutylamine;
- 20 (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-enylamine;
- (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-ynylamine;
- 25 (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 30 1-cyclopropyl-3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-amine;
- (1S,2S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylbutylamine;
- 35

- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole;
- 3-[1-[(2S)-2-amino-2-cyclopropylethanoyl]-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-isoleucyl-2,5-dihydro-1H-pyrrole;
- (2S)-4-(2,5-Difluorophenyl)-N-methyl-2-phenyl-N-(piperidin-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(5-Chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-[(3R)-pyrrolidin-3-yl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(5-chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(5-chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-[(3S)-pyrrolidin-3-yl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-pyrrolidin-3-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-N-(1-allylpiperidin-4-yl)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-Difluorophenyl)-N-methyl-N-[(1-methylpiperidin-3-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(pyridin-3-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;

- 4-(2,5-difluorophenyl)-N-methyl-N-[(1-methyl-1H-pyrazol-4-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 4-(2,5-difluorophenyl)-N-[2-(dimethylamino)ethyl]-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(2-pyridin-2-ylethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 (2S)-1-[4-(2,5-difluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methyl-1-oxobutan-2-amine;
- 20 N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-N-2,N-2-dimethylglycinamide;
- N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-N-2-methylglycinamide;
- 25 N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]glycinamide;
- N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-2-methylalaninamide;
- 30 N-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-2-pyrrolidin-1-ylacetamide;

- 2-azetidin-1-yl-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}acetamide;
- 5 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-morpholin-4-ylacetamide;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-piperazin-1-ylacetamide;
- 10 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(4-methylpiperazin-1-yl)acetamide;
- N-1-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N-2-isopropylglycinamide;
- 15 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}piperazine-1-carboxamide;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-piperidin-4-ylurea;
- 20 4-amino-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}piperidine-1-carboxamide;
- 25 N-(2-aminoethyl)-N'-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}urea;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-(3-morpholin-4-ylpropyl)urea;
- 30 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-[2-(dimethylamino)ethyl]urea;
- 2-azetidin-1-yl-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}ethanesulfonamide
- 35

- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(isopropylamino)ethanesulfonamide;
- 5 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-pyrrolidin-1-ylethanesulfonamide;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-morpholin-4-ylethanesulfonamide;
- 10 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-piperazin-1-ylethanesulfonamide;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(4-methylpiperazin-1-yl)ethanesulfonamide;
- 15 N-(tert-butyl)-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetamide;
- 20 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylacetamide;
- (2S)-1-(2-azetidin-1-yl-2-oxoethyl)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 25 (2S)-4-(2,5-difluorophenyl)-1-(2-oxo-2-pyrrolidin-1-ylethyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl}
- 30 morpholine;
- 1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl} piperazine;

- 1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl}-4-methylpiperazine;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-  
5 isopropylbutanamide;
- 4-{2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]butanoyl}morpholine;
- 10 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethylacetamide;
- N-cyclobutyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetamide;  
15 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethylpropanamide;
- N-cyclobutyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanamide;  
20 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methylpropanamide;
- 25 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylpropanamide;
- N-(tert-butyl)-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanamide;  
30 4-{2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoyl}morpholine;
- (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-  
35 N-ethyl-2,2-dimethyl-4-oxobutanamide;

- (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxo-N-piperidin-4-ylbutanamide;
- 5 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid;
- (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2,2-tetramethyl-4-oxobutanamide;
- 10 (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethyl-3-oxo-3-piperazin-1-ylpropylamine;
- (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropyl-2,2-dimethyl-4-oxobutanamide;
- 15 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,2,2-trimethyl-4-oxobutanamide;
- (3R)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2,2-tetramethyl-4-oxobutanamide;
- 20 (3R)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid;
- 25 (1R)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethyl-3-oxo-3-piperazin-1-ylpropylamine
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 4-methylpiperazine-1-carboxylate;
- 30 (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 1-methylpiperidin-4-ylcarbamate;

- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylmethyl(1-methylpiperidin-4-yl)carbamate;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl-4-dimethylamino)piperidine-1-carboxylate;
- (2S)-N,N-dimethyl-4-(4-methylpyridin-3-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-{[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]acetyl}morpholin-4-ium;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethyl-4-methylpiperazine-1-carboxylate;
- 3-{4-(2,5-difluorophenyl)-1-[(4-methylpiperazin-1-yl)carbonyl]-2,5-dihydro-1H-pyrrol-2-yl}phenol;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(1-methyl-1H-pyrazol-4-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluoro-phenyl)-2,5-dihydro-2-(3-hydroxyphenyl)-N-methyl-N--[2-(4-methyl-1-piperazinyl)-2-oxoethyl]-H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(5-oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-{[5-(methoxymethyl)-1H-pyrazol-3-yl]methyl}-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;



- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-2-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-1,2,4-triazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-pyrazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]butan-2-one;
- 4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methylbutan-2-one;
- 15 3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methylpropanamide;
- 3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N-dimethylpropanamide;
- 20 3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2-trimethylpropanamide;
- 4-{3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoyl}morpholine;
- 25 1-{3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoyl}-4-(methylsulfonyl)piperazine;
- 30 1-{3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoyl}piperidin-4-ol; and
- methyl 3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoate.
- 35

The compounds of the present invention may have asymmetric centers, chiral axes, and chiral planes (as described in: E.L. Eliel and S.H. Wilen, *Stereochemistry of Carbon Compounds*, John Wiley & Sons, New York, 1994, pages 1119-1190), and occur as racemates, racemic mixtures, and as individual diastereomers,

40 with all possible isomers and mixtures thereof, including optical isomers, all such stereoisomers being included in the present invention. In addition, the compounds disclosed herein may exist as tautomers and both tautomeric forms are intended to be

encompassed by the scope of the invention, even though only one tautomeric structure is depicted.

When any variable (e.g. R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, etc.) occurs more than one time in any constituent, its definition on each occurrence is independent at every other  
5 occurrence. Also, combinations of substituents and variables are permissible only if such combinations result in stable compounds. Lines drawn into the ring systems from substituents represent that the indicated bond may be attached to any of the substitutable ring atoms. If the ring system is polycyclic, it is intended that the bond be attached to any of the suitable carbon atoms on the proximal ring only.

10 It is understood that substituents and substitution patterns on the compounds of the instant invention can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques known in the art, as well as those methods set forth below, from readily available starting materials. If a substituent is itself substituted with more than one  
15 group, it is understood that these multiple groups may be on the same carbon or on different carbons, so long as a stable structure results. The phrase "optionally substituted with one or more substituents" should be taken to be equivalent to the phrase "optionally substituted with at least one substituent" and in such cases the preferred embodiment will have from zero to three substituents.

20 As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, C<sub>1</sub>-C<sub>10</sub>, as in "C<sub>1</sub>-C<sub>10</sub> alkyl" is defined to include groups having 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 carbons in a linear or branched arrangement. For example, "C<sub>1</sub>-C<sub>10</sub> alkyl" specifically includes methyl, ethyl, *n*-  
25 propyl, *i*-propyl, *n*-butyl, *t*-butyl, *i*-butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, and so on. The term "cycloalkyl" means a monocyclic saturated aliphatic hydrocarbon group having the specified number of carbon atoms. For example, "cycloalkyl" includes cyclopropyl, methyl-cyclopropyl, 2,2-dimethyl-cyclobutyl, 2-ethyl-cyclopentyl, cyclohexyl, and so on. In an embodiment of the invention the term  
30 "cycloalkyl" includes the groups described immediately above and further includes monocyclic unsaturated aliphatic hydrocarbon groups. For example, "cycloalkyl" as defined in this embodiment includes cyclopropyl, methyl-cyclopropyl, 2,2-dimethyl-cyclobutyl, 2-ethyl-cyclopentyl, cyclohexyl, cyclopentenyl, cyclobutenyl and so on.

35 The term "alkylene" means a hydrocarbon diradical group having the specified number of carbon atoms. For example, "alkylene" includes -CH<sub>2</sub>-,

-CH<sub>2</sub>CH<sub>2</sub>- and the like.

When used in the phrases "C<sub>1</sub>-C<sub>6</sub> aralkyl" and "C<sub>1</sub>-C<sub>6</sub> heteroaralkyl" the term "C<sub>1</sub>-C<sub>6</sub>" refers to the alkyl portion of the moiety and does not describe the number of atoms in the aryl and heteroaryl portion of the moiety.

5 "Alkoxy" represents either a cyclic or non-cyclic alkyl group of indicated number of carbon atoms attached through an oxygen bridge. "Alkoxy" therefore encompasses the definitions of alkyl and cycloalkyl above.

If no number of carbon atoms is specified, the term "alkenyl" refers to a non-aromatic hydrocarbon radical, straight, branched or cyclic, containing from 2 to  
10 10 carbon atoms and at least one carbon to carbon double bond. Preferably one carbon to carbon double bond is present, and up to four non-aromatic carbon-carbon double bonds may be present. Thus, "C<sub>2</sub>-C<sub>6</sub> alkenyl" means an alkenyl radical having from 2 to 6 carbon atoms. Alkenyl groups include ethenyl, propenyl, butenyl, 2-methylbutenyl and cyclohexenyl. The straight, branched or cyclic portion of the  
15 alkenyl group may contain double bonds and may be substituted if a substituted alkenyl group is indicated.

The term "alkynyl" refers to a hydrocarbon radical straight, branched or cyclic, containing from 2 to 10 carbon atoms and at least one carbon to carbon triple bond. Up to three carbon-carbon triple bonds may be present. Thus, "C<sub>2</sub>-C<sub>6</sub>  
20 alkynyl" means an alkynyl radical having from 2 to 6 carbon atoms. Alkynyl groups include ethynyl, propynyl, butynyl, 3-methylbutynyl and so on. The straight, branched or cyclic portion of the alkynyl group may contain triple bonds and may be substituted if a substituted alkynyl group is indicated.

In certain instances, substituents may be defined with a range of  
25 carbons that includes zero, such as (C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl. If aryl is taken to be phenyl, this definition would include phenyl itself as well as -CH<sub>2</sub>Ph, -CH<sub>2</sub>CH<sub>2</sub>Ph, CH(CH<sub>3</sub>)CH<sub>2</sub>CH(CH<sub>3</sub>)Ph, and so on.

As used herein, "aryl" is intended to mean any stable monocyclic or bicyclic carbon ring of up to 7 atoms in each ring, wherein at least one ring is  
30 aromatic. Examples of such aryl elements include phenyl, naphthyl, tetrahydronaphthyl, indanyl and biphenyl. In cases where the aryl substituent is bicyclic and one ring is non-aromatic, it is understood that attachment is via the aromatic ring.

The term heteroaryl, as used herein, represents a stable monocyclic or bicyclic ring of up to 7 atoms in each ring, wherein at least one ring is aromatic and contains from 1 to 4 heteroatoms selected from the group consisting of O, N and S. Heteroaryl groups within the scope of this definition include but are not limited to:

- 5 acridinyl, carbazolyl, cinnolinyl, quinoxaliny, pyrazolyl, indolyl, benzotriazolyl, furanyl, thienyl, benzothienyl, benzofuranyl, quinolinyl, isoquinolinyl, oxazolyl, isoxazolyl, indolyl, pyrazinyl, pyridazinyl, pyridinyl, pyrimidinyl, pyrrolyl, tetrahydroquinoline. As with the definition of heterocycle below, "heteroaryl" is also understood to include the N-oxide derivative of any nitrogen-containing heteroaryl.
- 10 In cases where the heteroaryl substituent is bicyclic and one ring is non-aromatic or contains no heteroatoms, it is understood that attachment is via the aromatic ring or via the heteroatom containing ring, respectively.

- The term "heterocycle" or "heterocyclyl" as used herein is intended to mean a 3- to 10-membered aromatic or nonaromatic heterocycle containing from 1 to 4 heteroatoms selected from the group consisting of O, N and S, and includes bicyclic groups. "Heterocyclyl" therefore includes the above mentioned heteroaryls, as well as dihydro and tetrahydro analogs thereof. Further examples of "heterocyclyl" include, but are not limited to the following: azetidiny, benzoimidazolyl, benzofuranyl, benzofurazanyl, benzopyrazolyl, benzotriazolyl, benzothiophenyl, benzoxazolyl, carbazolyl, carbolinyl, cinnolinyl, furanyl, imidazolyl, indolinyl, indolyl, indolaziny, indazolyl, isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthpyridinyl, oxadiazolyl, oxazolyl, oxazoline, isoxazoline, oxetanyl, pyranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridopyridinyl, pyridazinyl, pyridyl, pyrimidyl, pyrrolyl, quinazolinyl, quinolyl, quinoxaliny, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydroisoquinolinyl, tetrazolyl, tetrazolopyridyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, 1,4-dioxanyl, hexahydroazepinyl, piperazinyl, piperidinyl, pyridin-2-onyl, pyrrolidinyl, morpholinyl, thiomorpholinyl, dihydrobenzoimidazolyl, dihydrobenzofuranyl, dihydrobenzothiophenyl, dihydrobenzoxazolyl, dihydrofuranyl, dihydroimidazolyl, dihydroindolyl, dihydroisooxazolyl, dihydroisothiazolyl, dihydrooxadiazolyl, dihydrooxazolyl, dihydropyrazinyl, dihydropyrazolyl, dihydropyridinyl, dihydropyrimidinyl, dihydropyrrolyl, dihydroquinolinyl, dihydrotetrazolyl, dihydrothiadiazolyl, dihydrothiazolyl, dihydrothienyl, dihydrotriazolyl, dihydroazetidiny, methylenedioxybenzoyl, tetrahydrofuranyl, and tetrahydrothienyl, and N-oxides

thereof. Attachment of a heterocyclyl substituent can occur via a carbon atom or via a heteroatom.

In an embodiment, the term "heterocycle" or "heterocyclyl" as used herein is intended to mean a 5- to 10-membered aromatic or nonaromatic heterocycle containing from 1 to 4 heteroatoms selected from the group consisting of O, N and S, and includes bicyclic groups. "Heterocyclyl" in this embodiment therefore includes the above mentioned heteroaryls, as well as dihydro and tetrathydro analogs thereof. Further examples of "heterocyclyl" include, but are not limited to the following: benzoimidazolyl, benzofuranyl, benzofurazanyl, benzopyrazolyl, benzotriazolyl, benzothiophenyl, benzoxazolyl, carbazolyl, carbolinyl, cinnolinyl, furanyl, imidazolyl, indolinyl, indolyl, indolaziny, indazolyl, isobenzofuranyl, isoindolyl, isoquinolyl, isothiazolyl, isoxazolyl, naphthpyridinyl, oxadiazolyl, oxazolyl, oxazoline, isoxazoline, oxetanyl, pyranyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridopyridinyl, pyridazinyl, pyridyl, pyrimidyl, pyrrolyl, quinazolinyl, quinolyl, quinoxalinyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydroisoquinolyl, tetrazolyl, tetrazolopyridyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, azetidyl, 1,4-dioxanyl, hexahydroazepinyl, piperazinyl, piperidinyl, pyridin-2-onyl, pyrrolidinyl, morpholinyl, thiomorpholinyl, dihydrobenzoimidazolyl, dihydrobenzofuranyl, dihydrobenzothiophenyl, dihydrobenzoxazolyl, dihydrofuranyl, dihydroimidazolyl, dihydroindolyl, dihydroisooxazolyl, dihydroisothiazolyl, dihydrooxadiazolyl, dihydrooxazolyl, dihydropyrazinyl, dihydropyrazolyl, dihydropyridinyl, dihydropyrimidinyl, dihydropyrrolyl, dihydroquinolyl, dihydrotetrazolyl, dihydrothiadiazolyl, dihydrothiazolyl, dihydrothienyl, dihydrotriazolyl, dihydroazetidyl, methylenedioxybenzoyl, tetrahydrofuranyl, and tetrahydrothienyl, and N-oxides thereof. Attachment of a heterocyclyl substituent can occur via a carbon atom or via a heteroatom.

In another embodiment, heterocycle is selected from 2-azepinone, benzimidazolyl, 2-diazapinone, imidazolyl, 2-imidazolidinone, indolyl, isoquinolyl, morpholinyl, piperidyl, piperazinyl, pyridyl, pyrrolidinyl, 2-piperidinone, 2-pyrimidinone, 2-pyrrolidinone, quinolyl, tetrahydrofuryl, tetrahydroisoquinolyl, and thienyl.

As appreciated by those of skill in the art, "halo" or "halogen" as used herein is intended to include chloro, fluoro, bromo and iodo.

The alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heteroaryl and heterocyclyl substituents may be substituted or unsubstituted, unless specifically


- defined otherwise. For example, a (C<sub>1</sub>-C<sub>6</sub>)alkyl may be substituted with one, two or three substituents selected from OH, oxo, halogen, alkoxy, dialkylamino, or heterocyclyl, such as morpholinyl, piperidinyl, and so on. In this case, if one substituent is oxo and the other is OH, the following are included in the definition:
- 5 -C(=O)CH<sub>2</sub>CH(OH)CH<sub>3</sub>, -(C=O)OH, -CH<sub>2</sub>(OH)CH<sub>2</sub>CH(O), and so on.

The moiety formed when, in the definition of R<sup>4</sup> and R<sup>5</sup> and R<sup>8</sup> and R<sup>9</sup> on the same carbon atom are combined to form -(CH<sub>2</sub>)<sub>4</sub>- is illustrated by the following:



10


In addition, such cyclic moieties may optionally include a heteroatom(s). Examples of such heteroatom-containing cyclic moieties include, but are not limited to:




15

- In certain instances, R<sup>12</sup> and R<sup>13</sup> and R<sup>C</sup> and R<sup>C'</sup> are defined such that they can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said heterocycle optionally substituted with one or more substituents selected from R<sup>11</sup>. Examples of the heterocycles that can thus be formed include, but are not limited to the following, keeping in mind that the heterocycle is optionally
- 20

substituted with one or more (and preferably one, two or three) substituents chosen from R<sup>11</sup>:



- 5 In certain instances, R<sup>d</sup> and R<sup>d'</sup> are defined such that they can be taken together with the phosphorous to which they are attached to form a monocyclic heterocycle with 5-7 members in the ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N R<sup>e</sup>, O and S, said heterocycle optionally substituted with one or more substituents selected from R<sup>11</sup>.
- 10 Examples of the heterocycles that can thus be formed include, but are not limited to the following, keeping in mind that the heterocycle is optionally substituted with one or more (and preferably one or two) substituents chosen from R<sup>11</sup>:



In an embodiment, R<sup>1</sup> is selected from (C=O)NR<sup>c</sup>R<sup>c'</sup> and -SO<sub>2</sub>C<sub>1</sub>-C<sub>6</sub> alkyl, optionally substituted with one to three substituents selected from R<sup>10</sup>. In another aspect of this embodiment, R<sup>1</sup> is aminocarbonyl, N,N-dimethylaminocarbonyl, methylsulfonyl or ethylsulfonyl.

- In another embodiment, R<sup>1</sup> is selected from (C=O)NR<sup>c</sup>R<sup>c'</sup> and (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, optionally substituted with one to three substituents selected from R<sup>10</sup>. In another aspect of this embodiment, R<sup>1</sup> is selected from:
- 10 a) 1-cyclopropyl-2-oxoethanamine or 1-t-butyl-2-oxoethanamine, wherein the amine nitrogen is optionally substituted with R<sup>12</sup> and R<sup>13</sup>,
  - b) N,N-dimethylcarboxamide, and
  - c) N-methyl-N-(1-methylpiperidin-4-yl)carboxamide.

In an embodiment, R<sup>2</sup> is selected from aryl, optionally substituted with one to three substituents selected from R<sup>10</sup>. In another aspect of this embodiment, R<sup>2</sup> is phenyl, optionally substituted with one to three substituents selected from halo.

In another embodiment, R<sup>2</sup> is selected from aryl, optionally substituted with one to three substituents selected from R<sup>10</sup>. In another aspect of this embodiment, R<sup>2</sup> is phenyl and 3-hydrophenyl, optionally substituted with one to three substituents selected from halo.

In an embodiment, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are H.



In an embodiment, R<sup>3</sup> is selected from H and C<sub>1</sub>-C<sub>6</sub> alkyl, optionally substituted with one to two substituents selected from R<sup>10</sup>. In another aspect of this embodiment, R<sup>3</sup> is selected from H and methyl.

5 In an embodiment, R<sup>6</sup> is selected from aryl, optionally substituted with one to three substituents selected from R<sup>10</sup>. More preferably, R<sup>6</sup> is phenyl, optionally substituted with one to three substituents selected from halo.

Included in the instant invention is the free form of compounds of Formula I, as well as the pharmaceutically acceptable salts and stereoisomers thereof. Some of the specific compounds exemplified herein are the protonated salts of amine  
10 compounds. The term "free form" refers to the amine compounds in non-salt form. The encompassed pharmaceutically acceptable salts not only include the salts exemplified for the specific compounds described herein, but also all the typical pharmaceutically acceptable salts of the free form of compounds of Formula I. The free form of the specific salt compounds described may be isolated using techniques  
15 known in the art. For example, the free form may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous NaOH, potassium carbonate, ammonia and sodium bicarbonate. The free forms may differ from their respective salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the acid and base salts are otherwise pharmaceutically equivalent  
20 to their respective free forms for purposes of the invention.

The pharmaceutically acceptable salts of the instant compounds can be synthesized from the compounds of this invention which contain a basic or acidic moiety by conventional chemical methods. Generally, the salts of the basic  
25 compounds are prepared either by ion exchange chromatography or by reacting the free base with stoichiometric amounts or with an excess of the desired salt-forming inorganic or organic acid in a suitable solvent or various combinations of solvents. Similarly, the salts of the acidic compounds are formed by reactions with the appropriate inorganic or organic base.

Thus, pharmaceutically acceptable salts of the compounds of this  
30 invention include the conventional non-toxic salts of the compounds of this invention as formed by reacting a basic instant compound with an inorganic or organic acid. For example, conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like, as well as salts prepared from organic acids such as acetic, propionic, succinic,

glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, trifluoroacetic and the like.

5                   When the compound of the present invention is acidic, suitable “pharmaceutically acceptable salts” refers to salts prepared from pharmaceutically acceptable non-toxic bases including inorganic bases and organic bases. Salts derived from inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic salts, manganous, potassium, sodium, zinc and the  
10 like. Particularly preferred are the ammonium, calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as arginine, betaine caffeine, choline, N,N<sup>1</sup>-dibenzylethylenediamine,  
15 diethylamin, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine tripropylamine, tromethamine and the like. When the  
20 compound of the present invention is acidic, the term “free form” refers to the compound in its non-salt form, such that the acidic functionality is still protonated.

The preparation of the pharmaceutically acceptable salts described above and other typical pharmaceutically acceptable salts is more fully described by Berg *et al.*, “Pharmaceutical Salts,” *J. Pharm. Sci.*, 1977:66:1-19.

25                   It will also be noted that the compounds of the present invention may potentially be internal salts or zwitterions, since under physiological conditions a deprotonated acidic moiety in the compound, such as a carboxyl group, may be anionic, and this electronic charge might then be balanced off internally against the cationic charge of a protonated or alkylated basic moiety, such as a quaternary  
30 nitrogen atom. An isolated compound having internally balance charges, and thus not associated with a intermolecular counterion, may also be considered the “free form” of a compound.

The compounds of this invention may be prepared by employing reactions as shown in the following schemes, in addition to other standard  
35 manipulations that are known in the literature or exemplified in the experimental

procedures. The illustrative schemes below, therefore, are not limited by the compounds listed or by any particular substituents employed for illustrative purposes. Substituent numbering as shown in the schemes does not necessarily correlate to that used in the claims and often, for clarity, a single substituent is shown attached to the compound where multiple substituents are allowed under the definitions of Formula I  
5 hereinabove.

### SCHEMES

10 As shown in Scheme A, key dihydropyrrole intermediate A-4 may be obtained from readily available suitably substituted anilines and N-protected dihydropyrrole. Subsequent deprotection of the ring nitrogen allows functionalization with appropriately substituted electrophiles, such as the dialkylcarbamoyl chloride illustrated to give the instant compound A-6. Scheme A-1 shows an  
15 alternative synthesis of the A-4 intermediate.

As shown in Scheme B, use of a single phenyl pyrrole enantiomer having a suitably located hydroxyl moiety allows for the preparation of the enantiomerically pure intermediate B-5, which may then be deprotected and functionalized in a method analogous to that shown in Scheme A.

20 Scheme C illustrates attachment of a suitably substituted amino acid residue to the intermediate A-5. The group R<sup>sc</sup> represents a substituent sidechain, preferably a sidechain of an amino acid.

As illustrated in Schemes D and F, attachment of an amino carbonyl moiety to the dihydropyrrole ring nitrogen can also be accomplished by a two step  
25 synthesis, thereby allowing incorporation of a wide variety of substituted amines into the instant compounds.

Scheme E illustrates the syntheses of compounds of the instant invention that incorporate both a phenyl moiety and an alkyl moiety on the 2-position of the dihydropyrrole ring.


30 As illustrated in Scheme G, suitably substituted hydroxyalkylamides may also be prepared.


Other R<sup>6</sup> substituents may be incorporated into the instant compound as shown in Scheme H. Thus a suitable Grignard reagent may replace the aryl boronic acid that is utilized in Scheme B.

A Suzuki coupling may also be employed to incorporate an R<sup>6</sup> heteroaryl substituent, as illustrated in Scheme I.


SCHEME A

5





SCHEME A (continued)

10


SCHEME A-1

## SCHEME B




SCHEME C


5


SCHEME D


## SCHEME E







SCHEME E (continued)

SCHEME F

SCHEME G

5

SCHEME H

SCHEME I5 Utilities

The compounds of the invention find use in a variety of applications. As will be appreciated by those in the art, mitosis may be altered in a variety of ways; that is, one can affect mitosis either by increasing or decreasing the activity of a

10 component in the mitotic pathway. Stated differently, mitosis may be affected (e.g.,

disrupted) by disturbing equilibrium, either by inhibiting or activating certain components. Similar approaches may be used to alter meiosis.

5 In a preferred embodiment, the compounds of the invention are used to modulate mitotic spindle formation, thus causing prolonged cell cycle arrest in mitosis. By "modulate" herein is meant altering mitotic spindle formation, including increasing and decreasing spindle formation. By "mitotic spindle formation" herein is meant organization of microtubules into bipolar structures by mitotic kinesins. By "mitotic spindle dysfunction" herein is meant mitotic arrest and monopolar spindle formation.

10 The compounds of the invention are useful to bind to and/or modulate the activity of a mitotic kinesin. In a preferred embodiment, the mitotic kinesin is a member of the bimC subfamily of mitotic kinesins (as described in U.S. Patent No. 6,284,480, column 5). In a further preferred embodiment, the mitotic kinesin is human KSP, although the activity of mitotic kinesins from other organisms may also  
15 be modulated by the compounds of the present invention. In this context, modulate means either increasing or decreasing spindle pole separation, causing malformation, i.e., splaying, of mitotic spindle poles, or otherwise causing morphological perturbation of the mitotic spindle. Also included within the definition of KSP for these purposes are variants and/or fragments of KSP. See PCT Publ. WO 01/31335:  
20 "Methods of Screening for Modulators of Cell Proliferation and Methods of Diagnosing Cell Proliferation States", filed Oct. 27, 1999, hereby incorporated by reference in its entirety. In addition, other mitotic kinesins may be inhibited by the compounds of the present invention.

The compounds of the invention are used to treat cellular proliferation  
25 diseases. Disease states which can be treated by the methods and compositions provided herein include, but are not limited to, cancer (further discussed below), autoimmune disease, arthritis, graft rejection, inflammatory bowel disease, proliferation induced after medical procedures, including, but not limited to, surgery, angioplasty, and the like. It is appreciated that in some cases the cells may not be in a  
30 hyper- or hypoproliferation state (abnormal state) and still require treatment. For example, during wound healing, the cells may be proliferating "normally", but proliferation enhancement may be desired. Similarly, as discussed above, in the agriculture arena, cells may be in a "normal" state, but proliferation modulation may be desired to enhance a crop by directly enhancing growth of a crop, or by inhibiting  
35 the growth of a plant or organism which adversely affects the crop. Thus, in one

embodiment, the invention herein includes application to cells or individuals afflicted or impending affliction with any one of these disorders or states.

The compounds, compositions and methods provided herein are particularly deemed useful for the treatment of cancer including solid tumors such as skin, breast, brain, cervical carcinomas, testicular carcinomas, etc. More particularly, cancers that may be treated by the compounds, compositions and methods of the invention include, but are not limited to: Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma); Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocyoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondrogenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical

dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma),  
5 vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma];  
10 Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal glands: neuroblastoma. Thus, the term "cancerous cell" as provided herein, includes a cell afflicted by any one of the above-identified conditions.

The compounds of the instant invention may also be useful as  
15 antifungal agents, by modulating the activity of the fungal members of the bimC kinesin subgroup, as is described in U.S. Patent No. 6,284,480.

The compounds of this invention may be administered to mammals, preferably humans, either alone or, preferably, in combination with pharmaceutically acceptable carriers, excipients or diluents, in a pharmaceutical composition, according  
20 to standard pharmaceutical practice. The compounds can be administered orally or parenterally, including the intravenous, intramuscular, intraperitoneal, subcutaneous, rectal and topical routes of administration.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specific amounts, as well as any  
25 product which results, directly or indirectly, from combination of the specific ingredients in the specified amounts.

The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules,  
30 or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the  
35 active ingredient in admixture with non-toxic pharmaceutically acceptable excipients

which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, microcrystalline cellulose, sodium crosscarmellose, corn starch, or alginic acid; binding agents, for example starch, gelatin, polyvinyl-pyrrolidone or acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to mask the unpleasant taste of the drug or delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a water soluble taste masking material such as hydroxypropyl-methylcellulose or hydroxypropylcellulose, or a time delay material such as ethyl cellulose, cellulose acetate butyrate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water soluble carrier such as polyethyleneglycol or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active material in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl-cellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.



Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as butylated hydroxyanisol or alpha-tocopherol.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

The pharmaceutical compositions of the invention may also be in the form of an oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally occurring phosphatides, for example soy bean lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening, flavoring agents, preservatives and antioxidants.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, flavoring and coloring agents and antioxidant.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous solutions. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.

The sterile injectable preparation may also be a sterile injectable oil-in-water microemulsion where the active ingredient is dissolved in the oily phase. For example, the active ingredient may be first dissolved in a mixture of soybean oil and lecithin. The oil solution then introduced into a water and glycerol mixture and processed to form a microemulation.

The injectable solutions or microemulsions may be introduced into a patient's blood stream by local bolus injection. Alternatively, it may be advantageous to administer the solution or microemulsion in such a way as to maintain a constant circulating concentration of the instant compound. In order to maintain such a  
5 constant concentration, a continuous intravenous delivery device may be utilized. An example of such a device is the Deltec CADD-PLUS™ model 5400 intravenous pump.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension for intramuscular and subcutaneous  
10 administration. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butane diol. In addition, sterile, fixed oils are  
15 conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Compounds of Formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be  
20 prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

25 For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compound of Formula I are employed. (For purposes of this application, topical application shall include mouth washes and gargles.)

The compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles and delivery devices, or  
30 via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen. Compounds of the present invention may also be delivered as a suppository employing bases such as cocoa butter,

glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters of polyethylene glycol.

When a compound according to this invention is administered into a human subject, the daily dosage will normally be determined by the prescribing  
5 physician with the dosage generally varying according to the age, weight, sex and response of the individual patient, as well as the severity of the patient's symptoms.

In one exemplary application, a suitable amount of compound is administered to a mammal undergoing treatment for cancer. Administration occurs in an amount between about 0.1 mg/kg of body weight to about 60 mg/kg of body  
10 weight per day, preferably of between 0.5 mg/kg of body weight to about 40 mg/kg of body weight per day.

The instant compounds are also useful in combination with known therapeutic agents and anti-cancer agents. For example, instant compounds are useful in combination with known anti-cancer agents. Combinations of the presently  
15 disclosed compounds with other anti-cancer or chemotherapeutic agents are within the scope of the invention. Examples of such agents can be found in *Cancer Principles and Practice of Oncology* by V.T. Devita and S. Hellman (editors), 6<sup>th</sup> edition (February 15, 2001), Lippincott Williams & Wilkins Publishers. A person of ordinary skill in the art would be able to discern which combinations of agents would be useful  
20 based on the particular characteristics of the drugs and the cancer involved. Such anti-cancer agents include, but are not limited to, the following: estrogen receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic/cytostatic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors and other angiogenesis inhibitors,  
25 inhibitors of cell proliferation and survival signaling, and agents that interfere with cell cycle checkpoints. The instant compounds are particularly useful when co-administered with radiation therapy.

In an embodiment, the instant compounds are also useful in combination with known anti-cancer agents including the following: estrogen  
30 receptor modulators, androgen receptor modulators, retinoid receptor modulators, cytotoxic agents, antiproliferative agents, prenyl-protein transferase inhibitors, HMG-CoA reductase inhibitors, HIV protease inhibitors, reverse transcriptase inhibitors, and other angiogenesis inhibitors.

"Estrogen receptor modulators" refers to compounds that interfere  
35 with or inhibit the binding of estrogen to the receptor, regardless of mechanism.

Examples of estrogen receptor modulators include, but are not limited to, tamoxifen, raloxifene, idoxifene, LY353381, LY117081, toremifene, fulvestrant, 4-[7-(2,2-dimethyl-1-oxopropoxy-4-methyl-2-[4-[2-(1-piperidinyl)ethoxy]phenyl]-2H-1-benzopyran-3-yl)-phenyl-2,2-dimethylpropanoate, 4,4'-dihydroxybenzophenone-2,4-dinitrophenyl-hydrazone, and SH646.

"Androgen receptor modulators" refers to compounds which interfere or inhibit the binding of androgens to the receptor, regardless of mechanism.

Examples of androgen receptor modulators include finasteride and other 5 $\alpha$ -reductase inhibitors, nilutamide, flutamide, bicalutamide, liarozole, and abiraterone acetate.

"Retinoid receptor modulators" refers to compounds which interfere or inhibit the binding of retinoids to the receptor, regardless of mechanism. Examples of such retinoid receptor modulators include bexarotene, tretinoin, 13-cis-retinoic acid, 9-cis-retinoic acid,  $\alpha$ -difluoromethylornithine, ILX23-7553, trans-N-(4'-hydroxyphenyl) retinamide, and N-4-carboxyphenyl retinamide.

"Cytotoxic/cytostatic agents" refer to compounds which cause cell death or inhibit cell proliferation primarily by interfering directly with the cell's functioning or inhibit or interfere with cell myosis, including alkylating agents, tumor necrosis factors, intercalators, hypoxia activatable compounds, microtubule inhibitors/microtubule-stabilizing agents, inhibitors of mitotic kinesins, inhibitors of kinases involved in mitotic progression, antimetabolites; biological response modifiers; hormonal/anti-hormonal therapeutic agents, haematopoietic growth factors, monoclonal antibody targeted therapeutic agents, topoisomerase inhibitors, proteasome inhibitors and ubiquitin ligase inhibitors.

Examples of cytotoxic agents include, but are not limited to, sertenef, cachectin, ifosfamide, tasonermin, lonidamine, carboplatin, altretamine, prednimustine, dibromodulcitol, ranimustine, fotemustine, nedaplatin, oxaliplatin, temozolomide, heptaplatin, estramustine, improsulfan tosilate, trofosfamide, nimustine, dibrospidium chloride, pumitepa, lobaplatin, satraplatin, profiromycin, cisplatin, irofulven, dexifosfamide, cis-aminedichloro(2-methyl-pyridine)platinum, benzylguanine, glufosfamide, GPX100, (trans, trans, trans)-bis-mu-(hexane-1,6-diamine)-mu-[diamine-platinum(II)]bis[diamine(chloro)platinum (II)]tetrachloride, diarizidinylspermine, arsenic trioxide, 1-(11-dodecylamino-10-hydroxyundecyl)-3,7-dimethylxanthine, zorubicin, idarubicin, daunorubicin, bisantrene, mitoxantrone, pirarubicin, pinafide, valrubicin, amrubicin, antineoplaston, 3'-deamino-3'-morpholino-13-deoxo-10-hydroxycarminomycin, annamycin, galarubicin, elinafide,

MEN10755, and 4-demethoxy-3-deamino-3-aziridinyl-4-methylsulphonyl-daunorubicin (see WO 00/50032).

An example of a hypoxia activatable compound is tirapazamine.

Examples of proteasome inhibitors include but are not limited to  
5 lactacystin and MLN-341 (Velcade).

Examples of microtubule inhibitors/microtubule-stabilising agents include paclitaxel, vindesine sulfate, 3',4'-didehydro-4'-deoxy-8'-norvincal leukoblastine, docetaxol, rhizoxin, dolastatin, mivobulin isethionate, auristatin, cemadotin, RPR109881, BMS184476, vinflunine, cryptophycin, 2,3,4,5,6-  
10 pentafluoro-N-(3-fluoro-4-methoxyphenyl) benzene sulfonamide, anhydrovinblastine, N,N-dimethyl-L-valyl-L-valyl-N-methyl-L-valyl-L-prolyl-L-proline-t-butylamide, TDX258, the epothilones (see for example U.S. Pat. Nos. 6,284,781 and 6,288,237) and BMS188797. In an embodiment the epothilones are not included in the microtubule inhibitors/microtubule-stabilising agents.

Some examples of topoisomerase inhibitors are topotecan, hycaptamine, irinotecan, rubitecan, 6-ethoxypropionyl-3',4'-O-exo-benzylidene-chartreusin, 9-methoxy-N,N-dimethyl-5-nitropyrazolo[3,4,5-kl]acridine-2-(6H)propanamine, 1-amino-9-ethyl-5-fluoro-2,3-dihydro-9-hydroxy-4-methyl-1H,12H-benzo[de]pyrano[3',4':b,7]-indolizino[1,2b]quinoline-10,13(9H,15H)dione,  
20 lurtotecan, 7-[2-(N-isopropylamino)ethyl]-(20S)camptothecin, BNP1350, BNPI1100, BN80915, BN80942, etoposide phosphate, teniposide, sobuzoxane, 2'-dimethylamino-2'-deoxy-etoposide, GL331, N-[2-(dimethylamino)ethyl]-9-hydroxy-5,6-dimethyl-6H-pyrido[4,3-b]carbazole-1-carboxamide, asulacrine, (5a, 5aB, 8aa,9b)-9-[2-[N-[2-(dimethylamino)ethyl]-N-methylamino]ethyl]-5-[4-hydroxy-3,5-  
25 dimethoxyphenyl]-5,5a,6,8,8a,9-hexahydrofuro(3',4':6,7)naphtho(2,3-d)-1,3-dioxol-6-one, 2,3-(methylenedioxy)-5-methyl-7-hydroxy-8-methoxybenzo[c]-phenanthridinium, 6,9-bis[(2-aminoethyl)amino]benzo[g]isoguinoline-5,10-dione, 5-(3-aminopropylamino)-7,10-dihydroxy-2-(2-hydroxyethylaminomethyl)-6H-pyrazolo[4,5,1-de]acridin-6-one, N-[1-[2(diethylamino)ethylamino]-7-methoxy-9-  
30 oxo-9H-thioxanthen-4-ylmethyl]formamide, N-(2-(dimethylamino)ethyl)acridine-4-carboxamide, 6-[[2-(dimethylamino)ethyl]amino]-3-hydroxy-7H-indeno[2,1-c]quinolin-7-one, and dimesna.

Examples of inhibitors of mitotic kinesins, and in particular the human mitotic kinesin KSP, are described in PCT Publications WO 01/30768 and

WO 01/98278, and pending U.S. Ser. Nos. 60/338,779 (filed December 6, 2001), 60/338,344 (filed December 6, 2001), 60/338,383 (filed December 6, 2001), 60/338,380 (filed December 6, 2001), 60/338,379 (filed December 6, 2001) and 60/344,453 (filed November 7, 2001). In an embodiment inhibitors of mitotic  
 5 kinesins include, but are not limited to inhibitors of KSP, inhibitors of MKLP1, inhibitors of CENP-E, inhibitors of MCAK and inhibitors of Rab6-KIFL.


“Inhibitors of kinases involved in mitotic progression” include, but are not limited to, inhibitors of aurora kinase, inhibitors of Polo-like kinases (PLK) (in particular inhibitors of PLK-1), inhibitors of bub-1 and inhibitors of bub-R1.

10 “Antiproliferative agents” includes antisense RNA and DNA oligonucleotides such as G3139, ODN698, RVASKRAS, GEM231, and INX3001, and antimetabolites such as enocitabine, carmofur, tegafur, pentostatin, doxifluridine, trimetrexate, fludarabine, capecitabine, galocitabine, cytarabine ocfosfate, fosteabine sodium hydrate, raltitrexed, paltitrexid, emitefur, tiazofurin, decitabine, nolatrexed,  
 15 pemetrexed, nelzarabine, 2'-deoxy-2'-methylidenecytidine, 2'-fluoromethylene-2'-deoxycytidine, N-[5-(2,3-dihydro-benzofuryl)sulfonyl]-N'-(3,4-dichlorophenyl)urea, N6-[4-deoxy-4-[N2-[2(E),4(E)-tetradecadienoyl]glycylamino]-L-glycero-B-L-manno-heptopyranosyl]adenine, aplidine, ecteinascidin, troxacitabine, 4-[2-amino-4-oxo-4,6,7,8-tetrahydro-3H-pyrimidino[5,4-b][1,4]thiazin-6-yl-(S)-ethyl]-2,5-thienoyl-L-  
 20 glutamic acid, aminopterin, 5-fluorouracil, alanosine, 11-acetyl-8-(carbamoyloxymethyl)-4-formyl-6-methoxy-14-oxa-1,11-diazatetracyclo(7.4.1.0.0)-tetradeca-2,4,6-trien-9-yl acetic acid ester, swainsonine, lometrexol, dexrazoxane, methioninase, 2'-cyano-2'-deoxy-N4-palmitoyl-1-B-D-arabino furanosyl cytosine, 3-aminopyridine-2-carboxaldehyde thiosemicarbazone and trastuzumab.

25 Examples of monoclonal antibody targeted therapeutic agents include those therapeutic agents which have cytotoxic agents or radioisotopes attached to a cancer cell specific or target cell specific monoclonal antibody. Examples include Bexxar.

30 “HMG-CoA reductase inhibitors” refers to inhibitors of 3-hydroxy-3-methylglutaryl-CoA reductase. Compounds which have inhibitory activity for HMG-CoA reductase can be readily identified by using assays well-known in the art. For example, see the assays described or cited in U.S. Patent 4,231,938 at col. 6, and WO 84/02131 at pp. 30-33. The terms “HMG-CoA reductase inhibitor” and “inhibitor of HMG-CoA reductase” have the same meaning when used herein.

Examples of HMG-CoA reductase inhibitors that may be used include but are not limited to lovastatin (MEVACOR®; see U.S. Patent Nos. 4,231,938, 4,294,926 and 4,319,039), simvastatin (ZOCOR®; see U.S. Patent Nos. 4,444,784, 4,820,850 and 4,916,239), pravastatin (PRAVACHOL®; see U.S. Patent Nos. 4,346,227, 4,537,859, 4,410,629, 5,030,447 and 5,180,589), fluvastatin (LESCOL®; see U.S. Patent Nos. 5,354,772, 4,911,165, 4,929,437, 5,189,164, 5,118,853, 5,290,946 and 5,356,896), atorvastatin (LIPITOR®; see U.S. Patent Nos. 5,273,995, 4,681,893, 5,489,691 and 5,342,952) and cerivastatin (also known as rivastatin and BAYCHOL®; see US Patent No. 5,177,080). The structural formulas of these and additional HMG-CoA reductase inhibitors that may be used in the instant methods are described at page 87 of M. Yalpani, "Cholesterol Lowering Drugs", *Chemistry & Industry*, pp. 85-89 (5 February 1996) and US Patent Nos. 4,782,084 and 4,885,314. The term HMG-CoA reductase inhibitor as used herein includes all pharmaceutically acceptable lactone and open-acid forms (i.e., where the lactone ring is opened to form the free acid) as well as salt and ester forms of compounds which have HMG-CoA reductase inhibitory activity, and therefor the use of such salts, esters, open-acid and lactone forms is included within the scope of this invention. An illustration of the lactone portion and its corresponding open-acid-form is shown below as structures I and II.



20

In HMG-CoA reductase inhibitors where an open-acid form can exist, salt and ester forms may be formed from the open-acid, and all such forms are included within the meaning of the term "HMG-CoA reductase inhibitor" as used herein. In an embodiment, the HMG-CoA reductase inhibitor is selected from lovastatin and simvastatin, and in a further embodiment, simvastatin. Herein, the term "pharmaceutically acceptable salts" with respect to the HMG-CoA reductase

25

inhibitor shall mean non-toxic salts of the compounds employed in this invention which are generally prepared by reacting the free acid with a suitable organic or inorganic base, particularly those formed from cations such as sodium, potassium, aluminum, calcium, lithium, magnesium, zinc and tetramethylammonium, as well as those salts formed from amines such as ammonia, ethylenediamine, N-methylglucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chloroprocaine, diethanolamine, procaine, N-benzylphenethylamine, 1-p-chlorobenzyl-2-pyrrolidine-1'-yl-methylbenzimidazole, diethylamine, piperazine, and tris(hydroxymethyl) aminomethane. Further examples of salt forms of HMG-CoA reductase inhibitors may include, but are not limited to, acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynapthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylsulfate, mucate, napsylate, nitrate, oleate, oxalate, pamaote, palmitate, panthothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide, and valerate.

Ester derivatives of the described HMG-CoA reductase inhibitor compounds may act as prodrugs which, when absorbed into the bloodstream of a warm-blooded animal, may cleave in such a manner as to release the drug form and permit the drug to afford improved therapeutic efficacy.

"Prenyl-protein transferase inhibitor" refers to a compound which inhibits any one or any combination of the prenyl-protein transferase enzymes, including farnesyl-protein transferase (FPTase), geranylgeranyl-protein transferase type I (GGPTase-I), and geranylgeranyl-protein transferase type-II (GGPTase-II, also called Rab GGPTase). Examples of prenyl-protein transferase inhibiting compounds include (+)-6-[amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-4-(3-chlorophenyl)-1-methyl-2(1H)-quinolinone, (-)-6-[amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-4-(3-chlorophenyl)-1-methyl-2(1H)-quinolinone, (+)-6-[amino(4-chlorophenyl)(1-methyl-1H-imidazol-5-yl)methyl]-4-(3-chlorophenyl)-1-methyl-2(1H)-quinolinone, 5(S)-n-butyl-1-(2,3-dimethylphenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone, (S)-1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-5-[2-(ethanesulfonyl)methyl]-2-piperazinone,



5(S)-n-Butyl-1-(2-methylphenyl)-4-[1-(4-cyanobenzyl)-5-imidazolylmethyl]-2-piperazinone, 1-(3-chlorophenyl)-4-[1-(4-cyanobenzyl)-2-methyl-5-imidazolylmethyl]-2-piperazinone, 1-(2,2-diphenylethyl)-3-[N-(1-(4-cyanobenzyl)-1H-imidazol-5-ylethyl)carbamoyl]piperidine, 4-{5-[4-hydroxymethyl-4-(4-chloropyridin-2-ylmethyl)-piperidine-1-ylmethyl]-2-methylimidazol-1-ylmethyl} benzonitrile, 4-{5-[4-hydroxymethyl-4-(3-chlorobenzyl)-piperidine-1-ylmethyl]-2-methylimidazol-1-ylmethyl} benzonitrile, 4-{3-[4-(2-oxo-2H-pyridin-1-yl)benzyl]-3H-imidazol-4-ylmethyl} benzonitrile, 4-{3-[4-(5-chloro-2-oxo-2H-[1,2']bipyridin-5'-ylmethyl)-3H-imidazol-4-ylmethyl} benzonitrile, 4-{3-[4-(2-oxo-2H-[1,2'] bipyridin-5'-ylmethyl)-3H-imidazol-4-ylmethyl} benzonitrile, 4-[3-(2-oxo-1-phenyl-1,2-dihydropyridin-4-ylmethyl)-3H-imidazol-4-ylmethyl} benzonitrile, 18,19-dihydro-19-oxo-5*H*,17*H*-6,10:12,16-dimetheno-1H-imidazo[4,3-*c*][1,11,4]dioxazacyclo-nonadecine-9-carbonitrile, (±)-19,20-dihydro-19-oxo-5*H*-18,21-ethano-12,14-etheno-6,10-metheno-22*H*-benzo[*d*]imidazo[4,3-*k*][1,6,9,12]oxatriaza-cyclooctadecine-9-carbonitrile, 19,20-dihydro-19-oxo-5*H*,17*H*-18,21-ethano-6,10:12,16-dimetheno-22*H*-imidazo[3,4-*h*][1,8,11,14]oxatriazacycloeicosine-9-carbonitrile, and (±)-19,20-dihydro-3-methyl-19-oxo-5*H*-18,21-ethano-12,14-etheno-6,10-metheno-22*H*-benzo[*d*]imidazo[4,3-*k*][1,6,9,12]oxa-triazacyclooctadecine-9-carbonitrile.

Other examples of prenyl-protein transferase inhibitors can be found in the following publications and patents: WO 96/30343, WO 97/18813, WO 97/21701, WO 97/23478, WO 97/38665, WO 98/28980, WO 98/29119, WO 95/32987, U.S. Patent No. 5,420,245, U.S. Patent No. 5,523,430, U.S. Patent No. 5,532,359, U.S. Patent No. 5,510,510, U.S. Patent No. 5,589,485, U.S. Patent No. 5,602,098, European Patent Publ. 0 618 221, European Patent Publ. 0 675 112, European Patent Publ. 0 604 181, European Patent Publ. 0 696 593, WO 94/19357, WO 95/08542, WO 95/11917, WO 95/12612, WO 95/12572, WO 95/10514, U.S. Patent No. 5,661,152, WO 95/10515, WO 95/10516, WO 95/24612, WO 95/34535, WO 95/25086, WO 96/05529, WO 96/06138, WO 96/06193, WO 96/16443, WO 96/21701, WO 96/21456, WO 96/22278, WO 96/24611, WO 96/24612, WO 96/05168, WO 96/05169, WO 96/00736, U.S. Patent No. 5,571,792, WO 96/17861, WO 96/33159, WO 96/34850, WO 96/34851, WO 96/30017, WO 96/30018, WO 96/30362, WO 96/30363, WO 96/31111, WO 96/31477, WO 96/31478, WO 96/31501, WO 97/00252, WO 97/03047, WO 97/03050, WO 97/04785, WO 97/02920, WO 97/17070, WO 97/23478, WO 97/26246, WO 97/30053, WO 97/44350, WO 98/02436, and U.S. Patent No. 5,532,359.

For an example of the role of a prenyl-protein transferase inhibitor on angiogenesis see European J. of Cancer, Vol. 35, No. 9, pp.1394-1401 (1999).

“Angiogenesis inhibitors” refers to compounds that inhibit the formation of new blood vessels, regardless of mechanism. Examples of angiogenesis inhibitors include, but are not limited to, tyrosine kinase inhibitors, such as inhibitors of the tyrosine kinase receptors Flt-1 (VEGFR1) and Flk-1/KDR (VEGFR2), inhibitors of epidermal-derived, fibroblast-derived, or platelet derived growth factors, MMP (matrix metalloprotease) inhibitors, integrin blockers, interferon- $\alpha$ , interleukin-12, pentosan polysulfate, cyclooxygenase inhibitors, including nonsteroidal anti-inflammatories (NSAIDs) like aspirin and ibuprofen as well as selective cyclooxygenase-2 inhibitors like celecoxib and rofecoxib (PNAS, Vol. 89, p. 7384 (1992); JNCI, Vol. 69, p. 475 (1982); Arch. Ophthalmol., Vol. 108, p.573 (1990); Anat. Rec., Vol. 238, p. 68 (1994); FEBS Letters, Vol. 372, p. 83 (1995); Clin. Orthop. Vol. 313, p. 76 (1995); J. Mol. Endocrinol., Vol. 16, p.107 (1996); Jpn. J. Pharmacol., Vol. 75, p. 105 (1997); Cancer Res., Vol. 57, p. 1625 (1997); Cell, Vol. 93, p. 705 (1998); Intl. J. Mol. Med., Vol. 2, p. 715 (1998); J. Biol. Chem., Vol. 274, p. 9116 (1999)), steroidal anti-inflammatories (such as corticosteroids, mineralocorticoids, dexamethasone, prednisone, prednisolone, methylpred, betamethasone), carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl)-fumagillol, thalidomide, angiostatin, troponin-1, angiotensin II antagonists (see Fernandez et al., J. Lab. Clin. Med. 105:141-145 (1985)), and antibodies to VEGF (see, Nature Biotechnology, Vol. 17, pp.963-968 (October 1999); Kim et al., Nature, 362, 841-844 (1993); WO 00/44777; and WO 00/61186).

Other therapeutic agents that modulate or inhibit angiogenesis and may also be used in combination with the compounds of the instant invention include agents that modulate or inhibit the coagulation and fibrinolysis systems (see review in *Clin. Chem. La. Med.* 38:679-692 (2000)). Examples of such agents that modulate or inhibit the coagulation and fibrinolysis pathways include, but are not limited to, heparin (see *Thromb. Haemost.* 80:10-23 (1998)), low molecular weight heparins, GPIIb/IIIa antagonists (such as tirofiban), warfarin, thrombin inhibitors and carboxypeptidase U inhibitors (also known as inhibitors of active thrombin activatable fibrinolysis inhibitor [TAFIa]) (see *Thrombosis Res.* 101:329-354 (2001)). TAFIa inhibitors have been described in U.S. Serial Nos. 60/310,927 (filed August 8, 2001) and 60/349,925 (filed January 18, 2002).

“Agents that interfere with cell cycle checkpoints” refer to compounds that inhibit protein kinases that transduce cell cycle checkpoint signals, thereby sensitizing the cancer cell to DNA damaging agents. Such agents include inhibitors of ATR, ATM, the Chk1 and Chk2 kinases and cdk and cdc kinase inhibitors and are specifically exemplified by 7-hydroxystaurosporin, flavopiridol, CYC202 (Cyclacel) and BMS-387032.

“Inhibitors of cell proliferation and survival signalling pathway” refer to compounds that inhibit signal transduction cascades downstream of cell surface receptors. Such agents include inhibitors of serine/threonine kinases (including but not limited to inhibitors of Akt such as described in WO 02/083064, WO 02/083139, WO 02/083140 and WO 02/083138), inhibitors of Raf kinase (for example BAY-43-9006), inhibitors of MEK (for example CI-1040 and PD-098059), inhibitors of mTOR (for example Wyeth CCI-779), and inhibitors of PI3K (for example LY294002).

The combinations with NSAID's are directed to the use of NSAID's which are potent COX-2 inhibiting agents. For purposes of this specification an NSAID is potent if it possess an IC<sub>50</sub> for the inhibition of COX-2 of 1μM or less as measured by cell or microsomal assays.


The invention also encompasses combinations with NSAID's which are selective COX-2 inhibitors. For purposes of this specification NSAID's which are selective inhibitors of COX-2 are defined as those which possess a specificity for inhibiting COX-2 over COX-1 of at least 100 fold as measured by the ratio of IC<sub>50</sub> for COX-2 over IC<sub>50</sub> for COX-1 evaluated by cell or microsomal assays. Such compounds include, but are not limited to those disclosed in U.S. Patent 5,474,995, issued December 12, 1995, U.S. Patent 5,861,419, issued January 19, 1999, U.S. Patent 6,001,843, issued December 14, 1999, U.S. Patent 6,020,343, issued February 1, 2000, U.S. Patent 5,409,944, issued April 25, 1995, U.S. Patent 5,436,265, issued July 25, 1995, U.S. Patent 5,536,752, issued July 16, 1996, U.S. Patent 5,550,142, issued August 27, 1996, U.S. Patent 5,604,260, issued February 18, 1997, U.S. Patent 5,698,584, issued December 16, 1997, U.S. Patent 5,710,140, issued January 20, 1998, WO 94/15932, published July 21, 1994, U.S. Patent 5,344,991, issued June 6, 1994, U.S. Patent 5,134,142, issued July 28, 1992, U.S. Patent 5,380,738, issued January 10, 1995, U.S. Patent 5,393,790, issued February 20, 1995, U.S. Patent 5,466,823, issued November 14, 1995, U.S. Patent 5,633,272, issued May 27, 1997,

and U.S. Patent 5,932,598, issued August 3, 1999, all of which are hereby incorporated by reference.


Inhibitors of COX-2 that are particularly useful in the instant method of treatment are:

5

3-phenyl-4-(4-(methylsulfonyl)phenyl)-2-(5*H*)-furanone; and



10 5-chloro-3-(4-(methylsulfonyl)phenyl)-2-(2-methyl-5-pyridinyl)pyridine;






or a pharmaceutically acceptable salt thereof.

15

General and specific synthetic procedures for the preparation of the COX-2 inhibitor compounds described above are found in U.S. Patent No. 5,474,995, issued December 12, 1995, U.S. Patent No. 5,861,419, issued January 19, 1999, and U.S. Patent No. 6,001,843, issued December 14, 1999, all of which are herein incorporated by reference.

20

Compounds that have been described as specific inhibitors of COX-2 and are therefore useful in the present invention include, but are not limited to, the following:



or a pharmaceutically acceptable salt thereof.

- 5                   Compounds which are described as specific inhibitors of COX-2 and are therefore useful in the present invention, and methods of synthesis thereof, can be found in the following patents, pending applications and publications, which are herein incorporated by reference: WO 94/15932, published July 21, 1994, U.S. Patent No. 5,344,991, issued June 6, 1994, U.S. Patent No. 5,134,142, issued July 28, 1992, U.S. Patent No. 5,380,738, issued January 10, 1995, U.S. Patent No. 5,393,790, 10                   issued February 20, 1995, U.S. Patent No. 5,466,823, issued November 14, 1995, U.S. Patent No. 5,633,272, issued May 27, 1997, and U.S. Patent No. 5,932,598, issued August 3, 1999.

Compounds which are specific inhibitors of COX-2 and are therefore useful in the present invention, and methods of synthesis thereof, can be found in the following patents, pending applications and publications, which are herein incorporated by reference: U.S. Patent No. 5,474,995, issued December 12, 1995, U.S. Patent No. 5,861,419, issued January 19, 1999, U.S. Patent No. 6,001,843, issued December 14, 1999, U.S. Patent No. 6,020,343, issued February 1, 2000, U.S. Patent No. 5,409,944, issued April 25, 1995, U.S. Patent No. 5,436,265, issued July 25, 1995, U.S. Patent No. 5,536,752, issued July 16, 1996, U.S. Patent No. 5,550,142, issued August 27, 1996, U.S. Patent No. 5,604,260, issued February 18, 1997, U.S. Patent No. 5,698,584, issued December 16, 1997, and U.S. Patent No. 5,710,140, issued January 20, 1998.

Other examples of angiogenesis inhibitors include, but are not limited to, endostatin, ukrain, ranpirnase, IM862, 5-methoxy-4-[2-methyl-3-(3-methyl-2-butenyl)oxiranyl]-1-oxaspiro[2,5]oct-6-yl(chloroacetyl)carbamate, acetyldinanaline, 5-amino-1-[[3,5-dichloro-4-(4-chlorobenzoyl)phenyl]methyl]-1H-1,2,3-triazole-4-carboxamide, CM101, squalamine, combretastatin, RPI4610, NX31838, sulfated mannopentaose phosphate, 7,7-(carbonyl-bis[imino-N-methyl-4,2-pyrrolocarbonylimino[N-methyl-4,2-pyrrole]-carbonylimino]-bis-(1,3-naphthalene disulfonate), and 3-[(2,4-dimethylpyrrol-5-yl)methylene]-2-indolinone (SU5416).

As used above, "integrin blockers" refers to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the  $\alpha_v\beta_3$  integrin, to compounds which selectively antagonize, inhibit or counteract binding of a physiological ligand to the  $\alpha_v\beta_5$  integrin, to compounds which antagonize, inhibit or counteract binding of a physiological ligand to both the  $\alpha_v\beta_3$  integrin and the  $\alpha_v\beta_5$  integrin, and to compounds which antagonize, inhibit or counteract the activity of the particular integrin(s) expressed on capillary endothelial cells. The term also refers to antagonists of the  $\alpha_v\beta_6$ ,  $\alpha_v\beta_8$ ,  $\alpha_1\beta_1$ ,  $\alpha_2\beta_1$ ,  $\alpha_5\beta_1$ ,  $\alpha_6\beta_1$  and  $\alpha_6\beta_4$  integrins. The term also refers to antagonists of any combination of  $\alpha_v\beta_3$ ,  $\alpha_v\beta_5$ ,  $\alpha_v\beta_6$ ,  $\alpha_v\beta_8$ ,  $\alpha_1\beta_1$ ,  $\alpha_2\beta_1$ ,  $\alpha_5\beta_1$ ,  $\alpha_6\beta_1$  and  $\alpha_6\beta_4$  integrins.

Some specific examples of tyrosine kinase inhibitors include N-(trifluoromethylphenyl)-5-methylisoxazol-4-carboxamide, 3-[(2,4-dimethylpyrrol-5-yl)methylidenyl]indolin-2-one, 17-(allylamino)-17-demethoxygeldanamycin, 4-(3-chloro-4-fluorophenylamino)-7-methoxy-6-[3-(4-morpholinyl)propoxyl]quinazoline, N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)-4-quinazolinamine, BIBX1382, 2,3,9,10,11,12-hexahydro-10-(hydroxymethyl)-10-hydroxy-9-methyl-9,12-epoxy-1H-

diindolo[1,2,3-fg:3',2',1'-kl]pyrrolo[3,4-i][1,6]benzodiazocin-1-one, SH268, genistein, STI571, CEP2563, 4-(3-chlorophenylamino)-5,6-dimethyl-7H-pyrrolo[2,3-d]pyrimidinemethane sulfonate, 4-(3-bromo-4-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, 4-(4'-hydroxyphenyl)amino-6,7-dimethoxyquinazoline, 5 SU6668, STI571A, N-4-chlorophenyl-4-(4-pyridylmethyl)-1-phthalazinamine, and EMD121974.

Combinations with compounds other than anti-cancer compounds are also encompassed in the instant methods. For example, combinations of the instantly claimed compounds with PPAR- $\gamma$  (i.e., PPAR-gamma) agonists and PPAR- $\delta$  (i.e., 10 PPAR-delta) agonists are useful in the treatment of certain malignancies. PPAR- $\gamma$  and PPAR- $\delta$  are the nuclear peroxisome proliferator-activated receptors  $\gamma$  and  $\delta$ . The expression of PPAR- $\gamma$  on endothelial cells and its involvement in angiogenesis has been reported in the literature (see *J. Cardiovasc. Pharmacol.* 1998; 31:909-913; *J. Biol. Chem.* 1999;274:9116-9121; *Invest. Ophthalmol Vis. Sci.* 2000; 41:2309-2317). 15 More recently, PPAR- $\gamma$  agonists have been shown to inhibit the angiogenic response to VEGF in vitro; both troglitazone and rosiglitazone maleate inhibit the development of retinal neovascularization in mice. (*Arch. Ophthalmol.* 2001; 119:709-717). Examples of PPAR- $\gamma$  agonists and PPAR-  $\gamma/\alpha$  agonists include, but are not limited to, thiazolidinediones (such as DRF2725, CS-011, troglitazone, rosiglitazone, and 20 pioglitazone), fenofibrate, gemfibrozil, clofibrate, GW2570, SB219994, AR-H039242, JTT-501, MCC-555, GW2331, GW409544, NN2344, KRP297, NP0110, DRF4158, NN622, GI262570, PNU182716, DRF552926, 2-[(5,7-dipropyl-3-trifluoromethyl-1,2-benzisoxazol-6-yl)oxy]-2-methylpropionic acid (disclosed in USSN 09/782,856), and 2(R)-7-(3-(2-chloro-4-(4-fluorophenoxy) phenoxy)propoxy)- 25 2-ethylchromane-2-carboxylic acid (disclosed in USSN 60/235,708 and 60/244,697).

Another embodiment of the instant invention is the use of the presently disclosed compounds in combination with gene therapy for the treatment of cancer. For an overview of genetic strategies to treating cancer see Hall et al (*Am J Hum Genet* 61:785-789, 1997) and Kufe et al (*Cancer Medicine*, 5th Ed, pp 876-889, BC 30 Decker, Hamilton 2000). Gene therapy can be used to deliver any tumor suppressing gene. Examples of such genes include, but are not limited to, p53, which can be delivered via recombinant virus-mediated gene transfer (see U.S. Patent No. 6,069,134, for example), a uPA/uPAR antagonist ("Adenovirus-Mediated Delivery of a uPA/uPAR Antagonist Suppresses Angiogenesis-Dependent Tumor Growth and

Dissemination in Mice," *Gene Therapy*, August 1998;5(8):1105-13), and interferon gamma (*J Immunol* 2000;164:217-222).

The compounds of the instant invention may also be administered in combination with an inhibitor of inherent multidrug resistance (MDR), in particular  
5 MDR associated with high levels of expression of transporter proteins. Such MDR inhibitors include inhibitors of p-glycoprotein (P-gp), such as LY335979, XR9576, OC144-093, R101922, VX853 and PSC833 (valspodar).

A compound of the present invention may be employed in conjunction with anti-emetic agents to treat nausea or emesis, including acute, delayed, late-phase,  
10 and anticipatory emesis, which may result from the use of a compound of the present invention, alone or with radiation therapy. For the prevention or treatment of emesis, a compound of the present invention may be used in conjunction with other anti-emetic agents, especially neurokinin-1 receptor antagonists, 5HT3 receptor  
15 antagonists, such as ondansetron, granisetron, tropisetron, and zatisetron, GABAB receptor agonists, such as baclofen, a corticosteroid such as Decadron (dexamethasone), Kenalog, Aristocort, Nasalide, Preferid, Benecorten or others such as disclosed in U.S. Patent Nos. 2,789,118, 2,990,401, 3,048,581, 3,126,375, 3,929,768, 3,996,359, 3,928,326 and 3,749,712, an antidopaminergic, such as the phenothiazines (for example prochlorperazine, fluphenazine, thioridazine and  
20 mesoridazine), metoclopramide or dronabinol. For the treatment or prevention of emesis that may result upon administration of the instant compounds, conjunctive therapy with an anti-emesis agent selected from a neurokinin-1 receptor antagonist, a 5HT3 receptor antagonist and a corticosteroid is preferred.

Neurokinin-1 receptor antagonists of use in conjunction with the  
25 compounds of the present invention are fully described, for example, in U.S. Patent Nos. 5,162,339, 5,232,929, 5,242,930, 5,373,003, 5,387,595, 5,459,270, 5,494,926, 5,496,833, 5,637,699, 5,719,147; European Patent Publication Nos. EP 0 360 390, 0 394 989, 0 428 434, 0 429 366, 0 430 771, 0 436 334, 0 443 132, 0 482 539, 0 498 069, 0 499 313, 0 512 901, 0 512 902, 0 514 273, 0 514 274, 0 514 275,  
30 0 514 276, 0 515 681, 0 517 589, 0 520 555, 0 522 808, 0 528 495, 0 532 456, 0 533 280, 0 536 817, 0 545 478, 0 558 156, 0 577 394, 0 585 913, 0 590 152, 0 599 538, 0 610 793, 0 634 402, 0 686 629, 0 693 489, 0 694 535, 0 699 655, 0 699 674, 0 707 006, 0 708 101, 0 709 375, 0 709 376, 0 714 891, 0 723 959, 0 733 632 and 0 776 893; PCT International Patent Publication Nos. WO 90/05525,  
35 90/05729, 91/09844, 91/18899, 92/01688, 92/06079, 92/12151, 92/15585, 92/17449,



92/20661, 92/20676, 92/21677, 92/22569, 93/00330, 93/00331, 93/01159, 93/01165, 93/01169, 93/01170, 93/06099, 93/09116, 93/10073, 93/14084, 93/14113, 93/18023, 93/19064, 93/21155, 93/21181, 93/23380, 93/24465, 94/00440, 94/01402, 94/02461, 94/02595, 94/03429, 94/03445, 94/04494, 94/04496, 94/05625, 94/07843, 94/08997, 5 94/10165, 94/10167, 94/10168, 94/10170, 94/11368, 94/13639, 94/13663, 94/14767, 94/15903, 94/19320, 94/19323, 94/20500, 94/26735, 94/26740, 94/29309, 95/02595, 95/04040, 95/04042, 95/06645, 95/07886, 95/07908, 95/08549, 95/11880, 95/14017, 95/15311, 95/16679, 95/17382, 95/18124, 95/18129, 95/19344, 95/20575, 95/21819, 95/22525, 95/23798, 95/26338, 95/28418, 95/30674, 95/30687, 95/33744, 96/05181, 10 96/05193, 96/05203, 96/06094, 96/07649, 96/10562, 96/16939, 96/18643, 96/20197, 96/21661, 96/29304, 96/29317, 96/29326, 96/29328, 96/31214, 96/32385, 96/37489, 97/01553, 97/01554, 97/03066, 97/08144, 97/14671, 97/17362, 97/18206, 97/19084, 97/19942 and 97/21702; and in British Patent Publication Nos. 2 266 529, 2 268 931, 2 269 170, 2 269 590, 2 271 774, 2 292 144, 2 293 168, 2 293 169, and 2 302 689.

15 The preparation of such compounds is fully described in the aforementioned patents and publications, which are incorporated herein by reference.

In an embodiment, the neurokinin-1 receptor antagonist for use in conjunction with the compounds of the present invention is selected from: 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(3-(5-oxo- 20 1H,4H-1,2,4-triazolo)methyl)morpholine, or a pharmaceutically acceptable salt thereof, which is described in U.S. Patent No. 5,719,147.

A compound of the instant invention may also be administered with an agent useful in the treatment of anemia. Such an anemia treatment agent is, for example, a continuous erythropoiesis receptor activator (such as epoetin alfa).

25 A compound of the instant invention may also be administered with an agent useful in the treatment of neutropenia. Such a neutropenia treatment agent is, for example, a hematopoietic growth factor which regulates the production and function of neutrophils such as a human granulocyte colony stimulating factor, (G-CSF). Examples of a G-CSF include filgrastim.

30 A compound of the instant invention may also be administered with an immunologic-enhancing drug, such as levamisole, isoprinosine and Zadaxin.

Thus, the scope of the instant invention encompasses the use of the instantly claimed compounds in combination with a second compound selected from:

- 1) an estrogen receptor modulator,
- 35 2) an androgen receptor modulator,

- 3) retinoid receptor modulator,
- 4) a cytotoxic/cytostatic agent,
- 5) an antiproliferative agent,
- 6) a prenyl-protein transferase inhibitor,
- 5 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor,
- 10) an angiogenesis inhibitor,
- 11) a PPAR- $\gamma$  agonists,
- 10 12) a PPAR- $\delta$  agonists,
- 13) an inhibitor of inherent multidrug resistance,
- 14) an anti-emetic agent,
- 15) an agent useful in the treatment of anemia,
- 16) an agent useful in the treatment of neutropenia,
- 15 17) an immunologic-enhancing drug,
- 18) an inhibitor of cell proliferation and survival signaling, and
- 19) an agent that interferes with a cell cycle checkpoint.

The term "administration" and variants thereof (e.g., "administering" a compound) in reference to a compound of the invention means introducing the compound or a prodrug of the compound into the system of the animal in need of treatment. When a compound of the invention or prodrug thereof is provided in combination with one or more other active agents (e.g., a cytotoxic agent, etc.), "administration" and its variants are each understood to include concurrent and sequential introduction of the compound or prodrug thereof and other agents.

25 As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

30 The term "therapeutically effective amount" as used herein means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician.

The term "treating cancer" or "treatment of cancer" refers to administration to a mammal afflicted with a cancerous condition and refers to an

effect that alleviates the cancerous condition by killing the cancerous cells, but also to an effect that results in the inhibition of growth and/or metastasis of the cancer.

- In an embodiment, the angiogenesis inhibitor to be used as the second compound is selected from a tyrosine kinase inhibitor, an inhibitor of epidermal-  
 5 derived growth factor, an inhibitor of fibroblast-derived growth factor, an inhibitor of platelet derived growth factor, an MMP (matrix metalloprotease) inhibitor, an integrin blocker, interferon- $\alpha$ , interleukin-12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-chloroacetyl-carbonyl)-fumagillol, thalidomide, angiostatin, troponin-1, or an antibody to VEGF.  
 10 In an embodiment, the estrogen receptor modulator is tamoxifen or raloxifene.

Also included in the scope of the claims is a method of treating cancer that comprises administering a therapeutically effective amount of a compound of Formula I in combination with radiation therapy and/or in combination with a compound selected from:

- |    |  |
|----|--|
| 15 | 1) an estrogen receptor modulator,                                 |
|    | 2) an androgen receptor modulator,                                 |
|    | 3) a retinoid receptor modulator,                                  |
|    | 4) a cytotoxic/cytostatic agent,                                   |
|    | 5) an antiproliferative agent,                                     |
| 20 | 6) a prenyl-protein transferase inhibitor,                         |
|    | 7) an HMG-CoA reductase inhibitor,                                 |
|    | 8) an HIV protease inhibitor,                                      |
|    | 9) a reverse transcriptase inhibitor,                              |
|    | 10) an angiogenesis inhibitor,                                     |
| 25 | 11) PPAR- $\gamma$ agonists,                                       |
|    | 12) PPAR- $\delta$ agonists,                                       |
|    | 13) an inhibitor of inherent multidrug resistance,                 |
|    | 14) an anti-emetic agent,  |
|    | 15) an agent useful in the treatment of anemia,                    |
| 30 | 16) an agent useful in the treatment of neutropenia,               |
|    | 17) an immunologic-enhancing drug,                                 |
|    | 18) an inhibitor of cell proliferation and survival signaling, and |
|    | 19) an agent that interferes with a cell cycle checkpoint.         |

And yet another embodiment of the invention is a method of treating cancer that comprises administering a therapeutically effective amount of a compound of Formula I in combination with paclitaxel or trastuzumab.

The invention further encompasses a method of treating or preventing  
5 cancer that comprises administering a therapeutically effective amount of a compound of Formula I in combination with a COX-2 inhibitor.

The instant invention also includes a pharmaceutical composition useful for treating or preventing cancer that comprises a therapeutically effective amount of a compound of Formula I and a compound selected from:

- 10 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 3) a retinoid receptor modulator,
- 4) a cytotoxic/cytostatic agent,
- 5) an antiproliferative agent,
- 15 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor,
- 10) an angiogenesis inhibitor, and
- 20 11) a PPAR- $\gamma$  agonist,
- 12) a PPAR- $\delta$  agonists;
- 13) an inhibitor of cell proliferation and survival signaling, and
- 14) an agent that interferes with a cell cycle checkpoint.

The invention further comprises the use of the instant compounds in a  
25 method to screen for other compounds that bind to KSP. To employ the compounds of the invention in a method of screening for compounds that bind to KSP kinesin, the KSP is bound to a support, and a compound of the invention (which is a mitotic agent) is added to the assay. Alternatively, the compound of the invention is bound to the support and KSP is added. Classes of compounds among which novel binding  
30 agents may be sought include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. Of particular interest are screening assays for candidate agents that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including labeled *in vitro* protein-protein binding assays, electrophoretic mobility shift assays, immunoassays  
35 for protein binding, functional assays (phosphorylation assays, etc.) and the like.

The determination of the binding of the mitotic agent to KSP may be done in a number of ways. In a preferred embodiment, the mitotic agent (the compound of the invention) is labeled, for example, with a fluorescent or radioactive moiety and binding determined directly. For example, this may be done by attaching  
5 all or a portion of KSP to a solid support, adding a labeled mitotic agent (for example a compound of the invention in which at least one atom has been replaced by a detectable isotope), washing off excess reagent, and determining whether the amount of the label is that present on the solid support. Various blocking and washing steps may be utilized as is known in the art.

10 By "labeled" herein is meant that the compound is either directly or indirectly labeled with a label which provides a detectable signal, e.g., radioisotope, fluorescent tag, enzyme, antibodies, particles such as magnetic particles, chemiluminescent tag, or specific binding molecules, etc. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin etc. For the  
15 specific-binding members, the complementary member would normally be labeled with a molecule which provides for detection, in accordance with known procedures, as outlined above. The label can directly or indirectly provide a detectable signal.

In some embodiments, only one of the components is labeled. For example, the kinesin proteins may be labeled at tyrosine positions using  $^{125}\text{I}$ , or with  
20 fluorophores. Alternatively, more than one component may be labeled with different labels; using  $^{125}\text{I}$  for the proteins, for example, and a fluorophor for the mitotic agents.

The compounds of the invention may also be used as competitors to screen for additional drug candidates. "Candidate bioactive agent" or "drug candidate" or grammatical equivalents as used herein describe any molecule, e.g., protein,  
25 oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for bioactivity. They may be capable of directly or indirectly altering the cellular proliferation phenotype or the expression of a cellular proliferation sequence, including both nucleic acid sequences and protein sequences. In other cases, alteration of cellular proliferation protein binding and/or activity is screened. Screens  
30 of this sort may be performed either in the presence or absence of microtubules. In the case where protein binding or activity is screened, preferred embodiments exclude molecules already known to bind to that particular protein, for example, polymer structures such as microtubules, and energy sources such as ATP. Preferred  
35 embodiments of assays herein include candidate agents which do not bind the cellular proliferation protein in its endogenous native state termed herein as "exogenous"

agents. In another preferred embodiment, exogenous agents further exclude antibodies to KSP.

Candidate agents can encompass numerous chemical classes, though typically they are organic molecules, preferably small organic compounds having a molecular weight of more than 100 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding and lipophilic binding, and typically include at least an amine, carbonyl, hydroxyl, ether, or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof. Particularly preferred are peptides.

Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, amidification to produce structural analogs.

Competitive screening assays may be done by combining KSP and a drug candidate in a first sample. A second sample comprises a mitotic agent, KSP and a drug candidate. This may be performed in either the presence or absence of microtubules. The binding of the drug candidate is determined for both samples, and a change, or difference in binding between the two samples indicates the presence of an agent capable of binding to KSP and potentially modulating its activity. That is, if the binding of the drug candidate is different in the second sample relative to the first sample, the drug candidate is capable of binding to KSP.

In a preferred embodiment, the binding of the candidate agent is determined through the use of competitive binding assays. In this embodiment, the competitor is a binding moiety known to bind to KSP, such as an antibody, peptide,

binding partner, ligand, etc. Under certain circumstances, there may be competitive binding as between the candidate agent and the binding moiety, with the binding moiety displacing the candidate agent.

5 In one embodiment, the candidate agent is labeled. Either the candidate agent, or the competitor, or both, is added first to KSP for a time sufficient to allow binding, if present. Incubations may be performed at any temperature which facilitates optimal activity, typically between about 4 and about 40°C.

10 Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high throughput screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

15 In a preferred embodiment, the competitor is added first, followed by the candidate agent. Displacement of the competitor is an indication the candidate agent is binding to KSP and thus is capable of binding to, and potentially modulating, the activity of KSP. In this embodiment, either component can be labeled. Thus, for example, if the competitor is labeled, the presence of label in the wash solution indicates displacement by the agent. Alternatively, if the candidate agent is labeled, the presence of the label on the support indicates displacement.

20 In an alternative embodiment, the candidate agent is added first, with incubation and washing, followed by the competitor. The absence of binding by the competitor may indicate the candidate agent is bound to KSP with a higher affinity. Thus, if the candidate agent is labeled, the presence of the label on the support, coupled with a lack of competitor binding, may indicate the candidate agent is capable  
25 of binding to KSP.

It may be of value to identify the binding site of KSP. This can be done in a variety of ways. In one embodiment, once KSP has been identified as binding to the mitotic agent, KSP is fragmented or modified and the assays repeated to identify the necessary components for binding.

30 Modulation is tested by screening for candidate agents capable of modulating the activity of KSP comprising the steps of combining a candidate agent with KSP, as above, and determining an alteration in the biological activity of KSP. Thus, in this embodiment, the candidate agent should both bind to KSP (although this may not be necessary), and alter its biological or biochemical activity as defined  
35 herein. The methods include both in vitro screening methods and in vivo screening of

cells for alterations in cell cycle distribution, cell viability, or for the presence, morphology, activity, distribution, or amount of mitotic spindles, as are generally outlined above.

Alternatively, differential screening may be used to identify drug  
5 candidates that bind to the native KSP, but cannot bind to modified KSP.

Positive controls and negative controls may be used in the assays.  
Preferably all control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of all samples is for a time sufficient for the binding of the agent to the protein. Following incubation, all samples are washed free  
10 of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples may be counted in a scintillation counter to determine the amount of bound compound.

A variety of other reagents may be included in the screening assays.  
These include reagents like salts, neutral proteins, e.g., albumin, detergents, etc which  
15 may be used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may be used. The mixture of components may be added in any order that provides for the requisite binding.

20 These and other aspects of the invention will be apparent from the teachings contained herein.

## ASSAYS

25 The compounds of the instant invention described in the Examples were tested by the assays described below and were found to have kinesin inhibitory activity. Other assays are known in the literature and could be readily performed by those of skill in the art (see, for example, PCT Publication WO 01/30768, May 3, 2001, pages 18-22).

30

### I. Kinesin ATPase *In Vitro* Assay

**Cloning and expression of human poly-histidine tagged KSP motor domain (KSP(367H))**



Plasmids for the expression of the human KSP motor domain construct were cloned by PCR using a pBluescript full length human KSP construct (Blangy et al., Cell, vol.83, pp1159-1169, 1995) as a template. The N-terminal primer 5'-GCAACGATTAATATGGCGTCGCAGCCAAATTCGTCTGCGAAG (SEQ.ID.NO.: 1) and the C-terminal primer 5'-GCAACGCTCGAGTCAGTGAT GATGGTGGTGATGCTGATTCACTTCAGGCTTATTCAATAT (SEQ.ID.NO.: 2) were used to amplify the motor domain and the neck linker region. The PCR products were digested with AseI and XhoI, ligated into the NdeI/XhoI digestion product of pRSETa (Invitrogen) and transformed into E. coli BL21 (DE3).

Cells were grown at 37°C to an OD<sub>600</sub> of 0.5. After cooling the culture to room temperature expression of KSP was induced with 100µM IPTG and incubation was continued overnight. Cells were pelleted by centrifugation and washed once with ice-cold PBS. Pellets were flash-frozen and stored -80°C.

#### 15 Protein Purification

Cell pellets were thawed on ice and resuspended in lysis buffer (50mM K-HEPES, pH 8.0, 250mM KCl, 0.1% Tween, 10mM imidazole, 0.5mM Mg-ATP, 1mM PMSF, 2mM benzimidazole, 1x complete protease inhibitor cocktail (Roche)). Cell suspensions were incubated with 1mg/ml lysozyme and 5mM β-mercaptoethanol on ice for 10 minutes, followed by sonication (3x 30sec). All subsequent procedures were performed at 4°C. Lysates were centrifuged at 40,000x g for 40 minutes. Supernatants were diluted and loaded onto an SP Sepharose column (Pharmacia, 5ml cartridge) in buffer A (50mM K-HEPES, pH 6.8, 1mM MgCl<sub>2</sub>, 1mM EGTA, 10µM Mg-ATP, 1mM DTT) and eluted with a 0 to 750mM KCl gradient in buffer A. Fractions containing KSP were pooled and incubated with Ni-NTA resin (Qiagen) for one hour. The resin was washed three times with buffer B (Lysis buffer minus PMSF and protease inhibitor cocktail), followed by three 15-minute incubations and washes with buffer B. Finally, the resin was incubated and washed for 15 minutes three times with buffer C (same as buffer B except for pH 6.0) and poured into a column. KSP was eluted with elution buffer (identical to buffer B except for 150mM KCl and 250mM imidazole). KSP-containing fractions were pooled, made 10% in sucrose, and stored at -80°C.

Microtubules are prepared from tubulin isolated from bovine brain. Purified tubulin (> 97% MAP-free) at 1 mg/ml is polymerized at 37°C in the presence of 10 µM paclitaxel, 1 mM DTT, 1 mM GTP in BRB80 buffer (80 mM K-PIPES, 1

mM EGTA, 1 mM MgCl<sub>2</sub> at pH 6.8). The resulting microtubules are separated from non-polymerized tubulin by ultracentrifugation and removal of the supernatant. The pellet, containing the microtubules, is gently resuspended in 10  $\mu$ M paclitaxel, 1 mM DTT, 50  $\mu$ g/ml ampicillin, and 5  $\mu$ g/ml chloramphenicol in BRB80.

- 5                   The kinesin motor domain is incubated with microtubules, 1 mM ATP (1:1 MgCl<sub>2</sub>: Na-ATP), and compound at 23°C in buffer containing 80 mM K-HEPES (pH 7.0), 1 mM EGTA, 1 mM DTT, 1 mM MgCl<sub>2</sub>, and 50 mM KCl. The reaction is terminated by a 2-10 fold dilution with a final buffer composition of 80 mM HEPES and 50 mM EDTA (or, alternately, with a 1:1 addition of reaction volume to stop
- 10   buffer(1.8M KCl and 50 mM EDTA)). Free phosphate from the ATP hydrolysis reaction is measured via a quinaldine red/ammonium molybdate assay by adding a 1.5 times volume of quench C (e.g., to a mixture of 40  $\mu$ l reaction volume + 40  $\mu$ l stop buffer is then added 120  $\mu$ l quench C). Quench A contains 0.1 mg/ml quinaldine red and 0.14% polyvinyl alcohol; quench B contains 12.3 mM ammonium molybdate
- 15   tetrahydrate in 1.15 M sulfuric acid. Quench C is a 2:1 ratio of quench A:quench B. The reaction is incubated for 5-10 minutes at 23°C, and the absorbance of the phospho-molybdate complex is measured at 540 nm.

- The compounds 1-6, 2-6 to 2-14, 3-2, 4-2 to 4-7, 5-4, 6-1 to 6-7, 7-2, 8-2, 9-1, 10-1 to 10-3, 11-1, 12-1, 13-1, 14-1 to 14-2, 15-1 to 15-86, 16-1 to 16-16,
- 20   17-5 to 17-13, 18-4, 18a-6, 18b-8, 18-5 to 18-16, 19-2 to 19-15, 20-10 to 20-11, 21-4 to 21-7, 22-1 to 22-8, 24-1 to 24-9, 25-2 to 25-16, 26-2 to 26-7, 27-2 to 27-7, 28-2 to 28-18, 29-5 to 29-14, 30-2 to 30-21, 31-2 to 31-3, 32-3 to 32-3, 33-1 to 33-6, 33a-2 to 34-11, 35-2 to 35-11, 36-2 to 36-6, 37-1 to 37-8, 38-1, 39-2 to 39-8, 40-2 to 40-8 and 41-2 to 41-4 in the Examples were tested in the above assay and found to have an
- 25   IC<sub>50</sub>  $\leq$  50 $\mu$ M.

## II. Cell Proliferation Assay

- Cells are plated in 96-well tissue culture dishes at densities that allow for logarithmic growth over the course of 24, 48, and 72 hours and allowed to adhere
- 30   overnight. The following day, compounds are added in a 10-point, one-half log titration to all plates. Each titration series is performed in triplicate, and a constant DMSO concentration of 0.1% is maintained throughout the assay. Controls of 0.1% DMSO alone are also included. Each compound dilution series is made in media without serum. The final concentration of serum in the assay is 5% in a 200  $\mu$ L

volume of media. Twenty microliters of Alamar blue staining reagent is added to each sample and control well on the titration plate at 24, 48, or 72 hours following the addition of drug and returned to incubation at 37°C. Alamar blue fluorescence is analyzed 6-12 hours later on a CytoFluor II plate reader using 530-560 nanometer wavelength excitation, 590 nanometer emission.

A cytotoxic EC<sub>50</sub> is derived by plotting compound concentration on the x-axis and average percent inhibition of cell growth for each titration point on the y-axis. Growth of cells in control wells that have been treated with vehicle alone is defined as 100% growth for the assay, and the growth of cells treated with compounds is compared to this value. Proprietary in-house software is used to calculate percent cytotoxicity values and inflection points using logistic 4-parameter curve fitting. Percent cytotoxicity is defined as:

$$\% \text{ cytotoxicity} = \frac{(\text{Fluorescence}_{\text{control}}) - (\text{Fluorescence}_{\text{sample}})}{(\text{Fluorescence}_{\text{control}})} \times 100$$
  
The inflection point is reported as the cytotoxic EC<sub>50</sub>.

### III. Evaluation of mitotic arrest and apoptosis by FACS

FACS analysis is used to evaluate the ability of a compound to arrest cells in mitosis and to induce apoptosis by measuring DNA content in a treated population of cells. Cells are seeded at a density of 1.4x10<sup>6</sup> cells per 6cm<sup>2</sup> tissue culture dish and allowed to adhere overnight. Cells are then treated with vehicle (0.1% DMSO) or a titration series of compound for 8-16 hours. Following treatment, cells are harvested by trypsinization at the indicated times and pelleted by centrifugation. Cell pellets are rinsed in PBS and fixed in 70% ethanol and stored at 4°C overnight or longer.

For FACS analysis, at least 500,000 fixed cells are pelleted and the 70% ethanol is removed by aspiration. Cells are then incubated for 30 min at 4°C with RNase A (50 Kunitz units/ml) and propidium iodide (50 µg/ml), and analyzed using a Becton Dickinson FACSCaliber. Data (from 10,000 cells) is analyzed using the Modfit cell cycle analysis modeling software (Verity Inc.).

An EC<sub>50</sub> for mitotic arrest is derived by plotting compound concentration on the x-axis and percentage of cells in the G2/M phase of the cell cycle for each titration point (as measured by propidium iodide fluorescence) on the y-axis. Data analysis is performed using the SigmaPlot program to calculate an inflection point using logistic 4-parameter curve fitting. The inflection point is reported as the EC<sub>50</sub> for mitotic arrest. A similar method is used to determine the compound EC<sub>50</sub>

for apoptosis. Here, the percentage of apoptotic cells at each titration point (as determined by propidium iodide fluorescence) is plotted on the y-axis, and a similar analysis is carried out as described above.

5 VI. Immunofluorescence Microscopy to Detect Monopolar Spindles


Methods for immunofluorescence staining of DNA, tubulin, and  
pericentrin are essentially as described in Kapoor *et al.* (2000) J. Cell Biol. 150: 975-  
988. For cell culture studies, cells are plated on tissue culture treated glass chamber  
slides and allowed to adhere overnight. Cells are then incubated with the compound  
10 of interest for 4 to 16 hours. After incubation is complete, media and drug are  
aspirated and the chamber and gasket are removed from the glass slide. Cells are then  
permeabilized, fixed, washed, and blocked for nonspecific antibody binding  
according to the referenced protocol. Paraffin-embedded tumor sections are  
deparaffinized with xylene and rehydrated through an ethanol series prior to blocking.  
15 Slides are incubated in primary antibodies (mouse monoclonal anti- $\alpha$ -tubulin  
antibody, clone DM1A from Sigma diluted 1:500; rabbit polyclonal anti-pericentrin  
antibody from Covance, diluted 1:2000) overnight at 4°C. After washing, slides are  
incubated with conjugated secondary antibodies (FITC-conjugated donkey anti-mouse  
IgG for tubulin; Texas red-conjugated donkey anti-rabbit IgG for pericentrin) diluted  
20 to 15 $\mu$ g/ml for one hour at room temperature. Slides are then washed and  
counterstained with Hoechst 33342 to visualize DNA. Immunostained samples are  
imaged with a 100x oil immersion objective on a Nikon epifluorescence microscope  
using Metamorph deconvolution and imaging software.

25

EXAMPLES

Examples provided are intended to assist in a further understanding of  
the invention. Particular materials employed, species and conditions are intended to  
be illustrative of the invention and not limiting of the reasonable scope thereof.

30

SCHEME 1

5

Step 1:      2-chloro-5-fluorobenzenediazonium tetrafluoroborate (1-1)

Nitrosonium tetrafluoroborate (802 mg, 6.87 mmol) was added to a solution of 2-chloro-5-fluoroaniline (1.00 g, 6.87 mmol) in acetonitrile (50 mL) at 0°C. The resulting mixture was stirred for 1 hour, then diluted with ethyl ether (150

mL). The precipitate was filtered and air-dried to give 2-chloro-5-fluorobenzenediazonium tetrafluoroborate (1-1) as an off-white solid. <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.66 (ddd, 1H, *J* = 6.7, 2.1, 1.0 Hz), 8.16 (m, 2H).

5    Step 2:            tert-butyl 3-(2-chloro-5-fluorophenyl)-2,3-dihydro-1H-pyrrole-1-carboxylate (1-2)

Palladium(II) acetate (24 mg, 0.11 mmol, 0.020 equiv) was added to a vigorously stirred, deoxygenated mixture of tert-butyl 2,5-dihydro-1H-pyrrole-1-carboxylate (900 mg, 5.32 mmol) and 2-chloro-5-fluorobenzenediazonium tetrafluoroborate (1-1, 1.30 g, 5.32 mmol) in water and carbon tetrachloride (1:1, 50 mL) at 23°C, and the resulting mixture was stirred for 20 hours. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution (150 mL) and ethyl acetate (2 x 150 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was dissolved in toluene (100 mL), and the resulting solution concentrated *in vacuo* to facilitate azeotropic removal of residual water. 2,6-Lutidine (1.24 mL, 10.6 mmol, 2.00 equiv) and trifluoroacetic anhydride (0.558 mL, 2.66 mmol, 0.500 equiv) were then sequentially added to a solution of the residue in toluene (100 mL) at 0°C. The resulting mixture was stirred at 0°C for 8 hours, then warmed to 23°C and stirred an additional 8 hours. The reaction mixture was heated at reflux for 1 hour, then cooled to 23°C and concentrated. The residue was partitioned between ethyl acetate (100 mL) and saturated aqueous sodium bicarbonate solution (100 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes initially, grading to 60% EtOAc in hexanes) to give tert-butyl 3-(2-chloro-5-fluorophenyl)-2,3-dihydro-1H-pyrrole-1-carboxylate (1-2) as an orange oil. LRMS *m/z* (M+H-CH<sub>3</sub>) 283.0 found, 283.1 required.

30    Step 3:            tert-butyl 4-(2-chloro-5-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (1-4)


Tris(dibenzylideneacetone)dipalladium(0) (20 mg, 0.022 mmol, 0.020 equiv) was added to a deoxygenated mixture of tert-butyl 3-(2-chloro-5-fluorophenyl)-2,3-dihydro-1H-pyrrole-1-carboxylate (1-2, 330 mg, 1.11 mmol, 1 equiv), benzenediazonium tetrafluoroborate (1-3, prepared from aniline by the method described for 1-1, 213 mg, 1.11 mmol, 1.00 equiv), and sodium acetate trihydrate (459 mg, 3.32 mmol, 3.00 equiv) in acetonitrile (20 mL) at 23°C. The reaction mixture

was stirred for 16 hours, then partitioned between saturated aqueous sodium bicarbonate solution (50 mL) and ethyl acetate (2 x 70 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes initially, grading to 40% hexanes in EtOAc) to provide tert-butyl 4-(2-chloro-5-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (1-4) as a dark orange oil. LRMS  $m/z$  (M+H-CH<sub>3</sub>) 359.0 found, 359.1 required.

Step 4:            4-(2-chloro-5-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (1-5)  
10                    Trifluoroacetic acid (10 mL) was added to a solution of tert-butyl 4-(2-chloro-5-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (1-4, 320 mg, 0.856 mmol, 1 equiv) in dichloromethane (40 mL) at 23°C, and the resulting mixture was stirred for 30 min, then concentrated to give 4-(2-chloro-5-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (1-5) as a TFA salt (brown oil). LRMS  $m/z$  (M+H)  
15                    274.1 found, 274.1 required.

Step 5:            4-(2-chloro-5-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (1-6)  
                         Triethylamine (0.600 mL, 4.28 mmol, 5.00 equiv) and  
20                    dimethylcarbonyl chloride (0.080 mL, 0.86 mmol, 1.00 equiv) were added to a solution of 4-(2-chloro-5-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (1-5, TFA salt, 0.856 mmol, 1 equiv) in dichloromethane (50 mL) at 23°C, and the resulting mixture was stirred for 2 hours, then concentrated. The residue was partitioned between saturated aqueous sodium bicarbonate solution (75 mL) and ethyl acetate  
25                    (100 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was suspended in ethyl ether (2 mL) and the precipitate filtered to provide 4-(2-chloro-5-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (1-6) as an off-white solid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.40-7.30 (m, 5H), 7.26 (m, 1H), 7.03 (dd, 1H, *J* = 9.0, 2.9 Hz), 6.97 (m, 1H), 6.24 (m, 1H), 6.15  
30                    (m, 1H), 4.86 (ddd, 1H, *J* = 13.9, 5.6, 2.2 Hz), 4.49 (dt, 1H, *J* = 13.9, 2.0 Hz), 2.86 (s, 6H). LRMS  $m/z$  (M+H) 345.0 found, 345.1 required.

## SCHEME 2





Step 1: 2,5-difluorobenzenediazonium tetrafluoroborate (2-1)

Nitrosonium tetrafluoroborate (905 mg, 7.75 mmol, 1.00 equiv) was added to a solution of 2,5-difluoroaniline (0.780 mL, 7.75 mmol, 1 equiv) in acetonitrile (50 mL) at 0°C. The resulting mixture was stirred for 1 hour, then diluted with ethyl ether (150 mL). The precipitate was filtered and air dried to give 2,5-difluorobenzenediazonium tetrafluoroborate (2-1) as a tan solid. <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.54 (m, 1H), 8.24 (m, 1H), 7.95 (m, 1H).

Step 2: tert-butyl 3-(2,5-difluorophenyl)-2,3-dihydro-1H-pyrrole-1-carboxylate (2-2)

Palladium(II) acetate (67 mg, 0.30 mmol, 0.020 equiv) was added to a vigorously stirred, deoxygenated mixture of tert-butyl 2,5-dihydro-1H-pyrrole-1-carboxylate (2.59 mL, 15.0 mmol, 1 equiv) and 2,5-difluorobenzenediazonium tetrafluoroborate (2-1, 3.42 g, 15.0 mmol, 1.00 equiv) in water and carbon tetrachloride (1:1, 150 mL) at 23°C, and the resulting mixture was stirred for 20 hours. The reaction mixture was concentrated, and the residue partitioned between ethyl acetate (300 mL) and saturated aqueous sodium bicarbonate solution (75 mL). The organic layer was washed with brine, then dried over sodium sulfate and concentrated. The residue was dissolved in toluene (200 mL), and the resulting solution concentrated *in vacuo* to facilitate azeotropic removal of residual water. 2,6-Lutidine (3.50 mL, 30.0 mmol, 2.00 equiv) and trifluoroacetic anhydride (1.48 mL, 10.5 mmol, 0.700 equiv) were then sequentially added to a solution of the residue in toluene (100 mL) at -10°C. The resulting mixture was allowed to warm to 10°C over 16 hours, then heated at reflux for 1 hour. The reaction mixture was allowed to cool to 23°C, then concentrated. The residue was partitioned between ethyl acetate (300 mL) and saturated aqueous sodium bicarbonate solution (150 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes initially, grading to 20% EtOAc in hexanes) to give tert-butyl 3-(2,5-difluorophenyl)-2,3-dihydro-1H-pyrrole-1-carboxylate (2-2) as a red oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) major rotamer: δ 7.03-6.84 (m, 3H), 6.70 (br s, 1H), 5.01 (br s, 1H), 4.42 (m, 1H), 4.13 (m, 1H), 3.60 (m, 1H), 1.50 (s, 9H).

Step 3: tert-butyl 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (2-3)

Tris(dibenzylideneacetone)dipalladium(0) (59 mg, 0.064 mmol, 0.020 equiv) was added to a deoxygenated mixture of tert-butyl 3-(2,5-difluorophenyl)-2,3-dihydro-1H-pyrrole-1-carboxylate (2-2, 900 mg, 3.20 mmol, 1 equiv), benzenediazonium tetrafluoroborate (1-3, prepared by the method described above for 1-1, 614 mg, 3.20 mmol, 1.00 equiv), and sodium acetate trihydrate (1.32 g, 9.60 mmol, 3.00 equiv) in acetonitrile (70 mL) at 23°C. The reaction mixture was stirred for 16 h, then partitioned between saturated aqueous sodium bicarbonate solution and ethyl acetate (2 x 70 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes initially, grading to 40% hexanes in EtOAc) to provide tert-butyl 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (2-3) as an orange oil. LRMS  $m/z$  (M+H-CH<sub>3</sub>) 343.0 found, 343.1 required.

Step 4: 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (2-4)

Trifluoroacetic acid (20 mL) was added to a solution of tert-butyl 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (2-3, 700 mg, 1.96 mmol, 1 equiv) in dichloromethane (50 mL) at 23°C, and the resulting mixture was stirred for 30 minutes, then concentrated to give 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (2-4) as a TFA salt (brown oil). LRMS  $m/z$  (M+H) 258.1 found, 258.1 required.

Step 5: 4-(2,5-difluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (2-5)



Triethylamine (1.37 mL, 9.79 mmol, 5.00 equiv) and dimethylcarbonyl chloride (0.180 mL, 1.96 mmol, 1.00 equiv) were added to a solution of 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (2-4, 1.96 mmol) in dichloromethane (50 mL) at 23°C, and the resulting mixture was stirred for 2 hours, then concentrated. The residue was partitioned between saturated aqueous sodium bicarbonate solution (75 mL) and ethyl acetate (100 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide 4-(2,5-difluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (2-5) as an off-white solid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.35-7.29 (m, 4H), 7.25 (m, 1H), 7.05 (m, 1H), 7.00




(m, 1H), 6.96 (m, 1H), 6.40 (br s, 1H), 6.13 (m, 1H), 4.88 (ddd, 1H,  $J = 13.7, 5.6, 2.0$  Hz), 4.52 (d, 1H,  $J = 13.7$  Hz), 2.88 (s, 6H). LRMS  $m/z$  (M+H) 329.1 found, 329.1 required.

- 5 Step 6: Enantiomers of 4-(2,5-difluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (2-6 and 2-7)
- Resolution of enantiomers of racemic 4-(2,5-difluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (2-5) by chiral normal-phase HPLC (Chiralcel OD column: 0.1 % diethylamine in 40% ethanol in hexanes)
- 10 provided in order of elution 2-6 (-) and 2-7 (+).

The following compounds were prepared by using appropriately substituted anilines in place of 2,5-difluoroaniline in Step 1 and simple modifications of the above procedure.


15

Cmpd	Structure	Name	$^1\text{H}$ NMR/LRMS $m/z$ (M+H)
2-8		4-(5-chloro-2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	$^1\text{H}$ NMR (500 MHz, $\text{CDCl}_3$ ) $\delta$ 7.36-7.20 (m, 7H), 7.04 (dd, 1H, $J = 10.7, 8.8$ Hz), 6.40 (br s, 1H), 6.13 (m, 1H), 4.87 (ddd, 1H, $J = 13.7, 5.6, 2.0$ Hz), 4.52 (d, 1H, $J = 13.7$ Hz), 2.88 (s, 6H). LRMS $m/z$ (M+H) 345.0 found, 345.1 required.
2-9		4-(2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	$^1\text{H}$ NMR (300 MHz, $\text{CDCl}_3$ ) $\delta$ 7.38-7.20 (m, 8H), 7.18-7.05 (m, 2H), 6.36 (br s, 1H), 6.12 (m, 1H), 4.92 (ddd, 1H, $J = 13.7, 5.6, 2.0$ Hz), 4.56 (d, 1H, $J = 13.7$ Hz), 2.88 (s, 6H). LRMS $m/z$ (M+H) 311.0 found, 311.1 required.

2-10		4-(2-fluoro-5-methylphenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	<sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 7.36-7.29 (m, 4H), 7.24 (m, 1H), 7.11 (dd, 1H, <i>J</i> = 7.3, 2.0 Hz), 7.05 (m, 1H), 6.97 (dd, 1H, <i>J</i> = 11.2, 8.3 Hz), 6.33 (br s, 1H), 6.11 (m, 1H), 4.89 (ddd, 1H, <i>J</i> = 13.7, 5.6, 2.0 Hz), 4.54 (d, 1H, <i>J</i> = 13.9 Hz), 2.88 (s, 6H), 2.32 (s, 3H). LRMS <i>m/z</i> (M+H) 325.0 found, 325.2 required.
2-11		4-(5-bromo-2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	<sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 7.42 (dd, 1H, <i>J</i> = 6.6, 2.4 Hz), 7.37 (ddd, 1H, <i>J</i> = 6.8, 4.4, 2.4 Hz), 7.34-7.23 (m, 5H), 6.99 (dd, 1H, <i>J</i> = 11.0, 8.8 Hz), 6.40 (br s, 1H), 6.13 (m, 1H), 4.87 (ddd, 1H, <i>J</i> = 13.7, 5.6, 2.0 Hz), 4.52 (d, 1H, <i>J</i> = 13.7 Hz), 2.88 (s, 6H). LRMS <i>m/z</i> (M+H) 390.9 found, 391.1 required.
2-12		4-{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}morpholine	<sup>1</sup> H NMR (500 MHz, CD <sub>3</sub> OD) δ 7.33 (m, 3H), 7.25 (m, 2H), 7.14 (m, 1H), 7.07 (m, 1H), 6.42 (br s, 1H), 6.07 (m, 1H), 4.99 (ddd, 1H, <i>J</i> = 13.9, 5.6, 2.0 Hz), 4.62 (br d, 1H, <i>J</i> = 13.9 Hz), 3.68 (m, 2H), 3.58 (m, 2H), 3.46 (m, 2H), 3.21 (m, 2H). LRMS <i>m/z</i> (M+H) 387.1 found, 387.1 required.

2-13		4-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)morpholine	<sup>1</sup> H NMR (500 MHz, CD <sub>3</sub> OD) δ 7.46 (dd, 1H, <i>J</i> = 8.8, 5.1 Hz), 7.35 (m, 3H), 7.27 (tt, 1H, <i>J</i> = 6.8, 2.2 Hz), 7.23 (dd, 1H, <i>J</i> = 8.8, 6.4, 3.2 Hz), 7.09 (dd, 1H, <i>J</i> = 4.9, 3.2 Hz), 6.26 (m, 1H), 6.08 (m, 1H), 4.97 (ddd, 1H, <i>J</i> = 13.9, 5.6, 2.0 Hz), 4.58 (dt, 1H, <i>J</i> = 10.3, 1.9 Hz), 3.65 (m, 2H), 3.57 (m, 2H), 3.43 (m, 2H), 3.19 (m, 2H). LRMS <i>m/z</i> (M+H) 371.0 found, 370.1 required.
2-14		N,N-dimethyl-2,4-diphenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	<sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 7.42 (br d, 2H, <i>J</i> = 7.6 Hz), 7.36 (br t, 2H, <i>J</i> = 7.1 Hz), 7.33-7.28 (m, 5H), 7.24 (m, 1H), 6.18 (m, 1H), 6.09 (m, 1H), 4.90 (ddd, 1H, <i>J</i> = 13.7, 5.6, 2.0 Hz), 4.52 (dt, 1H, <i>J</i> = 13.7, 2.0 Hz), 2.88 (s, 6H). LRMS <i>m/z</i> (M+H) 293.15 found, 293.16 required.

## SCHEME 3



Step 1: tert-butyl 3-[2-fluoro-5-(trifluoromethyl)phenyl]-2,3-dihydro-1H-pyrrole-1-carboxylate (3-1)


- 5 A deoxygenated solution of tert-butyl 2,5-dihydro-1H-pyrrole-1-carboxylate (3.38 g, 20.0 mmol, 10.0 equiv), 2-bromo-1-fluoro-4-(trifluoromethyl)benzene (486 mg, 2.00 mmol, 1 equiv), *N,N*-diisopropylethylamine (1.39 mL, 8.00 mmol, 4.00 equiv), tri-*o*-tolylphosphine (67 mg, 0.22 mmol, 0.11 equiv), palladium(II) acetate (22 mg, 0.10 mmol, 0.050 equiv), and silver carbonate
- 10 (386 mg, 1.40 mmol, 0.700 equiv) in DMF (8.0 mL) was heated under nitrogen at 100 °C for 20 h. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution (50 mL) and ethyl acetate (50 mL). The organic layer was washed with brine, dried over magnesium sulfate, and concentrated. The residue was purified by flash column chromatography (20% CH<sub>2</sub>Cl<sub>2</sub> in hexanes, grading to 50%
- 15 CH<sub>2</sub>Cl<sub>2</sub> in hexanes) to give tert-butyl 3-[2-fluoro-5-(trifluoromethyl)phenyl]-2,3-dihydro-1H-pyrrole-1-carboxylate (3-1) as a colorless gum. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) major rotamer: δ 7.51 (m, 2H), 7.15 (t, 1H, *J* = 9.1 Hz), 6.74 (br s, 1H), 5.03 (br s, 1H), 4.48 (m, 1H), 4.17 (m, 1H), 3.61 (m, 1H), 1.50 (s, 9H).

Step 2: 3-[2-fluoro-5-(trifluoromethyl)phenyl]-N,N-dimethyl-5-phenyl-2,3-dihydro-1H-pyrrole-1-carboxamide (3-2)

3-[2-fluoro-5-(trifluoromethyl)phenyl]-N,N-dimethyl-5-phenyl-2,3-dihydro-1H-pyrrole-1-carboxamide (3-2) was prepared from tert-butyl 3-[2-fluoro-5-(trifluoromethyl)phenyl]-2,3-dihydro-1H-pyrrole-1-carboxylate (3-1) following the method of Scheme 1. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.56 (m, 2H), 7.33 (m, 3H), 7.26 (m, 1H), 7.21 (dd, 1H, *J* = 10.7, 8.8 Hz), 6.46 (br s, 1H), 6.15 (m, 1H), 4.93 (ddd, 1H, *J* = 13.7, 5.6, 2.0 Hz), 4.58 (d, 1H, *J* = 13.7 Hz), 2.89 (s, 6H). LRMS *m/z* (M+H) 379.0 found, 379.1 required.

10

## SCHEME 4



Step 1: tert-butyl 4-(2,5-difluorophenyl)-2-(3-fluorophenyl)-2,3-dihydro-1H-pyrrole-1-carboxylate (4-1)

15

To a flame dried flask equipped with stir bar was added tert-butyl 3-(2,5-difluorophenyl)-2,3-dihydro-1H-pyrrole-1-carboxylate (2-2, 0.088 g, 0.31 mmol), 3-fluoro-1-iodobenzene (0.070 g, 0.31 mmol), triphenylarsine (0.038 g, 0.13 mmol), tributylamine (0.15 mL, 0.62 mmol) and anhydrous DMF (2.0 mL). The resulting




solution was purged, and filled with a nitrogen atmosphere. Palladium acetate (0.014 g, 0.06 mmol) was added in one portion, and the reaction was carefully returned to a nitrogen atmosphere. The reaction stirred under nitrogen for 24 h at 65 °C. The reaction was then diluted with EtOAc (10 mL) and poured into 5% aq. NaHCO<sub>3</sub> in a separatory funnel. The organic layer was washed with brine, separated and dried over magnesium sulfate. The organic layer was then filtered and concentrated to provide crude product. Purification by flash column chromatography (SiO<sub>2</sub>, 0-20% EtOAc/hexanes gradient) provided tert-butyl 4-(2,5-difluorophenyl)-2-(3-fluorophenyl)-2,5-dihydro-1H-pyrrole-1-carboxylate (4-1). LRMS *m/z* (M+H) 361.0 found, 361.1 required (M – 15). Further transformations followed those described in Scheme 1.



Step 2:            2-(3-fluorophenyl)-4-(2,5-difluorophenyl)- N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide (4-2)

Using the procedures described in Steps 4 and 5 of Scheme 1, but substituting intermediate 4-1 for intermediate 1-4 provided the title compound. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.29 (m, 1H), 7.14-6.90 (m, 6H), 6.37 (m, 1H), 6.14 (m, 1H), 4.86 (ddd, *J* = 13.7, 5.5, 1.2 Hz, 1H), 4.51 (d, *J* = 13.7 Hz, 1H), 2.91 (s, 6H). LRMS *m/z* (M+H) 347.0 found, 347.1 required.


The following compounds were prepared by simple modifications of the above procedures.



Cmpd	Structure	Name	<sup>1</sup> H NMR/LRMS <i>m/z</i> (M+H)
4-3		4-(2,5-difluorophenyl)-2-(4-fluorophenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide	<sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 7.33-7.29 (m, 2H), 7.09-6.94 (m, 5H), 6.37 (s, 1H), 6.13 (m, 1H), 4.85 (ddd, <i>J</i> = 13.5, 5.5, 2.0 Hz, 1H), 4.50 (d, <i>J</i> = 13.5 Hz, 1H), 2.88 (s, 6H). LRMS <i>m/z</i> (M+H) 347.0 found, 347.1 required.
4-4		4-(2,5-difluorophenyl)-2-(2-fluorophenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide	<sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 7.31-7.20 (m, 2H), 7.12-6.93 (m, 5H), 6.40 (s, 1H), 6.37 (m, 1H), 4.89 (ddd, <i>J</i> = 13.5, 5.5, 2.0 Hz, 1H), 4.56 (d, <i>J</i> = 13.5 Hz, 1H), 2.92 (s, 6H). LRMS <i>m/z</i> (M+H) 346.9 found, 347.1 required.
4-5		2-(3-bromophenyl)-4-(2,5-difluorophenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide	<sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 7.45 (s, 1H), 7.38 (d, <i>J</i> = 7.5 Hz, 1H), 7.29 (d, <i>J</i> = 7.5 Hz, 1H), 7.19 (t, <i>J</i> = 7.5 Hz, 1H), 7.09-7.04 (m, 1H), 7.02-6.95 (m, 2H), 6.36 (s, 1H), 6.13 (t, <i>J</i> = 3.0 Hz, 1H), 4.85 (ddd, <i>J</i> = 13.0, 5.5, 1.5 Hz, 1H), 4.51 (d, <i>J</i> = 13.5, 1H), 2.91 (s, 6H). LRMS <i>m/z</i> (M+H) 408.9 found,

			409.1 required.
4-6		2-(3-aminophenyl)-4-(2,5-difluorophenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide	<sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 8.06 (br s, 3H), 7.43 (s, 1H), 7.32 (t, <i>J</i> = 7.5 Hz, 1H), 7.27 (d, <i>J</i> = 4 Hz, 1H), 7.22 (d, <i>J</i> = 7.5 Hz, 1H), 7.07-7.02 (m, 1H), 6.98-6.94 (m, 2H), 6.31 (s, 1H), 6.07 (m, 1H), 4.86 (dd, <i>J</i> = 13.5, 4.0 Hz, 1H), 4.47 (d, <i>J</i> = 13.5 Hz, 1H), 2.82 (s, 6H). LRMS <i>m/z</i> (M+H) 344.0 found, 344.1 required.
4-7		4-(2,5-difluorophenyl)-2-(3-methylphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide	<sup>1</sup> H NMR (500 MHz, CDCl <sub>3</sub> ) δ 7.21 (t, <i>J</i> = 7.5 Hz, 1H), 7.12-6.94 (m, 6H), 6.38 (s, 1H), 6.08 (m, 1H), 4.90 (ddd, <i>J</i> = 14.0, 5.5, 2.0 Hz, 1H), 4.53 (d, <i>J</i> = 14.0 Hz, 1H), 2.89 (s, 6H), 2.33 (s, 3H). LRMS <i>m/z</i> (M+H) 343.1 found, 343.2 required.

## SCHEME 5



- 5 Step 1: (2S,4S)-tert-Butyl 4-hydroxy-2-phenylpyrrolidine-1-carboxylate (5-2)  
 To a flame dried flask equipped with stir bar was added tert-butyl  
 (2S,4S)-4-[[tert-butyl(dimethyl)silyl]oxy]-2-phenylpyrrolidine-1-carboxylate (5-1,  
 prepared from (S)-(-)-4-chloro-3-hydroxybutyronitrile by the method of Maeda, *et al*

*Synlett* 2001, 1808-1810, 7.8 g, 20.7 mmol) and anhydrous acetonitrile (20.0 mL). The resulting solution was treated with triethylamine trihydrofluoride (10.1 mL, 62.0 mmol) while stirring under N<sub>2</sub>. The reaction stirred 12 hours at 40°C. The reaction was then diluted with EtOAc (100 mL) and poured into 5% aq. NaHCO<sub>3</sub>. Following  
5 cessation of gas evolution, the organic layer was washed three addition times with 5% aq. NaHCO<sub>3</sub>. The organic layer was dried over magnesium sulfate, filtered and concentrated to provide crude product. Recrystallization was effected from EtOAc/hexanes to provide (2S,4S)-tert-butyl 4-hydroxy-2-phenylpyrrolidine-1-carboxylate (5-2) as a white crystalline solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) rotamers δ  
10 7.38-7.18 (m, 5H), 4.90 (m, 1H), 4.42 (m, 1H), 3.88 (m, 1H), 3.56 (dd, *J* = 11.5, 4.0 Hz, 1H), 2.60 (m, 1H), 2.03 (m, 1H), 1.50 and 1.20 (br s, 9H); MS 208.0 found, 208.1 (M – C(CH<sub>3</sub>)<sub>3</sub>) required.

Step 2: (2S)-tert-butyl 4-oxo-2-phenylpyrrolidine-1-carboxylate (5-3)  
15 To a flame dried flask equipped with stir bar was added 150 mL anhydrous dichloromethane which was cooled to –78°C. Oxalyl chloride (3.8 mL, 44 mmol) and DMSO (4.8 mL, 61 mmol) were added sequentially and the reaction stirred for 10 minutes. (2S,4S)-tert-Butyl 4-hydroxy-2-phenylpyrrolidine-1-carboxylate (5-2, 2.28 g, 8.73 mmol) in 10 mL anhydrous dichloromethane was added  
20 dropwise and stirred 1 hour at –78°C. Triethylamine (12 mL, 87mmol) was added and the reaction was warmed to 0°C over 1 hour. Upon completion, the reaction was washed with 5% NaHCO<sub>3</sub>, brine and dried over MgSO<sub>4</sub>. The organic layer was concentrated to provide crude (2S)-tert-butyl 4-oxo-2-phenylpyrrolidine-1-carboxylate (5-3). Recrystallization was effected with EtOAc/hexanes. <sup>1</sup>H NMR (300 MHz,  
25 CDCl<sub>3</sub>) δ 7.35 (m, 3H), 7.17 (m, 2H), 5.38 (m, 1H), 4.08 (d, *J* = 19.5 Hz, 1H), 3.90 (d, *J* = 19.3 Hz, 1H), 3.13 (dd, *J* = 18.8, 9.8 Hz, 1H), 2.58 (dd, *J* = 18.6, 2.4 Hz, 1H), 1.40 (br s, 9H); MS 206.0 found, 206.1 (M – C(CH<sub>3</sub>)<sub>3</sub>) required.

Step 3: (2S)-tert-butyl 2-phenyl-4-[[trifluoromethyl)sulfonyl]oxy}-2,5-dihydro-1H-pyrrole-1-carboxylate (5-4)  
30 To a flame dried flask equipped with stir bar was added ketone (2S)-tert-butyl 4-oxo-2-phenylpyrrolidine-1-carboxylate (5-3, 0.16 g, 0.62 mmol) and anhydrous THF (2 mL). The resulting solution was cooled to –78°C, and treated dropwise with lithium hexamethyldisilylamide (LHMDS, 0.68 mL, 1M in THF, 0.68

mmol). The reaction stirred 1 hour at  $-78^{\circ}\text{C}$ , and N-(5-chloropyridin-2-yl)-1,1,1-trifluoro-N-[(trifluoromethyl)sulfonyl]methanesulfonamide (0.27 g, 0.68 mmol) was added neat in one portion. The reaction was allowed to warm to  $0^{\circ}\text{C}$  and stirred 4 hours total. The reaction was diluted with  $\text{Et}_2\text{O}$  (10 mL) and washed successively with  
5  $\text{H}_2\text{O}$  (10 mL) and brine (10 mL). The organic layer was dried over  $\text{MgSO}_4$ , filtered and concentrated. The crude residue was purified by flash column chromatography (0-20%  $\text{EtOAc}$ /hexanes gradient, 15 min) to provide (2S)-tert-butyl 2-phenyl-4-  
10 {[(trifluoromethyl)sulfonyl]oxy}-2,5-dihydro-1H-pyrrole-1-carboxylate (5-4).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ) major rotamer:  $\delta$  7.30 (m, 5H), 5.72 (m, 1H), 5.48 (m, 1H), 4.42 (m, 2H), 1.18 (s, 9H); MS 379.0 found 379.1 ( $\text{M} - \text{CH}_3$ ) required.


Step 4: (2S)-4-(2,5-difluorophenyl)-2-phenyl-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide (2-6)

To a flame dried flask equipped with stir bar was added (2S)-tert-butyl  
15 2-phenyl-4-[(trifluoromethyl)sulfonyl]oxy}-2,5-dihydro-1H-pyrrole-1-carboxylate (5-4, 0.250 g, 0.636 mmol), 2,5-difluorophenyl boronic acid (0.251 g, 1.59 mmol),  $\text{Na}_2\text{CO}_3$  (0.202 g, 1.91 mmol), and  $\text{LiCl}$  (0.081 g, 1.91 mmol). The solids were dissolved in 20 mL 4:1  $\text{DME}/\text{H}_2\text{O}$  and degassed with nitrogen.  $\text{Pd}(\text{PPh}_3)_4$  (0.037 g, 0.032 mmol) was added and the reaction was sealed under nitrogen and heated to  
20  $90^{\circ}\text{C}$  for 2 hours. Upon completion, the reaction was partitioned between 5% aq.  $\text{NaHCO}_3$  and  $\text{EtOAc}$  (3 x 50 mL), and the combined organic layers were dried over  $\text{MgSO}_4$ . Following filtration, the organic layer was concentrated and purified via flash column chromatography ( $\text{SiO}_2$ , 0-20%  $\text{EtOAc}$ /hexanes gradient) to provide  
25 (2S)-tert-butyl 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (5-5). Further transformations followed those described in Scheme 1 to provide compound 2-6.

The following compounds were prepared by simple modifications of the above procedures.

Cmpd	Structure	Name	<sup>1</sup> H NMR/LRMS <i>m/z</i> (M+H)
5-6		(2S)-4-(5-chloro-2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	<sup>1</sup> H NMR (300 MHz, CDCl <sub>3</sub> ) δ 7.34-7.22 (m, 6H), 6.92-6.82 (m, 2H), 6.29 (m, 1H), 6.10 (m, 1H), 4.88 (ddd, 1H, <i>J</i> = 13.7, 5.6, 2.0 Hz), 4.53 (d, 1H, <i>J</i> = 13.7 Hz), 2.87 (s, 6H). LRMS <i>m/z</i> (M+H) 329.0 found, 329.1 required.

## SCHEME 6



5



4-(2,5-difluorophenyl)-1-(methanesulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole (6-1)




Methanesulfonyl chloride (40  $\mu$ L, 0.51 mmol, 1.00 equiv) was added to a solution of 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (2-4, 130 mg, 0.505 mmol, 1 equiv) and triethylamine (350  $\mu$ L, 2.52 mmol, 5.00 equiv) in dichloromethane (30 mL) at 23°C, and the resulting reaction mixture was stirred for 16 hours. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution (50 mL) and ethyl acetate (3 x 50 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide 4-(2,5-difluorophenyl)-1-(methanesulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole (6-1) as a tan solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.39-7.30 (m, 5H), 7.09 (td, 1H, *J* = 9.8, 4.6 Hz), 7.01 (m, 2H), 6.36 (br s, 1H), 5.76 (m, 1H), 4.84 (br d, 1H, *J* = 13.2 Hz), 4.64

(ddd, 1H,  $J = 13.2, 5.6, 2.0$  Hz), 2.55 (s, 3H). LRMS  $m/z$  (M+H) 336.0 found, 336.1 required.


The following compounds were prepared by simple modifications of the above procedure.

5


Cmpd	Structure	Name	$^1\text{H}$ NMR/LRMS $m/z$ (M+H)
6-2		4-(2,5-difluorophenyl)-1-(ethylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole	$^1\text{H}$ NMR (300 MHz, $\text{CDCl}_3$ ) $\delta$ 7.40-7.30 (m, 5H), 7.08 (td, 1H, $J = 9.8, 4.6$ Hz), 7.00 (m, 2H), 6.35 (br s, 1H), 5.81 (m, 1H), 4.95 (br d, 1H, $J = 13.2$ Hz), 4.62 (ddd, 1H, $J = 13.2, 5.6, 2.0$ Hz), 2.65 (m, 1H), 2.57 (m, 1H), 1.16 (t, 3H, $J = 7.3$ Hz) LRMS $m/z$ (M+H) 350.0 found, 350.1 required.
6-3		4-(2,5-difluorophenyl)-1-(propylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole	$^1\text{H}$ NMR (300 MHz, $\text{CDCl}_3$ ) $\delta$ 7.40-7.30 (m, 5H), 7.08 (td, 1H, $J = 9.8, 4.6$ Hz), 6.98 (m, 2H), 6.35 (br s, 1H), 5.80 (m, 1H), 4.92 (br d, 1H, $J = 13.2$ Hz), 4.61 (ddd, 1H, $J = 13.2, 5.6, 2.0$ Hz), 2.59 (m, 1H), 2.48 (m, 1H), 1.67 (m, 2H), 0.82 (t, 3H, $J = 7.3$ Hz) LRMS $m/z$ (M+H) 364.0 found, 364.1 required.

6-4		4-(2,5-difluorophenyl)-1-(isopropylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole	<sup>1</sup> H NMR (300 MHz, CDCl <sub>3</sub> ) δ 7.40-7.30 (m, 5H), 7.08 (td, 1H, <i>J</i> = 9.8, 4.6 Hz), 6.98 (m, 2H), 6.35 (br s, 1H), 5.87 (m, 1H), 5.01 (br d, 1H, <i>J</i> = 13.2 Hz), 4.58 (ddd, 1H, <i>J</i> = 13.2, 5.6, 2.0 Hz), 2.57 (pentet, 1H, <i>J</i> = 6.7 Hz), 1.18 (d, 3H, <i>J</i> = 7.0 Hz), 1.13 (d, 3H, <i>J</i> = 6.7 Hz) LRMS <i>m/z</i> (M+H) 364.0 found, 364.1 required.
6-5		4-(5-chloro-2-fluorophenyl)-1-(methylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole	<sup>1</sup> H NMR (500 MHz, CD <sub>3</sub> OD) δ 7.49 (dd, 1H, <i>J</i> = 8.8, 5.1 Hz), 7.45 (br d, 2H, <i>J</i> = 7.3 Hz), 7.38 (br t, 2H, <i>J</i> = 7.3 Hz), 7.32 (br t, 1H, <i>J</i> = 7.1 Hz), 7.26 (dd, 1H, <i>J</i> = 9.3, 2.9 Hz), 7.12 (td, 1H, <i>J</i> = 8.6, 2.9 Hz), 6.22 (m, 1H), 5.75 (m, 1H), 4.74 (dt, 1H, <i>J</i> = 14.2, 2.0 Hz), 4.68 (ddd, 1H, <i>J</i> = 14.2, 5.4, 2.0 Hz), 2.74 (s, 3H). LRMS <i>m/z</i> (M+H) 352.0 found, 352.05 required.
6-6		4-(2-fluoro-5-methylphenyl)-1-(isopropylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 360.1 found, 360.1 required.



6-7		4-(5-chloro-2-fluorophenyl)-1-(isopropylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole	<sup>1</sup> H NMR (500 MHz, CD <sub>3</sub> OD) $\delta$ 7.49 (dd, 1H, $J$ = 8.8, 5.1 Hz), 7.44 (br d, 2H, $J$ = 7.3 Hz), 7.39 (br t, 2H, $J$ = 7.3 Hz), 7.32 (br t, 1H, $J$ = 7.1 Hz), 7.24 (dd, 1H, $J$ = 9.3, 2.9 Hz), 7.11 (td, 1H, $J$ = 8.6, 2.9 Hz), 6.19 (m, 1H), 5.84 (m, 1H), 4.89 (dt, 1H, $J$ = 14.2, 2.0 Hz), 4.64 (ddd, 1H, $J$ = 14.2, 5.4, 2.0 Hz), 2.76 (m, 1H), 1.15 (d, 3H, $J$ = 6.8 Hz), 1.14 (d, 3H, $J$ = 6.8 Hz). LRMS $m/z$ (M+H) 379.9 found, 380.1 required.
-----	---	---	---

## SCHEME 7




5 2-({[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]sulfonyl}ethanamine (7-2))

A solution of hydrazine hydrate (20  $\mu$ L, 0.35 mmol, 4.0 equiv) and 2-({[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]sulfonyl}ethyl)-1H-indole-1,3(2H)-dione (7-1, prepared by the method of Scheme 6 from the

- corresponding sulfonyl chloride, 43 mg, 0.087 mmol, 1 equiv) in ethanol (5 mL) was heated at reflux for 5 hours. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution (50 mL) and ethyl acetate (3 x 50 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide 2-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]sulfonyl}ethanamine (7-2) as a TFA salt (white solid). <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.42 (m, 5H), 7.30-7.05 (br m, 3H), 6.42 (br s, 1H), 5.93 (m, 1H), 4.83 (m, 2H), 3.35-3.02 (br m, 4H). LRMS *m/z* (M+H) 365.0 found, 365.1 required.

10


SCHEME 8

- 15 Step 1: 4-(2,5-difluorophenyl)-2-phenyl-1-(vinylsulfonyl)-2,5-dihydro-1H-pyrrole (8-1)
- 
- Triethylamine (122 μL, 0.874 mmol, 5.00 equiv) and 2-chloroethanesulfonyl chloride (29 mg, 0.175 mmol, 1.00 equiv) were added sequentially to a solution of 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole

(2-4, 45 mg, 0.175 mmol, 1 equiv) in dichloromethane (10 mL) at 23°C, and the resulting mixture was stirred for 1 hour. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution (50 mL) and ethyl acetate (2 x 50 mL). The combined organic layers were dried over sodium sulfate and were  
 5 concentrated to provide 4-(2,5-difluorophenyl)-2-phenyl-1-(vinylsulfonyl)-2,5-dihydro-1H-pyrrole (8-1) as a brown oil. LRMS  $m/z$  (M+H) 348.0 found, 348.1 required.


Step 2: 2-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]sulfonyl}-N,N-dimethylethanamine (8-2)  
 10 Dimethylamine hydrochloride (7.0 mg, 0.086 mmol, 1.0 equiv) was added to a solution of 4-(2,5-difluorophenyl)-2-phenyl-1-(vinylsulfonyl)-2,5-dihydro-1H-pyrrole (8-1, 30 mg, 0.086 mmol, 1 equiv) and *N,N*-diisopropylethylamine (75  $\mu$ L, 0.43 mmol, 5.0 equiv) in dichloromethane (5 mL) at 23°C, and the resulting mixture  
 15 was stirred for 16 hours. Additional dimethylamine hydrochloride (21 mg, 0.26 mmol, 3.0 equiv) was added the and reaction mixture was stirred for 3 hours, then partitioned between saturated aqueous sodium bicarbonate solution (50 mL) and ethyl acetate (3 x 50 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC ( $H_2O/CH_3CN$  gradient  
 20 w/ 0.1 % TFA present) to provide 2-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]sulfonyl}-N,N-dimethylethanamine (8-2) as a TFA salt (white solid).  $^1H$  NMR (300 MHz,  $CD_3OD$ )  $\delta$  7.43 (m, 5H), 7.30-7.10 (br m, 3H), 6.42 (br s, 1H), 5.86 (m, 1H), 4.94-4.74 (br m, 2H), 3.46-3.14 (br m, 4H), 2.78 (s, 6H). LRMS  $m/z$  (M+H) 393.0 found, 393.1 required.

25

SCHEME 9

1-acetyl-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (9-1)

Acetic anhydride (44  $\mu$ L, 0.47 mmol, 2.0 equiv) and triethylamine (163  $\mu$ L, 1.17 mmol, 5.00 equiv) were added sequentially to a solution of 4-(2-chloro-5-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (2-4, 60 mg, 0.233 mmol, 1 equiv) in dichloromethane (10 mL) at 23°C, and the resulting mixture was stirred for 1 hour. The reaction mixture was concentrated and the residue purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide 1-acetyl-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (9-1) as a colorless oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) major rotamer:  $\delta$  7.40-7.24 (m, 5H), 7.11-6.94 (m, 3H), 6.38 (br s, 1H), 5.69 (br s, 1H), 4.91 (d, 1H,  $J$  = 15.9 Hz), 4.79 (dd, 1H,  $J$  = 15.9, 4.4 Hz), 1.89 (s, 3H). LRMS  $m/z$  (M+H) 300.0 found, 300.1 required.

SCHEME 104-(2-chloro-5-fluorophenyl)-1-pivaloyl-2-phenyl-2,5-dihydro-1H-pyrrole (10-1)


Triethylamine (149  $\mu$ L, 1.07 mmol, 10.00 equiv) and pivaloyl chloride (26  $\mu$ L, 0.214 mmol, 2.00 equiv) were added sequentially to a solution of 4-(2-chloro-5-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (1-5, 0.107 mmol, 1 equiv) in dichloromethane (10 mL) at 23°C, and the resulting mixture was stirred for 20 minutes. The reaction mixture was concentrated and the residue purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide 4-(2-chloro-5-fluorophenyl)-1-(2,2-dimethylpropanoyl)-2-phenyl-2,5-dihydro-1H-pyrrole (10-1) as a tan solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.39 (dd, 1H,  $J$  = 8.8, 5.2 Hz), 7.33 (m, 4H), 7.26 (m, 1H), 7.05 (dd, 1H,  $J$  = 9.2, 3.1 Hz), 6.99 (m, 1H), 6.20 (m, 1H), 5.98 (m, 1H), 5.00 (m, 2H), 1.28 (s, 9H). LRMS  $m/z$  (M+H) 358.0 found, 358.1 required.

The following compounds were prepared by simple modifications of the above procedure.

Cmpd	Structure	Name	<sup>1</sup> H NMR/LRMS <i>m/z</i> (M+H)
10-2		4-(2-chloro-5-fluorophenyl)-1-isobutyryl-2-phenyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 344.0 found, 344.1 required.
10-3		4-(2,5-difluorophenyl)-1-(2,2-dimethylpropanoyl)-2-phenyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 342.1 found, 342.2 required.

5

### SCHEME 11




10 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-ol (11-1)


*N,N*-Diisopropylethylamine (200  $\mu$ L, 1.2 mmol, 5.0 equiv) and 2-chloro-1,1-dimethyl-2-oxoethyl acetate (50  $\mu$ L, 0.35 mmol, 1.5 equiv) were added sequentially to a solution of 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole

- (2-4, 60 mg, 0.23 mmol, 1 equiv) in dichloromethane (12 mL) at 23°C, and the resulting mixture was stirred for 35 minutes. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution (40 mL) and dichloromethane (2 x 35 mL). The combined organic layers were dried over sodium sulfate and concentrated. Potassium hydroxide (65 mg, 1.2 mmol, 5.0 equiv) was added to a solution of the residue in 4:1 mixture of methanol and water (20 mL) and the resulting mixture was stirred at 23°C for 20 hours. The reaction mixture was partitioned between half-saturated aqueous sodium chloride solution (55 mL) and ethyl acetate (50 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-ol (11-1) as an off-white solid. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) δ 7.30 (m, 4H), 7.22 (m, 2H), 7.14 (m, 1H), 7.08 (m, 1H), 6.40 (m, 1H), 5.91 (m, 1H), 5.27 (d, 1H, *J* = 14.9 Hz), 5.19 (br d, 1H, *J* = 14.9 Hz), 1.43 (s, 3H), 1.41 (s, 3H).
- LRMS *m/z* (M+H) 344.1 found, 344.1 required.

The following compounds were prepared by simple modifications of the above procedure.

Cmpd	Structure	Name	<sup>1</sup> H NMR/LRMS <i>m/z</i> (M+H)
11-2		1-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-ol	<sup>1</sup> H NMR (500 MHz, CD <sub>3</sub> OD) δ 7.46 (dd, 1H, <i>J</i> = 8.8, 5.1 Hz), 7.36 (br d, 2H, <i>J</i> = 7.6 Hz), 7.32 (br t, 2H, <i>J</i> = 7.3 Hz), 7.22 (m, 2H), 7.09 (td, 1H, <i>J</i> = 8.6, 2.9 Hz), 6.21 (m, 1H), 5.91 (m, 1H), 5.24 (d, 1H, <i>J</i> = 14.9 Hz), 5.19 (ddd, 1H, <i>J</i> = 14.9, 5.6, 2.0 Hz), 1.42 (s, 3H), 1.38 (s, 3H). LRMS <i>m/z</i> (M+H) 360.0 found, 360.1 required.

## SCHEME 12




5

1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine (12-1)

A solution of N-(tert-butoxycarbonyl)-2-methylalanine (65 mg, 0.32 mmol, 1.5 equiv), 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (2-4, 55 mg, 0.21 mmol, 1 equiv), triethylamine (75  $\mu$ L, 0.54 mmol, 2.5 equiv), and PYBOP (145 mg, 0.279 mmol, 1.30 equiv) in DMF (15 mL) was stirred at 23°C for 20 hours. Additional triethylamine (75  $\mu$ L, 0.54 mmol, 2.5 equiv), and PYBOP (145 mg, 0.279 mmol, 1.30 equiv) were added and the resulting mixture stirred for 5 hours. The reaction mixture was partitioned between water (50 mL) and dichloromethane (2 x 50 mL). The combined organic layers were dried over sodium sulfate and concentrated. A solution of the residue in a 1:1 mixture (10 mL) of TFA and dichloromethane was stirred for 1.5 hours, then concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-

amine (12-1) as a TFA salt (light yellow oil).  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.28-6.98 (m, 8H), 6.33 (br s, 1H), 5.90 (m, 1H), 4.89 (m, 4H), 1.66 (s, 3H), 1.61 (s, 3H). LRMS  $m/z$  (M+H) 343.1 found, 343.15 required.

5

SCHEME 13

4-(2-fluoro-5-isocyanophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (13-1)

10 A deoxygenated solution of 4-(5-bromo-2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (2-11, 50 mg, 0.13 mmol, 1 equiv), zinc cyanide (10 mg, 0.85 mmol, 0.66 equiv), tris(dibenzylideneacetone)-dipalladium (12 mg, 0.013 mmol, 0.10 equiv), and 1,1'-bis(diphenylphosphino)-ferrocene (17 mg, 0.031 mmol, 0.24 equiv) in DMF (3 mL) was heated at 110°C for 3

15 hours. The reaction mixture was partitioned between brine (50 mL) and ethyl acetate (50 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography ( $\text{CH}_2\text{Cl}_2$  initially, grading to 10% EtOAc in  $\text{CH}_2\text{Cl}_2$ ) to provide 4-(2-fluoro-5-isocyanophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (13-1) as a pale yellow solid.  $^1\text{H}$  NMR


20 (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.62 (dd, 1H,  $j = 6.8, 2.0$  Hz), 7.59 (m, 1H), 7.33 (m, 4H), 7.26 (m, 1H), 7.22 (dd, 1H,  $J = 10.7, 8.8$  Hz), 6.49 (br s, 1H), 6.16 (m, 1H), 4.89 (ddd, 1H,



$J = 13.7, 5.6, 2.0$  Hz), 4.55 (d, 1H,  $J = 13.7$  Hz), 2.89 (s, 6H). LRMS  $m/z$  (M+H) 336.05 found, 336.14 required.

# SCHEME 14


5




## 4-(2,5-difluorophenyl)-2-phenyl-1-(trifluoroacetyl)-2,5-dihydro-1H-pyrrole (14-1)

- N,N*-Diisopropylethylamine (190  $\mu$ L, 1.1 mmol, 5.0 equiv) and dimethylphosphinic chloride (36 mg, 0.32 mmol, 1.5 equiv) were added sequentially to a solution of the TFA salt of 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (2-4, 55 mg, 0.21 mmol, 1 equiv) in dichloromethane (10 mL) at 23°C, and the resulting mixture was stirred for 20 hours. The reaction mixture was then partitioned between saturated aqueous sodium bicarbonate solution (40 mL) and dichloromethane (2 x 35 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC ( $H_2O/CH_3CN$  gradient w/ 0.1 % TFA present) to provide 4-(2,5-difluorophenyl)-2-phenyl-1-(trifluoroacetyl)-2,5-dihydro-1H-pyrrole (14-1) as an off-white solid.  $^1H$  NMR (500 MHz,  $CD_3OD$ ) major rotamer:  $\delta$  7.33 (m, 4H), 7.28 (m, 2H), 7.20 (m, 1H), 7.12 (m, 1H), 6.45 (m, 1H), 5.96 (m, 1H), 5.14 (br d, 1H,  $J = 14.9$  Hz), 5.05 (br d, 1H,  $J = 14.9$  Hz). LRMS  $m/z$  (M+H) 354.0 found, 354.1 required.

The following compounds were prepared by simple modifications of the above procedure.

Cmpd	Structure	Name	<sup>1</sup> H NMR/LRMS <i>m/z</i> (M+H)
14-2		4-(5-chloro-2-fluorophenyl)-2-phenyl-1-(trifluoroacetyl)-2,5-dihydro-1H-pyrrole	<sup>1</sup> H NMR (500 MHz, CD <sub>3</sub> OD) major rotamer: δ 7.49 (dd, 1H, <i>J</i> = 8.8, 5.1 Hz), 7.39 (br d, 2H, <i>J</i> = 7.3 Hz), 7.35 (br t, 2H, <i>J</i> = 7.3 Hz), 7.32 (br t, 1H, <i>J</i> = 7.1 Hz), 7.25 (dd, 1H, <i>J</i> = 9.3, 2.9 Hz), 7.11 (td, 1H, <i>J</i> = 8.6, 2.9 Hz), 6.25 (m, 1H), 5.96 (m, 1H), 5.12 (br d, 1H, <i>J</i> = 14.9 Hz), 4.99 (br d, 1H, <i>J</i> = 14.9 Hz). LRMS <i>m/z</i> (M+H) 370.0 found, 370.05 required.





## SCHEME 15















- 5 (1S)-1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2-methylpropylamine (15-1)
- N-(tert-butoxycarbonyl)-L-valine (101 mg, 0.464 mmol, 2.00 equiv), PYBOP (243 mg, 0.467 mmol, 2.00 equiv), and triethylamine (0.163 mL, 1.17 mmol, 5.00 equiv) were added to a solution of 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole (2-4, 0.233 mmol, 1 equiv) in DMF (10 mL) at 23°C, and the resulting mixture was stirred for 20 hours. The reaction mixture was partitioned between a 3:1 mixture of saturated aqueous sodium bicarbonate solution and brine (50 mL) and ethyl acetate (50 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was dissolved in a 1:1 mixture of trifluoroacetic acid and dichloromethane (10 mL), and the resulting solution was allowed to stand for 45 minutes, then concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide (1S)-1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2-methylpropylamine (15-1) as a 1:1 mixture of diastereomers (tan gum) (isolated as the TFA salt). LRMS *m/z* (M+H) 357.2 found, 357.2 required.







The following compounds were prepared by simple modifications of the above procedure. Unless otherwise indicated, the compounds in the table were isolated as the TFA salt.






5

Cmpd	Structure	Name	LRMS $m/z$ (M+H)
15-2		(1R)-1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2-methylpropylamine	LRMS $m/z$ (M+H) 357.2 found, 357.2 required.
15-3		4-(2,5-difluorophenyl)-2-phenyl-1-L-prolyl-2,5-dihydro-1H-pyrrole	LRMS $m/z$ (M+H) 355.2 found, 355.2 required
15-4		4-(2,5-difluorophenyl)-2-phenyl-1-D-prolyl-2,5-dihydro-1H-pyrrole	LRMS $m/z$ (M+H) 355.2 found, 355.2 required
15-5		(4R)-4-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-1,3-thiazolidine	LRMS $m/z$ (M+H) 373.1 found, 373.1 required

15-6		methyl (3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutanoate	LRMS <i>m/z</i> (M+H) 387.2 found, 387.1 required
15-7		(4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanamide	LRMS <i>m/z</i> (M+H) 386.1 found, 386.2 required
15-8		(1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-(methylthio)propylamine	LRMS <i>m/z</i> (M+H) 389.2 found, 389.1 required
15-9		(1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-(methylsulfonyl)propylamine	LRMS <i>m/z</i> (M+H) 421.1 found, 421.1 required
15-10		(2S)-2-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}piperidine	LRMS <i>m/z</i> (M+H) 369.2 found, 369.2 required
15-11		(1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}pentylamine	LRMS <i>m/z</i> (M+H) 371.2 found, 371.2 required







15-12		(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(thien-2-ylmethyl)ethylamine	LRMS <i>m/z</i> (M+H) 411.1 found, 411.1 required
15-13		4-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-1,1-dioxidotetrahydro-2H-thiopyran-4-ylamine	LRMS <i>m/z</i> (M+H) 433.1 found, 433.1 required
15-14		(2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methyl-1-oxopropan-2-amine	LRMS <i>m/z</i> (M+H) 343.1 found, 343.2 required
15-15		(1S)-1-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]propylamine	LRMS <i>m/z</i> (M+H) 343.1 found, 343.2 required
15-16		(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-phenylethanamine	LRMS <i>m/z</i> (M+H) 391.0 found, 391.2 required
15-17		2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methyl-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 329.1 found, 329.1 required






15-18		3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-amine	LRMS <i>m/z</i> (M+H) 329.1 found, 329.1 required
15-19		(1S,2S)-1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2-methylbutylamine	LRMS <i>m/z</i> (M+H) 371.1 found, 371.2 required
15-20		(1S)-1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)butylamine	LRMS <i>m/z</i> (M+H) 357.1 found, 357.2 required
15-21		(1S)-1-cyclopropyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 355.1 found, 355.2 required
15-22		1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)cyclopropanamine	LRMS <i>m/z</i> (M+H) 340.9 found, 341.1 required
15-23		1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-oxopropan-2-amine	LRMS <i>m/z</i> (M+H) 329.2 found, 329.1 required

15-24		(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-methyl-2-oxoethylamine	LRMS $m/z$ (M+H) 329.1 found, 329.1 required
15-25		(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(pyridin-2-ylmethyl)ethylamine	LRMS $m/z$ (M+H) 406.1 found, 406.2 required
15-26		(1S)-1-cyclohexyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine	LRMS $m/z$ (M+H) 397.2 found, 397.2 required
15-27		(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(4-iodobenzyl)-2-oxoethylamine	LRMS $m/z$ (M+H) 531.0 found, 531.1 required
15-28		(1S)-1-benzyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine	LRMS $m/z$ (M+H) 405.2 found, 405.2 required








15-29		4-{(2S)-2-amino-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropyl}phenol	LRMS <i>m/z</i> (M+H) 421.1 found, 421.2 required
15-30		(3S)-3-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-1,2,3,4-tetrahydroisoquinoline	LRMS <i>m/z</i> (M+H) 417.2 found, 417.2 required
15-31		(1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-phenylpropylamine	LRMS <i>m/z</i> (M+H) 419.2 found, 419.2 required
15-32		(1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-methylbutylamine	LRMS <i>m/z</i> (M+H) 371.2 found, 371.2 required
15-33		(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(pyridin-3-ylmethyl)ethylamine	LRMS <i>m/z</i> (M+H) 406.1 found, 406.2 required






15-34		1-[(2S)-azetidin-2-ylcarbonyl]-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 341.1 found, 341.1 required
15-35		(3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutanamide	LRMS <i>m/z</i> (M+H) 372.1 found, 372.1 required
15-36		4-(2,5-difluorophenyl)-1-[(2S)-2,5-dihydro-1H-pyrrol-2-ylcarbonyl]-2-phenyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 353.1 found, 353.1 required
15-37		4-(2,5-difluorophenyl)-1-[(2-methylazetidin-2-yl)carbonyl]-2-phenyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 355.1 found, 355.2 required
15-38		(1S)-1- {[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylamine	LRMS <i>m/z</i> (M+H) 371.2 found, 371.2 required
15-39		methyl (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanoate	LRMS <i>m/z</i> (M+H) 401.1 found, 401.2 required





15-40		4-(2,5-difluorophenyl)-2-phenyl-1-([(2S,3S)-2-phenylpyrrolidin-3-yl]carbonyl)-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 431.2 found, 431.2 required
15-41		4-(2,5-difluorophenyl)-2-phenyl-1-[(5-phenylpyrrolidin-3-yl)carbonyl]-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 431.2 found, 431.2 required
15-42		(2S)-2-amino-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-ol	LRMS <i>m/z</i> (M+H) 345.1 found, 345.1 required
15-43		(2R,3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutan-2-ol	LRMS <i>m/z</i> (M+H) 359.1 found, 359.2 required
15-44		(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(methoxymethyl)-2-oxoethylamine	LRMS <i>m/z</i> (M+H) 359.1 found, 359.2 required

15-45		4-(2,5-difluorophenyl)-2-phenyl-1-(pyrrolidin-3-ylcarbonyl)-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 355.1 found, 355.2 required
15-46		4-(2,5-difluorophenyl)-2-phenyl-1-[(3-phenylpyrrolidin-3-yl)acetyl]-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 445.1 found, 445.2 required
15-47		(1S)-1-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-3,3-difluoropropylamine	LRMS <i>m/z</i> (M+H) 379.1 found, 379.1 required
15-48		(1S)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxo-1-phenylpropan-1-amine	LRMS <i>m/z</i> (M+H) 405.2 found, 405.2 required
15-49		4-(2,5-difluorophenyl)-2-phenyl-1-[(4S)-4-phenyl-L-prolyl]-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 431.3 found, 431.2 required






15-50		1-{2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-cyclohexanamine	LRMS <i>m/z</i> (M+H) 397.0 found, 397.2 required
15-51		2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 315.0 found, 315.1 required
15-52		4-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}piperidin-4-amine	LRMS <i>m/z</i> (M+H) 384.1 found, 384.2 required
15-53		(1S,3R)-3-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopentanamine	LRMS <i>m/z</i> (M+H) 369.1 found, 369.2 required
15-54		(1R,4S)-4-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-2-en-1-amine	LRMS <i>m/z</i> (M+H) 367.1 found, 367.2 required
15-55		(1S,4R)-4-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-2-	LRMS <i>m/z</i> (M+H) 367.1 found, 367.2 required






		en-1-amine	
15-56		(1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-ynylamine	LRMS <i>m/z</i> (M+H) 353.1 found, 353.1 required
15-57		(1R)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxo-1-phenylpropan-1-amine	LRMS <i>m/z</i> (M+H) 405.1 found, 405.2 required
15-58		3-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-phenylpiperidine	LRMS <i>m/z</i> (M+H) 445.1 found, 445.2 required
15-59		(1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-enylamine	LRMS <i>m/z</i> (M+H) 355.1 found, 355.2 required
15-60		(2S)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-(methylamino)-3-oxopropan-1-ol	LRMS <i>m/z</i> (M+H) 359.1 found, 359.2 required






15-61		(3R,5S)-5-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)pyrrolidin-3-ol	LRMS <i>m/z</i> (M+H) 371.1 found, 371.2 required
15-62		(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(1,3-thiazol-4-ylmethyl)ethylamine	LRMS <i>m/z</i> (M+H) 412.0 found, 412.1 required
15-63		(1R)-1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)but-3-enylamine	LRMS <i>m/z</i> (M+H) 355.1 found, 355.2 required
15-64		(2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,3-dimethyl-1-oxobutan-2-amine	LRMS <i>m/z</i> (M+H) 371.0 found, 371.2 required
15-65		(2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,4-dimethyl-1-oxopentan-2-amine	LRMS <i>m/z</i> (M+H) 385.1 found, 385.2 required


15-66		(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-[(1-methyl-1H-imidazol-4-yl)methyl]-2-oxoethylamine	LRMS <i>m/z</i> (M+H) 409.2 found, 409.2 required
15-67		4-(2,5-difluorophenyl)-1-(N~6~-formyl-L-lysyl)-2-phenyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 414.2 found, 414.2 required
15-68		(2S,3S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,3-dimethyl-1-oxopentan-2-amine	LRMS <i>m/z</i> (M+H) 385.2 found, 385.2 required
15-69		(1S)-1-(cyclohexylmethyl)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine	LRMS <i>m/z</i> (M+H) 411.2 found, 411.2 required




15-70		(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxoethylamine	LRMS $m/z$ (M+H) 444.1 found, 444.2 required
15-71		(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(isocyanomethyl)-2-oxoethylamine	LRMS $m/z$ (M+H) 354.1 found, 354.1 required
15-72		(1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3,3-dimethylbutylamine	LRMS $m/z$ (M+H) 385.0 found, 385.2 required
15-73		1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,3-dimethyl-1-oxobutan-2-amine	LRMS $m/z$ (M+H) 371.0 found, 371.2 required
15-74		1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclohexanamine	LRMS $m/z$ (M+H) 383.1 found, 383.2 required

15-75		1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)cyclopentylamine	LRMS <i>m/z</i> (M+H) 369.1 found, 369.2 required
15-76		(1S)-3-(benzyloxy)-1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)propylamine	LRMS <i>m/z</i> (M+H) 449.1 found, 449.2 required
15-77		1-([4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)cyclopent-3-en-1-amine	LRMS <i>m/z</i> (M+H) 367.0 found, 367.2 required
15-78		(1S)-1-cyclopentyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 383.1 found, 383.2 required
15-79		4-(2,5-difluorophenyl)-1-(2-methylpropyl)-2-phenyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 369.1 found, 369.2 required

15-80		1-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine; isolated as the free base	LRMS <i>m/z</i> (M+H) 359.0 found, 359.1 required
15-81		(1S)-1-([4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2-methylpropylamine; isolated as the free base	LRMS <i>m/z</i> (M+H) 373.0 found, 373.1 required
15-82		(1S)-2-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-cyclopropyl-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 371.1 found, 371.1 required
15-83		(1S,2S)-1-([4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2-methylbutylamine	LRMS <i>m/z</i> (M+H) 387.0 found, 387.2 required
15-84		(1S)-1-([4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-pentylamine; isolated as the free base	LRMS <i>m/z</i> (M+H) 387.0 found, 387.2 required

<p>15-85</p> 	<p>(1S)-1-{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3,3-dimethylbutylamine; isoalted as the free base</p>	<p>LRMS <i>m/z</i> (M+H) 401.0 found, 401.2 required</p>
--	---	--

**SCHEME 16**










(1S)-1-[[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-2,2-dimethylpropylamine (16-1)






A solution of tert-butyl (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (5-5, 400 mg, 1.12 mmol, 1 equiv) in a 4:1 mixture of dichloromethane and trifluoroacetic acid (20 mL) was stirred for 30 minutes, then concentrated. The residue was dried via toluene azeotrope (2 x 10 mL), then dissolved in DMF (5 mL). Triethylamine (0.780 mL, 5.60 mmol, 5.00 equiv), N-(tert-butoxycarbonyl)-3-methyl-L-valine (388 mg, 1.68 mmol, 1.50 equiv), and PYBOP (874 mg, 1.68 mmol, 1.50 equiv) were added, and the resulting mixture was stirred at 23°C for 24 hours. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution and ethyl acetate (2 x 50 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes initially, grading to 40% ethyl acetate in





hexanes) to yield the desired coupled, N-Boc-protected intermediate, which was dissolved in a 2:1 mixture of dichloromethane and trifluoroacetic acid. The resulting solution was allowed to stand for 30 minutes, then concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present). The desired fractions were partitioned between saturated aqueous sodium bicarbonate solution and ethyl acetate (2 x 50 mL), then the combined organic layers were dried over sodium sulfate and concentrated to provide (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylamine (16-1) as a white solid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) major rotamer: δ 7.40 (t, 2H, *J* = 7.6 Hz), 7.31 (m, 1H), 7.26 (d, 2H, *J* = 7.8 Hz), 7.11-6.93 (m, 3H), 6.38 (s, 1H), 5.81 (m, 1H), 4.88 (m, 2H), 3.03 (s, 1H), 1.00 (s, 9H). LRMS *m/z* (M+H) 371.0 found, 371.2 required.

The following compounds were prepared by simple modifications of the above procedures. Unless noted otherwise, the compounds in the table were isolated as the free base.


Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
16-2		(1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylpropylamine	LRMS <i>m/z</i> (M+H) 357.1 found, 357.2 required.
16-3		(1S)-1-cyclohexyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 397.1 found, 397.2 required

16-4		(1S)-1-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)but-3-enylamine; isolated as the TFA salt	LRMS $m/z$ (M+H) 355.0 found, 355.2 required
16-5		(1S)-1-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)but-3-ynylamine; isolated as the TFA salt	LRMS $m/z$ (M+H) 353.0 found, 353.1 required
16-6		(1S)-1-benzyl-2-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine	LRMS $m/z$ (M+H) 405.0 found, 405.2 required
16-7		(1S)-1-cyclopropyl-2-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine; isolated as the TFA salt	LRMS $m/z$ (M+H) 355.0 found, 355.2 required
16-8		1-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine	LRMS $m/z$ (M+H) 343.0 found, 343.2 required


16-9		(1S)-1-([(2S)-4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2,2-dimethylpropylamine	LRMS <i>m/z</i> (M+H) 387.0 found, 387.2 required
16-10		(1S)-1-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-pentylamine	LRMS <i>m/z</i> (M+H) 371.1 found, 371.2 required
16-11		(1S)-1-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-3-methylbutylamine	LRMS <i>m/z</i> (M+H) 371.1 found, 371.2 required
16-12		(1S)-1-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-3,3-dimethylbutylamine	LRMS <i>m/z</i> (M+H) 385.1 found, 385.2 required
16-13		1-cyclopropyl-3-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-amine; isolated as the TFA salt	LRMS <i>m/z</i> (M+H) 369.0 found, 369.2 required

16-14		(1S)-2-[(2S)-4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-cyclopropyl-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 371.0 found, 371.1 required
16-15		(1S)-1-[(2S)-4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylpropylamine	LRMS <i>m/z</i> (M+H) 372.9 found, 373.1 required
16-16		(1S,2S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylbutylamine; isolated as the TFA salt	LRMS <i>m/z</i> (M+H) 371.1 found, 371.2 required
16-17		(2S)-4-(2,5-difluorophenyl)-1-[(methylsulfonyl)acetyl]-2-phenyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 378.3 found, 378.1 required.



16-18		(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-1-[(phenylsulfonyl)acetyl]-2,5-dihydro-1 <i>H</i> -pyrrole	LRMS <i>m/z</i> (M+H) 440.4 found, 440.1 required.
-------	---	--	--

## SCHEME 17



Step 1:        tert-butyl 3-(5-chloro-2-fluorophenyl)-2,3-dihydro-1H-pyrrole-1-carboxylate (17-1)

Palladium(II) acetate (106 mg, 0.473 mmol, 0.020 equiv) was added to a vigorously stirred, deoxygenated mixture of tert-butyl 2,5-dihydro-1H-pyrrole-1-carboxylate (4.00 g, 23.6 mmol, 1 equiv) and 5-chloro-2-fluorobenzenediazonium tetrafluoroborate (10.4 g, 42.5 mmol, 1.80 equiv) in water and carbon tetrachloride (1:1, 150 mL) at 23°C, and the resulting mixture was stirred for 20 hours. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution (200 mL) and dichloromethane (2 x 200 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was dissolved in toluene (200 mL), and the resulting solution concentrated *in vacuo* to facilitate azeotropic removal of residual water. 2,6-Lutidine (5.51 mL, 47.3 mmol, 2.00 equiv) and trifluoroacetic anhydride (1.67 mL, 11.8 mmol, 0.500 equiv) were then sequentially added to a solution of the residue in toluene (150 mL) at -20°C. The resulting mixture was kept at -20°C for 5 hours, then allowed to slowly warm to 23°C. After 16 hours at 23°C, the reaction mixture was heated at reflux for 30 minutes. The reaction mixture was allowed to cool to 23°C, then partitioned between ethyl acetate (100 mL) and saturated aqueous sodium bicarbonate solution (100 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes initially, grading to 30% EtOAc in hexanes) to give tert-butyl 3-(5-chloro-2-fluorophenyl)-2,3-dihydro-1H-pyrrole-1-carboxylate (17-1) as an orange oil. LRMS *m/z* (M+H-CH<sub>3</sub>) 283.0 found, 283.1 required.

Step 2:        tert-butyl 2-(3-([tert-butyl(dimethyl)silyl]oxy)phenyl)-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrole-1-carboxylate (17-2)

A deoxygenated mixture of tert-butyl 3-(5-chloro-2-fluorophenyl)-2,3-dihydro-1H-pyrrole-1-carboxylate (17-1, 1.100 g, 3.69 mmol, 1 equiv), tert-butyl(3-iodophenoxy)dimethylsilane (1.85 g, 5.54 mmol, 1.50 equiv), tributylamine (1.76 mL, 7.39 mmol, 2.00 equiv), triphenylarsine (0.453 g, 1.48 mmol, 0.400 equiv), and palladium acetate (0.166 g, 0.739 mmol, 0.200 equiv) in DMF (10 mL) was heated at 65°C for 20 hours. The reaction mixture was partitioned between water (100 mL) and ethyl acetate (2 x 85 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography (100% hexanes initially, grading to 50% ethyl acetate in hexanes) to provide tert-butyl

2-(3-([tert-butyl(dimethyl)silyl]oxy)phenyl)-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrole-1-carboxylate (17-2) as an orange oil. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) major rotamer: δ 7.11-6.53 (m, 7H), 6.11 (s, 1H), 5.32 (m, 1H), 4.53 (m, 2H), 1.09 (s, 9H), 0.79 (s, 9H), 0.00 (s, 6H). LRMS *m/z* (M+H) 504.1 found, 504.2 required.

5

Step 3: tert-butyl 2-[2-(3-([tert-butyl(dimethyl)silyl]oxy)phenyl)-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrol-1-yl]-1,1-dimethyl-2-oxoethylcarbamate (17-3)

A solution of tert-butyl 2-(3-([tert-butyl(dimethyl)silyl]oxy)phenyl)-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrole-1-carboxylate (17-2, 276 mg, 0.683 mmol, 1 equiv), triethylamine (277 mg, 2.73 mmol, 4 equiv), PYBOP (711 mg, 1.37 mmol, 2 equiv) and 2-[(tert-butoxycarbonyl)amino]-2-methylpropanoic acid (278 mg, 1.37 mmol, 2 equiv) in DMF (10 mL) was warmed at 60°C for 18 hours. The solution was concentrated and the residue was partitioned between ethyl acetate and saturated aqueous sodium bicarbonate solution. The organic layer was washed with brine, dried over magnesium sulfate and concentrated. The residue was purified by flash column chromatography. Elution with 10% ethyl acetate/hexanes to 20% ethyl acetate/hexanes gave tert-butyl 2-[2-(3-([tert-butyl(dimethyl)silyl]oxy)phenyl)-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrol-1-yl]-1,1-dimethyl-2-oxoethylcarbamate (17-3) as a pale yellow gum. LRMS *m/z* (M+H) 588.9 found, 589.0 required.



Step 4: 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole (17-4)






A solution of tert-butyl 2-[2-(3-([tert-butyl(dimethyl)silyl]oxy)phenyl)-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrol-1-yl]-1,1-dimethyl-2-oxoethylcarbamate (17-3, 247 mg, 0.419 mmol, 1 equiv) and trifluoroacetic acid (4 mL) in dichloromethane (10 mL) was stirred under ambient conditions for 1 hour. The reaction was concentrated at 23°C and the residue was partitioned between ethyl acetate and saturated sodium bicarbonate solution. The organic layer was washed with brine, dried over magnesium sulfate and concentrated to yield a yellow gum. A solution of the gum in methanol (5 mL) was treated with potassium carbonate (290 mg, 2.10 mmol, 5.00 equiv) and water (1 mL). The reaction mixture was stirred under ambient conditions for 2 hours, then concentrated. The aqueous solution was acidified with aqueous 1 N HCl solution to pH 8 and the mixture was extracted with ethyl






acetate. The organic layer was washed with brine, dried over magnesium sulfate and concentrated to provide 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole (17-4) as a cream-colored solid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.30 (dd, 1H, *J* = 6.6, 2.7 Hz), 7.30 (m, 1H), 7.18 (t, 1H, *J* = 7.8 Hz), 7.05 (t, 1H, *J* = 9.2 Hz), 6.83 (br d, 1H, *J* = 7.6 Hz), 6.78 (br s, 1H), 6.70 (dd, 1H, *J* = 8.1, 1.8 Hz), 6.38 (s, 1H), 5.92 (br s, 1H), 5.30 (m, 1H), 5.08 (m, 1H), 1.46 (s, 6H). LRMS *m/z* (M+H) 375.0 found, 375.0 required.





**Step 5:** (2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole (17-5)  
Resolution of enantiomers of racemic 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole (17-4) by chiral normal-phase HPLC (Chiralcel OD column: 0.1% diethylamine in 10% ethanol in hexanes initially, grading to 30% ethanol in hexanes) provided (2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole (17-5) as the first eluting, minus (-) enantiomer.

The following compounds were prepared by simple modifications of the above procedures. Unless otherwise indicated, the compounds in the table were isolated as the free base.

Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
17-6		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole; isolated as the TFA salt	LRMS <i>m/z</i> (M+H) 373.0 found, 373.2 required.
17-7		4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 389.0 found, 389.1 required


17-8		(2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole	LRMS $m/z$ (M+H) 389.0 found, 389.1 required
17-9		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole; isolated as the TFA salt	LRMS $m/z$ (M+H) 359.0 found, 359.2 required
17-10		3-[1-[(2S)-2-amino-2-cyclopropylethanoyl]-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol; isolated as the TFA salt	LRMS $m/z$ (M+H) 386.9 found, 387.1 required
17-11		4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-isoleucyl-2,5-dihydro-1H-pyrrole; isolated as the TFA salt	LRMS $m/z$ (M+H) 403.0 found, 403.2 required
17-12		4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-norleucyl-2,5-dihydro-1H-pyrrole	LRMS $m/z$ (M+H) 403.0 found, 403.2 required

17-13		(2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(3-methyl-L-valyl)-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 403.0 found, 403.2 required
17-14		(2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 373.0 found, 373.2 required
17-15		3-[(2S)-1-[(2S)-2-amino-2-cyclopropylethanoyl]-4-(2,5-difluorophenyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol	LRMS <i>m/z</i> (M+H) 370.9 found, 371.2 required
17-16		3-[(2S)-1-[(2S)-2-amino-2-cyclopropylethanoyl]-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol	LRMS <i>m/z</i> (M+H) 387.1 found, 387.1 required
17-17		(2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-norleucyl-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 403.3 found, 403.2 required

17-18		(2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(3-methyl-L-valyl)-2,5-dihydro-1H-pyrrole	LRMS <i>m/z</i> (M+H) 413.4 found, 413.2 required
17-19		N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)acetamide	LRMS <i>m/z</i> (M+H) 403.0 found, 403.2 required
17-20		3-[(2S)-1-[(2S)-2-cyclopropyl-2-hydroxyethanoyl]-4-(2,5-difluorophenyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol	LRMS <i>m/z</i> (M+H) 372.3 found, 372.4 required
17-21		3-[(2S)-4-(2,5-difluorophenyl)-1-[(2S)-2-hydroxy-3,3-dimethylbutanoyl]-2,5-dihydro-1H-pyrrol-2-yl]phenol	LRMS <i>m/z</i> (M+H) 388.3 found, 388.4 required



## SCHEME 18



Step 1: 1-[(2S)-4-(2,5-Difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-methyl-1H-imidazol-3-ium (18-2)

5

To a flame-dried flask equipped with stir bar under nitrogen was charged 5-5 (0.63 g, 1.75 mmol) and anhydrous CH<sub>2</sub>Cl<sub>2</sub> (10 mL). The resulting solution was treated with trifluoroacetic acid (5 mL) and stirred 1.5 hours at 25°C. Upon completion, the reaction was concentrated, taken up in CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and washed with 5% NaHCO<sub>3</sub> (50 mL). The organic layer was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The resulting free-amine was dissolved in anhydrous THF (10 mL) and treated with carbonyl diimidazole (0.31 g, 1.93 mmol). The resulting solution was refluxed for 4 hours until completion. The reaction was concentrated, taken up in EtOAc (50 mL) and washed with H<sub>2</sub>O and brine. The combined organic layers were dried (MgSO<sub>4</sub>) and concentrated. The crude acyl

10

15

imidazole was dissolved in anhydrous CH<sub>3</sub>CN and treated with MeI (2.2 mL, 36 mmol). The resulting solution was stirred at 25°C overnight. Upon completion, the reaction was concentrated to give **18-2** as an orange colored solid: LRMS *m/z* (M+H) 365.9 found, 366.1 required.

5

Step 2:           tert-Butyl 4-([[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]amino)piperidine-1-carboxylate (**18-3**)

Crude **18-2** (115 mg, 0.24 mmol) was suspended in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (3 mL) and treated successively with 4-amino-1-tert-butyloxycarbonyl-piperidine (235 mg, 1.17 mmol) and Et<sub>3</sub>N (0.1 mL, 0.7 mmol). The suspension was stirred for 6 hours and complete dissolution was observed. Upon completion, the reaction was diluted further with CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and washed with 1N HCl to remove all unbound amine and Et<sub>3</sub>N. The organic layer was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure to yield crude **18-3** as a yellow oil: LRMS *m/z* (M-C(CH<sub>3</sub>)<sub>3</sub>+H) 427.9 found, 428.1 required.

15

Step 3:           (2S)-4-(2,5-Difluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide (**18-4**)


Crude **18-3** (100 mg, 0.2 mmol) was dissolved in anhydrous DMF (3 mL) and cooled to 0°C under N<sub>2</sub>. NaH (28 mg, 1.17 mmol) was added neat and the reaction was stirred 30 minutes at 0°C. Following cessation of H<sub>2</sub> evolution, MeI (0.15 mL, 2.4 mmol) was added and the reaction was stirred 12 hours at 25°C. Upon completion, the reaction was diluted with EtOAc (10 mL), and the organic layer was washed successively with 5% NH<sub>4</sub>Cl (1 x 10 mL), H<sub>2</sub>O (3 x 5 mL) and brine (1 x 10 mL). The organic layer was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The crude oil was dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (10 mL). The resulting solution was treated with trifluoroacetic acid (5 mL) and stirred 1 hour at 25°C. Upon completion, the reaction was concentrated and directly subjected to reverse-phase column chromatography to provide **18-4**: <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.33 (m, 4H), 7.25 (m, 2H), 7.17 (m, 1H), 7.08 (m, 1H), 6.41 (s, 1H), 6.00 (t, *J* = 2.7 Hz, 1H), 5.07 (ddd, *J* = 14.0, 5.6, 1.7 Hz, 1H), 4.55 (d, *J* = 14.0 Hz, 1H), 3.82 (m, 1H), 3.40 (m, 2H), 3.03 (dt, *J* = 12.5, 3.9 Hz, 1H), 2.88 (s, 3H), 2.81 (dt, *J* = 13.1, 2.9 Hz, 1H), 1.90 (m, 4H); LRMS *m/z* (M+H) 398.0 found, 398.2 required.

20

25

30

## SCHEME 18a



5


(2S)-4-(2,5-Difluorophenyl)-N-methyl-2-phenyl-N-(piperidin-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide (**18a-6**)

To a flame-dried, N<sub>2</sub> purged flask equipped with stir bar was added

10 **18a-5** (prepared in the same way as shown in Scheme 18 but utilizing N-methyl-N-(1-allyloxycarbonyl-4-piperidinylmethyl)-amine) (22 mg, .044 mmol) in anhydrous THF (1 mL). Pd(PPh<sub>3</sub>)<sub>4</sub> (1 mg, .001 mmol) was added neat followed by NaBH<sub>4</sub> (3 mg, .088 mmol) and the reaction was stirred overnight at 25°C. Upon completion, piperidine (10 μL) was added and the reaction was diluted with EtOAc (10 mL), washed with

15 5% aq. NaHCO<sub>3</sub> (10 mL), brine (10 mL) and dried (MgSO<sub>4</sub>). Following filtration and concentration under reduced pressure, the crude was subjected to reverse phase column chromatography to provide pure **18a-6** as the TFA salt: LRMS *m/z* (M+H) 412.0 found, 412.2 required.

## SCHEME 18b



5






(2S)-4-(5-Chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-[(3R)-pyrrolidin-3-yl]-2,5-dihydro-1H-pyrrole-1-carboxamide (**18b-8**)




To a flame-dried, N<sub>2</sub> purged flask equipped with stir bar was added **18b-7** (prepared in the same way as shown in Scheme 18 but utilizing N-methyl-N-(3-pyrrolyl)-amine) (14 mg, 0.026 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 mL). Cooled to 0°C prior to the addition of BBr<sub>3</sub> (0.10 mL, 0.10 mmol). The reaction stirred 1.5 hours at 0°C. Upon completion, piperidine (10 μL) was added and the reaction was diluted with CH<sub>2</sub>Cl<sub>2</sub> (10 mL), and washed with 5% aq. NaHCO<sub>3</sub>. The organic layer was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. Reverse phase chromatography provided **18b-8** as a white solid (isolated as the TFA salt): <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 11.86 (s, 1H), 8.06 (s, 1H), 7.30 (m, 7H), 7.08 (m, 1H), 6.37 (s, 1H), 5.88 (s, 1H), 4.94 (dd, J=5.4, 13.9 Hz, 1H), 4.48 (d, J=14.1, 1H), 3.79 (m, 1H), 3.49 (m, 1H), 3.36 (s, 2H), 3.00 (s, 3H), 2.23 (m, 1H), 1.77 (m, 1H). LRMS *m/z* (M+H) 400.0 found, 400.1 required

The following compounds were prepared by simple modifications of the above procedures. Unless otherwise indicated, the compounds in the table were isolated as the free base.


5

Cmpd	Structure	Name	LRMS $m/z$ (M+H)
18-5		(2S)-4-(5-chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide; isolated as the TFA salt	LRMS $m/z$ (M+H) 414.0 found, 414.1 required.
18-6		(2S)-4-(5-chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-[(3S)-pyrrolidin-3-yl]-2,5-dihydro-1H-pyrrole-1-carboxamide; isolated as the TFA salt	LRMS $m/z$ (M+H) 400.0 found, 400.1 required
18-7		(2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-pyrrolidin-3-yl-2,5-dihydro-1H-pyrrole-1-carboxamide; isolated as the TFA salt	LRMS $m/z$ (M+H) 384.0 found, 384.2 required.
18-8		(2S)-4-(2,5-difluorophenyl)-N-methyl-N-[(3S)-1-methylpyrrolidin-3-yl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 398.1 found, 398.2 required.

18-9		(2S)-4-(2,5-difluorophenyl)-N-methyl-N-[(3R)-1-methylpyrrolidin-3-yl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 398.1 found, 398.2 required.
18-10		(2S)-4-(2,5-difluorophenyl)-N-methyl-N-[(1-methyl-5-oxopyrrolidin-2-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 426.1 found, 426.2 required.
18-11		(2S)-4-(2,5-difluorophenyl)-N-(1,3-dioxolan-2-ylmethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 401.0 found, 401.2 required.
18-12		(2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-tetrahydrofuran-3-yl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 385.0 found, 385.2 required.
18-13		(2S)-N-(1-allylpiperidin-4-yl)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide; isolated as the TFA salt	LRMS <i>m/z</i> (M+H) 438.1 found, 438.2 required.

18-14		allyl 4-{{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}(methyl)amino]piperidine-1-carboxylate	LRMS $m/z$ (M+H) 482.0 found, 482.2 required.
18-15		allyl 4-{{[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}(methyl)amino]methyl}piperidine-1-carboxylate	LRMS $m/z$ (M+H) 496.1 found, 496.2 required.
18-16		(2S)-4-(2,5-difluorophenyl)-N-methyl-N-(1-methylpiperidin-4-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 412.4 found, 412.5 required.

## SCHEME 19



**Step 1:** 4-Nitrophenyl 4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (**19-1**)

5 To a flame-dried flask equipped with stir bar under nitrogen was charged 2-3 (0.71 g, 1.98 mmol) and anhydrous CH<sub>2</sub>Cl<sub>2</sub> (50 mL). The resulting solution was treated with trifluoroacetic acid (10 mL) and stirred 1 hour at 25°C. Upon completion, the reaction was concentrated, taken up in CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and washed with 5% NaHCO<sub>3</sub> (50 mL). The organic layer was dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure. The resulting free-amine was dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and treated with p-nitrophenyl chloroformate (0.44 g, 2.17 mmol). The resulting solution was stirred 4 hours at 25°C until completion. The

10





reaction was concentrated to give **19-1** as a yellow oil:  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.22 (d,  $J = 9.4$  Hz, 1H), (d,  $J = 9.1$  Hz, 1H), 7.38 (m, 5H), 7.02 (m, 5H), 6.43 (m, 1H), 5.85 (m, 1H), 4.90 (m, 2H); LRMS  $m/z$  (M+H) 422.9 found, 423.1 required.






5 Step 2: 4-(2,5-Difluorophenyl)-N-methyl-N-[(1-methylpiperidin-3-yl)methyl]-  
2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (**19-2**)






**19-1** (20 mg, 0.05 mmol) was dissolved in anhydrous DMF (1.0 mL) in a microwave reaction tube. The resulting solution was treated with 4-aminomethyl-N,N'-dimethylpiperidine (13 mg, 0.09 mmol). The solution was stirred for 6 minutes at 250°C  
10 under microwave irradiation. The dark brown reaction was directly purified by reverse phase column chromatography to yield 4-(2,5-difluorophenyl)-N-methyl-N-[(1-methylpiperidin-3-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide **19-2** (mix of four diastereomers) (isolated as the TFA salt): LRMS  $m/z$  (M+H) 426.2 found, 426.2 required.


15

The following compounds were prepared by simple modifications of the above procedures. Unless otherwise indicated, the compounds in the table were isolated as the free base.


Cmpd	Structure	Name	LRMS $m/z$ (M+H)
19-3		4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(pyridin-3-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide; isolated as the TFA salt	LRMS $m/z$ (M+H) 406.1 found, 406.2 required.
19-4		N-benzyl-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 405.2 found, 405.2 required.

19-5		4-(2,5-difluorophenyl)-N-methyl-N-[(1-methyl-1H-pyrazol-4-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide; isolated as the TFA salt	LRMS $m/z$ (M+H) 409.2 found, 409.2 required.
19-6		4-(2,5-difluorophenyl)-N-[2-(dimethylamino)ethyl]-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide; isolated as the TFA salt	LRMS $m/z$ (M+H) 386.2 found, 386.2 required.
19-7		4-(2,5-difluorophenyl)-N-(2-hydroxyethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 359.1 found, 359.0 required.
19-8		4-(2,5-difluorophenyl)-N-isobutyl-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 371.2 found, 371.2 required.
19-9		4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(2-pyridin-2-ylethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide; isolated as the TFA salt	LRMS $m/z$ (M+H) 420.2 found, 420.2 required.


19-10		4-(2,5-difluorophenyl)- N-(2-methoxyethyl)-N- methyl-2-phenyl-2,5- dihydro-1H-pyrrole-1- carboxamide	LRMS $m/z$ (M+H) 373.1 found, 373.2 required.
19-11		4-(2,5-difluorophenyl)- N-(2,3-dihydroxypropyl)- N-methyl-2-phenyl-2,5- dihydro-1H-pyrrole-1- carboxamide	LRMS $m/z$ (M+H) 389.1 found, 389.2 required.
19-12		4-(2,5-difluorophenyl)- N-methyl-2-phenyl-N-(2- phenylethyl)-2,5-dihydro- 1H-pyrrole-1- carboxamide	LRMS $m/z$ (M+H) 419.1 found, 419.2 required.
19-13		4-(2,5-difluorophenyl)-2- phenyl-N-propyl-2,5- dihydro-1H-pyrrole-1- carboxamide	LRMS $m/z$ (M+H) 343.1 found, 343.1 required.
19-14		4-(5-chloro-2- fluorophenyl)-2-(3- hydroxyphenyl)-N- methyl-N-piperidin-4-yl- 2,5-dihydro-1H-pyrrole- 1-carboxamide; isolated as the TFA salt	LRMS $m/z$ (M+H) 430.0 found, 429.2 required.

19-15		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide; isolated as the TFA salt	LRMS $m/z$ (M+H) 414.0 found, 413.2 required.
-------	---	---	---

## SCHEME 20



## SCHEME 20 (continued)



5 Step 1: 2,2,2-Trichloro-N-(1-methyl-1-phenylprop-2-enyl)-N-(2-methylprop-2-enyl)acetamide (**20-2**)

To a flame-dried flask equipped with stir bar under nitrogen was charged Z-3-phenyl-buten-1-ol (3.1 g, 21.0 mmol) and anhydrous  $\text{Et}_2\text{O}$  (100 mL). The resulting solution was treated with  $\text{NaH}$  (0.05 g, 2.1 mmol) and stirred 30 minutes at  
 10 0°C. Trichloroactonitrile (4.12 mL, 41.1 mmol) was added to the cooled solution and stirred an additional 2 hours. The reaction mixture was concentrated under reduced pressure then pentane (100 mL), containing 0.1 equivalent of methanol was added to

the residue and stirred vigorously for 1 hour. The resulting solution was concentrated and washed with pentane (3 x 20 mL). The filtrate was concentrated under reduced pressure to give 2-(E)-3-phenylbut-2-enyl 2,2,2-trichloroethanimidoate. Xylene (80 mL) was added to the crude residue and the solution was heated 3 hours at 150°C.

- 5 The warm solution was concentrated under reduced pressure and purified by flash column chromatography (SiO<sub>2</sub>, 0-20% EtOAc/Hexanes gradient) to give 2,2,2-trichloro-N-(1-methyl-1-phenylprop-2-enyl)-N-(2-methylprop-2-enyl)acetamide, **20-2**.  
<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.39 (m, 5H), 6.98 (s, 1H), 6.29 (dd, J=17.4, 10.6 Hz, 1H), 5.30 (d, J=10.6 Hz, 1H), 5.26 (d, J=17.4 Hz, 1H) 1.89 (s, 3H); LRMS *m/z*  
 10 (M+H) 294.2 found, 294.6 required.

Step 2: tert-Butyl allyl(1-methyl-1-phenylprop-2-enyl)carbamate (20-3)

- Trichloroacetamide (**20-2**) (0.98 g, 3.9 mmol) was stirred 72 h. in a 1:1 solution of 6M NaOH/EtOH (100 mL). The solution was concentrated under  
 15 reduced pressure and purified by flash column chromatography (SiO<sub>2</sub>, 0-20% EtOAc/Hexanes gradient). The purified amine was combined with BOC<sub>2</sub>O (3.81 g, 17.4 mmol) and Et<sub>3</sub>N (1.76 g, 17.4 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (100 mL). The solution was then heated at 35°C for 24 hours. Partitioned between 5% aq. NaHCO<sub>3</sub> (150 mL) and EtOAc (3 x 75 mL). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered,  
 20 and concentrated under reduced pressure. The residue was purified by flash column chromatography (SiO<sub>2</sub>, 0-20% EtOAc/Hexanes gradient) to give tert-butyl-1-methyl-1-phenylprop-2-enylcarbamate. The BOC-protected amine was dissolved in anhydrous DMF (100 mL) under N<sub>2</sub> at 0°C. NaH (0.19 g, 7.9 mmol) was added and the solution was stirred at 0°C for 30 minutes. Allyl bromide (1.37 mL, 15.8 mmol)  
 25 was added neat, and the reaction stirred 24 hours. at 25°C. The reaction mixture was partitioned between 5 % aq. NaHCO<sub>3</sub> (150 mL) and Et<sub>2</sub>O (3 x 75 mL). The combined organic layers were then dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated under reduced pressure. The residue was purified by flash column chromatography (SiO<sub>2</sub>, 0-20% EtOAc/Hexanes gradient) to give tert-butyl allyl(1-methyl-1-phenylprop-2-  
 30 enyl)carbamate, **20-3**. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.20 (m, 5H), 6.34 (dd, J=10.6, 17.4 Hz, 1H), 6.00 (ddt, J=5.8, 11.3, 21.7 Hz, 1H), 5.18 (m, 2H), 5.09 (d, J=20.7 Hz, 1H), 5.01 (d, J=16.8 Hz, 1H), 4.07 (d, J=5.5 Hz, 2H), 1.76 (s, 3H), 1.19 (s, 9H); LRMS *m/z* (M+H-C(CH<sub>3</sub>)<sub>3</sub>) 232.1 found, 232.2 required.

Step 3:            tert-Butyl 2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (20-4)

A solution of **20-3** (0.78 g, 2.7 mmol) and tricyclohexylphosphine[1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene] [benzylidene]ruthenium (IV)dichloride (0.04 g, 0.05 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (500 mL) was stirred 1 hour. at 40°C under N<sub>2</sub>. The reaction mixture was concentrated under reduced pressure and purified by flash column chromatography (SiO<sub>2</sub>, 0-30% EtOAc/Hexanes gradient) to give tert-butyl 2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate, **20-4**. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.25 (m, 5H), 5.79 (m, 1H), 5.58 (m, 1H), 4.32 (m, 1H), 4.43 (m, 1H), 1.8 (s, 3H), 1.5 (s, 9 H); LRMS *m/z* (M+H) 260.1 found, 260.2 required.

Step 4:            tert-Butyl 4-hydroxy-2-methyl-2-phenylpyrrolidine-1-carboxylate (20-5)

A solution of **20-4** (0.50 g, 1.9 mmol) in anhydrous THF (80 mL) at 0°C was treated with BH<sub>3</sub>:THF (1.93 mL, 1.9 mmol) then stirred for 1 hour. at 25°C. Water (0.35 mL, 19.2 mmol) was added, followed by 3 N NaOH (0.23 mL, 5.8 mmol). The resulting solution was cooled to 0°C prior to the addition of 30% H<sub>2</sub>O<sub>2</sub> (0.22 mL, 1.9 mmol). The reaction was partitioned between sat. aq. NaHCO<sub>3</sub> (100 mL) and EtOAc (2 x 100 mL). The combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated under reduced pressure. The residue was purified by flash column (SiO<sub>2</sub>, 0-100% EtOAc/Hexanes gradient) to yield tert-butyl 4-hydroxy-2-methyl-2-phenylpyrrolidine-1-carboxylate, **20-5**: LRMS *m/z* (M+H) 278.2 found, 278.1 required.

Step 5:            tert-Butyl 2-methyl-4-oxo-2-phenylpyrrolidine-1-carboxylate (20-6)

To a flame-dried flask equipped with stir bar under nitrogen was charged anhydrous CH<sub>2</sub>Cl<sub>2</sub> (5 mL). The flask was immersed in a cooling bath at -78°C. Oxalyl chloride (0.24 mL, 2.7 mmol) was added neat followed by DMSO (0.29 mL, 4.1 mmol) dropwise. The reaction was stirred 10 minutes at -78°C prior to the addition of alcohol **20-5** (0.19 g, 0.7 mmol) in minimal CH<sub>2</sub>Cl<sub>2</sub>. The resulting solution was stirred an additional 1 hour at -78°C prior to the addition of Et<sub>3</sub>N (0.76 mL, 5.4 mmol) neat and dropwise. The reaction warmed to 0°C over 2 hour. Upon completion, the reaction solution was washed with 5% aq. NaHCO<sub>3</sub>, dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was carried crude into the vinyl triflate formation: LRMS *m/z* (M-C(CH<sub>3</sub>)<sub>3</sub> + H) 220.1 found, 220.2 required.

Step 6:           tert-Butyl 2-methyl-2-phenyl-4-[[trifluoromethyl)sulfonyl]oxy]-2,5-dihydro-1H-pyrrole-1-carboxylate (**20-7**)

To a solution of ketone **20-6** (70 mg, 25  $\mu$ mol) in anhydrous THF (2 mL) at  $-78^{\circ}\text{C}$  was added LiHMDS (254 mL, 25  $\mu$ mol), and the resulting yellow-tinted solution was stirred for 1 hours. Comins reagent (110 mg, 28  $\mu$ mol) was added neat and the reaction was allowed to warm to  $0^{\circ}\text{C}$  over 1 hour. The reaction mixture was poured into a separatory funnel containing 5%  $\text{NH}_4\text{Cl}$ , and was then extracted with  $\text{CH}_2\text{Cl}_2$  (2 x 10 mL). The combined organic layers were dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to provide the crude triflate **20-7**, which was carried on directly to the Suzuki coupling: LRMS  $m/z$  ( $\text{M}-\text{CH}_3 + \text{H}$ ) 393.0 found, 393.2 required.

Step 7:           tert-Butyl 4-(2,5-difluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (**20-9**)

Crude intermediate **20-7** (100 mg, 25  $\mu$ mol) in degassed 4:1 DME/ $\text{H}_2\text{O}$  (2 mL) was treated with 2,5-difluorophenyl boronic acid (**20-8**, 100 mg, 61  $\mu$ mol), sodium carbonate (78 mg, 74  $\mu$ mol), LiCl (31 mg, 74  $\mu$ mol) and  $\text{Pd}(\text{PPh}_3)_4$  (13 mg, 12  $\mu$ mol). The reaction was kept under  $\text{N}_2$  and heated to  $90^{\circ}\text{C}$  for 1 hour. Upon completion, the reaction was diluted with  $\text{CH}_2\text{Cl}_2$  (10 mL), washed with sat. aq.  $\text{NaHCO}_3$  (10 mL), and brine (10 mL). The organic layer was dried ( $\text{MgSO}_4$ ), filtered and concentrated under reduced pressure. The residue was purified by flash column chromatography ( $\text{SiO}_2$ , 0-30% EtOAc/hexanes gradient) to provide pure **20-9** as a clear oil: LRMS  $m/z$  ( $\text{M}+\text{H}$ ) 408.0 found, 408.2 required.

Step 9:           1-Acetyl-4-(2,5-difluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole (**20-10**)

To a flame-dried flask equipped with stir bar under nitrogen was charged **20-9** (14 mg, 38  $\mu$ mol) and anhydrous  $\text{CH}_2\text{Cl}_2$  (1 mL). The resulting solution was treated with trifluoroacetic acid (0.2 mL) and stirred 1 hour at  $25^{\circ}\text{C}$ . Upon completion, the reaction was concentrated, diluted with toluene (5 mL) and again concentrated. The TFA salt was then suspended in  $\text{CH}_2\text{Cl}_2$  at  $25^{\circ}\text{C}$ , and treated successively with  $\text{Et}_3\text{N}$  (50  $\mu\text{L}$ , 380  $\mu$ mol) and acetic anhydride (25  $\mu\text{L}$ , 20  $\mu$ mol). After 1 hour, the reaction was complete. The reaction was concentrated and flash column chromatography ( $\text{SiO}_2$ , 0-50% EtOAc/hex gradient) provided **20-10** as a clear




oil:  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.36 (m, 5H), 7.03 (m, 3H), 6.22 (m, 1H), 4.82 (m, 2H), 2.15 and 2.05 (s, 3H, rotamers), 1.98 and 1.74 (s, 3H, rotamers); LRMS  $m/z$  (M+H) 314.0 found, 314.2 required.

- 5 The following compound was prepared by simple modifications of the above procedures and isolated as the TFA salt.

Cmp	Structure	Name	LRMS $m/z$ (M+H)
20-11		(2S)-1-[4-(2,5-difluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methyl-1-oxobutan-2-amine	LRMS $m/z$ (M+H) 371.1 found, 371.2 required.

#### SCHEME 21

10



Step 1:        tert-butyl (2S,4S)-4-[[tert-butyl(dimethyl)silyl]oxy]-2-methyl-2-phenylpyrrolidine-1-carboxylate (21-1)

A solution of *sec*-BuLi in cyclohexane (1.4 M, 2.65 mL, 3.71 mmol, 1.40 equiv) was added to a solution of tetramethylethylenediamine (0.880 mL, 5.83 mmol, 2.20 equiv) and tert-butyl (2S,4S)-4-[[tert-butyl(dimethyl)silyl]oxy]-2-phenylpyrrolidine-1-carboxylate (5-1, 1.00 g, 2.65 mmol, 1 equiv) in THF (150 mL) at -78°C. The resulting bright yellow mixture was stirred for 1 hour, then methyl trifluoromethanesulfonate (1.20 mL, 10.6 mmol, 4.00 equiv) was added. The mixture was warmed to 0°C and stirred for 30 minutes, then partitioned between saturated sodium chloride solution (100 mL) and ethyl acetate (2 x 75 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes, grading to 50% ethyl acetate in hexanes) to provide tert-butyl (2S,4S)-4-[[tert-butyl(dimethyl)silyl]oxy]-2-methyl-2-phenylpyrrolidine-1-carboxylate (21-1) as a colorless oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.38-7.18 (m, 5H), 4.39 (m, 1H), 3.92 (dd, 1H, *J* = 10.8, 5.1 Hz), 3.65 (br d, 1H, *J* = 10.8 Hz), 2.28 (dd, 1H, *J* = 13.0, 4.5 Hz), 2.13 (br d, 1H, *J* = 12.8 Hz), 1.92 (s, 3H), 1.12 (s, 9H), 0.91 (s, 9H), 0.03 (s, 3H), 0.02 (s, 3H). LRMS *m/z* (M+H-Boc) 292.2 found, 292.2 required.


Step 2:        tert-butyl (2S,4S)-4-hydroxy-2-methyl-2-phenylpyrrolidine-1-carboxylate (21-2)



A solution of (2S,4S)-4-[[tert-butyl(dimethyl)silyl]oxy]-2-methyl-2-phenylpyrrolidine-1-carboxylate (21-1, 250 mg, 0.638 mmol, 1 equiv) and triethylamine trihydrofluoride (0.310 mL, 1.91 mmol, 3.00 equiv) in acetonitrile (10 mL) was stirred at 23°C for 16 hours. The reaction mixture was partitioned between aqueous saturated sodium bicarbonate solution (100 mL) and ethyl acetate (2 x 50 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes initially, grading to 50% ethyl acetate in hexanes) to provide tert-butyl (2S,4S)-4-hydroxy-2-methyl-2-phenylpyrrolidine-1-carboxylate (21-2) as a colorless oil. LRMS *m/z* (M+H-Boc) 278.2 found, 278.2 required.

Step 3: (2S)-4-(2,5-difluorophenyl)-N,N,2-trimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (21-4)


A solution of tert-butyl (2S)-4-(2,5-difluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (21-3, prepared from tert-butyl (2S,4S)-4-hydroxy-2-methyl-2-phenylpyrrolidine-1-carboxylate (21-2) by the procedures of Scheme 20, 24 mg, 0.065 mmol, 1 equiv) in a 1:1 mixture of trifluoroacetic acid and dichloromethane (10 mL) was stirred for 30 minutes, then concentrated. The residue was dried via toluene azeotrope (2 x 10 mL), then dissolved in dichloromethane (10 mL). Triethylamine (0.090 mL, 0.65 mmol, 10 equiv) and dimethylcarbamoyl chloride (0.030 mL, 0.33 mmol, 5.0 equiv) were added, and the resulting mixture was stirred at 40°C for 6 hours, then 23°C for 2 days. The reaction mixture was partitioned between aqueous saturated sodium bicarbonate solution (30 mL) and ethyl acetate (2 x 25 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography (hexanes initially, grading to 40% ethyl acetate in hexanes) to provide (2S)-4-(2,5-difluorophenyl)-N,N,2-trimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (21-4) as an off-white solid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 7.40 (br d, 2H, *J* = 7.7 Hz), 7.32 (br t, 2H, *J* = 7.8 Hz), 7.22 (t, 1H, *J* = 7.5 Hz), 7.08-6.91 (m, 3H), 6.24 (m, 1H), 4.82 (dd, 1H, *J* = 13.9, 1.5 Hz), 4.77 (dd, 1H, *J* = 13.9, 1.5 Hz), 2.77 (s, 6H), 2.02 (s, 3H). LRMS *m/z* (M+H) 343.0 found, 343.2 required.

The following compounds were prepared by simple modifications of the above procedures.

Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
21-5		(2S)-4-(5-chloro-2-fluorophenyl)-N,N,2-trimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 359.0 found, 359.1 required.

21-6		(2S)-4-(5-chloro-2-fluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 331.0 found, 331.1 required
21-7		(2S)-4-(5-chloro-2-fluorophenyl)-N-ethyl-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 359.0 found, 359.1 required



SCHEME 22









5 (2S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3,3-dimethyl-1-oxobutan-2-ol (22-1)

A solution of tert-butyl (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (5-5, 113 mg, 0.316 mmol, 1 equiv) in a 1:1 mixture of dichloromethane and trifluoroacetic acid (20 mL) was stirred for 30 minutes, then concentrated. The residue was dried via toluene azeotrope (1 x 10 mL), then dissolved in DMF (4 mL). Triethylamine (0.176 mL, 1.27 mmol, 4.00 equiv), (2S)-2-hydroxy-3,3-dimethylbutanoic acid (84 mg, 0.632 mmol, 2.00 equiv), and


- PYBOP (329 mg, 0.632 mmol, 2.00 equiv) were added, and the resulting mixture was stirred at 23°C for 24 hours. Additional triethylamine (0.176 mL, 1.27 mmol, 4.00 equiv), (2S)-2-hydroxy-3,3-dimethylbutanoic acid (84 mg, 0.632 mmol, 2.00 equiv), and PYBOP (329 mg, 0.632 mmol, 2.00 equiv) were added, and the resulting mixture
- 5 was stirred at 23°C for 20 hours. The reaction mixture was partitioned between water (70 mL) and ethyl acetate (2 x 55 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide (2S)-1-[(2S)-4-(2,5-
- 10 difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3,3-dimethyl-1-oxobutan-2-ol (22-1) as a white solid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) rotamers: δ 7.43-7.02 (m, 8H), 6.42 (m, 1H), 5.93 (m, 1H), 5.01-4.82 (m, 2H), 4.16 (major rotamer, s, 1H), 3.85 (minor rotamer, s, 1H), 1.00 (minor, s, 9H), 0.92 (major, s, 9H). LRMS *m/z* (M+H) 372.0 found, 372.2 required.
- 15 The following compounds were prepared by simple modifications of the above procedures.

Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
22-2		(2S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methyl-1-oxobutan-2-ol	LRMS <i>m/z</i> (M+H) 358.0 found, 358.2 required.
22-3		(2S,3S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methyl-1-oxopentan-2-ol	LRMS <i>m/z</i> (M+H) 372.0 found, 372.2 required

22-4		1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-oxohexan-2-ol	LRMS <i>m/z</i> (M+H) 372.0 found, 372.2 required
22-5		(2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-oxo-3-phenylpropan-2-ol	LRMS <i>m/z</i> (M+H) 406.1 found, 406.2 required
22-6		(2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-methyl-1-oxopentan-2-ol	LRMS <i>m/z</i> (M+H) 372.0 found, 372.2 required
22-7		(1S)-1-cyclohexyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanol	LRMS <i>m/z</i> (M+H) 398.1 found, 398.2 required

22-8		(2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3,3-dimethyl-1-oxobutan-2-ol	LRMS <i>m/z</i> (M+H) 372.0 found, 372.2 required
22-9		(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanol	LRMS <i>m/z</i> (M+H) 356.3 found, 356.1 required

## SCHEME 23



5

Step 1: tert-butyl (2S,4S)-4-hydroxy-2-phenylpyrrolidine-1-carboxylate (5-2)



To a flame dried flask equipped with stir bar was added tert-butyl (2S,4S)-4-{{tert-butyl(dimethyl)silyl}oxy}-2-phenylpyrrolidine-1-carboxylate (5-1, prepared from (S)-(-)-4-chloro-3-hydroxybutyronitrile by the method of Maeda, *et al Synlett* 2001, 1808-1810, 7.8 g, 20.7 mmol) and anhydrous acetonitrile (20.0 mL).

- 5 The resulting solution was treated with triethylamine trihydrofluoride (10.1 mL, 62.0 mmol) while stirring under N<sub>2</sub>. The reaction stirred 12 hours at 40°C. The reaction was then diluted with EtOAc (100 mL) and poured into 5% aq. NaHCO<sub>3</sub>. Following cessation of gas evolution, the organic layer was washed three addition times with 5% aq. NaHCO<sub>3</sub>. The organic layer was dried over magnesium sulfate, filtered and
- 10 concentrated to provide crude product. Recrystallization was effected from EtOAc/hexanes to provide tert-butyl (2S,4S)-4-hydroxy-2-phenylpyrrolidine-1-carboxylate (5-2) as a white crystalline solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) rotamers δ 7.38-7.18 (m, 5H), 4.90 (m, 1H), 4.42 (m, 1H), 3.88 (m, 1H), 3.56 (dd, *J* = 11.5, 4.0 Hz, 1H), 2.60 (m, 1H), 2.03 (m, 1H), 1.50 and 1.20 (br s, 9H); MS 208.0 found, 208.1
- 15 (M – C(CH<sub>3</sub>)<sub>3</sub>) required.

Step 2: tert-butyl (2S)-4-oxo-2-phenylpyrrolidine-1-carboxylate (5-3)

- To a flame dried flask equipped with stir bar was added 150 mL anhydrous dichloromethane which was cooled to –78°C. Oxalyl chloride (3.8 mL, 44
- 20 mmol) and DMSO (4.8 mL, 61 mmol) were added sequentially and the reaction stirred for 10 min. tert-butyl (2S,4S)-4-hydroxy-2-phenylpyrrolidine-1-carboxylate (5-2, 2.28 g, 8.73 mmol) in 10 mL anhydrous dichloromethane was added dropwise and stirred 1 hour at –78°C. Triethylamine (12 mL, 87mmol) was added and the reaction was warmed to 0°C over 1 hour. Upon completion, the reaction was washed with 5%
- 25 NaHCO<sub>3</sub>, brine and dried over MgSO<sub>4</sub>. The organic layer was concentrated to provide crude tert-butyl (2S)-4-oxo-2-phenylpyrrolidine-1-carboxylate (5-3). Recrystallization was effected with EtOAc/hexanes. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.35 (m, 3H), 7.17 (m, 2H), 5.38 (m, 1H), 4.08 (d, *J* = 19.5 Hz, 1H), 3.90 (d, *J* = 19.3 Hz, 1H), 3.13 (dd, *J* = 18.8, 9.8 Hz, 1H), 2.58 (dd, *J* = 18.6, 2.4 Hz, 1H), 1.40 (br s,
- 30 9H); MS 206.0 found, 206.1 (M – C(CH<sub>3</sub>)<sub>3</sub>) required.

Step 3: tert-butyl (2S)-2-phenyl-4-[[trifluoromethyl)sulfonyl]oxy}-2,5-dihydro-1H-pyrrole-1-carboxylate (5-4)

- To a flame dried flask equipped with stir bar was added ketone tert-
- 35 butyl (2S)-4-oxo-2-phenylpyrrolidine-1-carboxylate (5-3, 2.00 g, 7.65 mmol) and

anhydrous THF (100 mL). The resulting solution was cooled to  $-78^{\circ}\text{C}$ , and treated dropwise with sodium hexamethyldisilylamide (NaHMDS, 8.42 mL, 1M in THF, 8.42 mmol). The reaction was stirred 1 hour at  $-78^{\circ}\text{C}$ , and a solution of 1,1,1-trifluoro-N-phenyl-N-[(trifluoromethyl)sulfonyl]methanesulfonamide (3.01 g, 8.42 mmol) in THF  
 5 (30 mL) was added via cannula. The reaction mixture was warmed to  $0^{\circ}\text{C}$  and stirred 30 minutes. The reaction mixture was then partitioned between brine (200 mL) and a 1:1 mixture of ethyl acetate and hexane (200 mL). The organic layer was dried over sodium sulfate and concentrated to provide crude tert-butyl (2S)-2-phenyl-4-  
 10 {[(trifluoromethyl)sulfonyl]oxy}-2,5-dihydro-1H-pyrrole-1-carboxylate (5-4) as an orange oil.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ) major rotamer:  $\delta$  7.30 (m, 5H), 5.72 (m, 1H), 5.48 (m, 1H), 4.42 (m, 2H), 1.18 (s, 9H); MS 379.0 found 379.1 (M –  $\text{CH}_3$ ) required.

Step 4:           tert-butyl (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (5-5)

15           A deoxygenated mixture of crude tert-butyl (2S)-2-phenyl-4-  
 {[(trifluoromethyl)sulfonyl]oxy}-2,5-dihydro-1H-pyrrole-1-carboxylate (5-4, 7.65 mmol), 2,5-difluorophenyl boronic acid (1.81 g, 11.5 mmol), aqueous  $\text{Na}_2\text{CO}_3$  solution (2 M, 11.5 mL, 23.0 mmol), and  $\text{Pd}(\text{PPh}_3)_4$  (0.442 g, 0.383 mmol) in dioxane (100 mL) was heated at  $90^{\circ}\text{C}$  for 45 minutes. The reaction mixture was cooled, then  
 20 partitioned between brine (200 mL) and a 1:1 mixture of ethyl acetate and hexane (200 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography ( $\text{SiO}_2$ , 0-50% EtOAc/hexanes gradient) to provide tert-butyl (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (5-5) as a white solid. LRMS  $m/z$  (M+H- $\text{CH}_3$ ) 358.0 found,  
 25 358.2 required.


Step 5:           (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine (16-7)

30           A solution of tert-butyl (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (5-5, 1.00 g, 2.80 mmol, 1 equiv) in a 4:1 mixture of dichloromethane and trifluoroacetic acid (20 mL) was stirred for 30 minutes, then concentrated. The residue was dried via benzene azeotrope (1 x 50 mL), then dissolved in DMF (30 mL). (2S)-[(tert-butoxycarbonyl)amino](cyclopropyl)ethanoic acid (0.903 g, 4.20 mmol, 1.50 equiv), 1-(3-dimethylaminopropyl)-3-

ethylcarbodiimide hydrochloride (EDC, 0.805 g, 4.20 mmol, 1.50 equiv), 1-hydroxy-7-azabenzotriazole (0.571, 4.20 mmol, 1.50 equiv) and triethylamine (1.95 mL, 14.0 mmol, 5.00 equiv) were added, and the resulting mixture was stirred at 23°C for 5 days. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution and ethyl acetate (2 x 100 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was dissolved in a 4:1 mixture of dichloromethane and trifluoroacetic acid. The resulting solution was allowed to stand for 25 minutes, then concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present). The desired fractions were partitioned between saturated aqueous sodium bicarbonate solution and ethyl acetate (2 x 50 mL), and the combined organic layers were washed with aqueous sodium bicarbonate solution (2 x 100 mL), then brine (100 mL), and dried over sodium sulfate and concentrated to provide (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine (16-7) as a white solid. <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) major rotamer: δ 7.41 (t, 2H, *J* = 7.6 Hz), 7.33 (m, 3H), 7.26 (m, 1H), 7.18 (m, 1H), 7.09 (m, 1H), 6.46 (s, 1H), 5.91 (m, 1H), 5.02 (d, 2H, *J* = 13.9 Hz), 4.96 (ddd, 1H, *J* = 13.9, 4.4, 1.7 Hz), 3.08 (d, 1H, *J* = 7.6 Hz), 1.04 (m, 1H), 0.52 (m, 3H), 0.37 (m, 1H). LRMS *m/z* (M+H) 355.0 found, 355.2 required.

20




## SCHEME 24








25 N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-N,N-dimethylglycinamide (24-1)

A solution of (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine (16-7, 30 mg, 0.085 mmol, 1 equiv), (N,N-dimethylglycine (17 mg, 0.17 mmol, 2.0 equiv), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC, 32 mg, 0.17 mmol, 2.0 equiv), 1-hydroxy-7-azabenzotriazole (23 mg, 0.17 mmol, 2.0 equiv) and triethylamine (35  $\mu$ L, 0.25 mmol, 3.0 equiv) in DMF (1 mL) was stirred at 23°C for 20 hours. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution and ethyl acetate (2 x 20 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-N-2,N-2-dimethylglycinamide (24-1) as a colorless gum (TFA salt). <sup>1</sup>H NMR (500 MHz, CD<sub>3</sub>OD) major rotamer:  $\delta$  7.33 (m, 4H), 7.26 (m, 2H), 7.26 (m, 1H), 7.19 (m, 1H), 7.10 (m, 1H), 6.46 (s, 1H), 5.91 (m, 1H), 5.23 (d, 2H, *J* = 13.9 Hz), 5.02 (ddd, 1H, *J* = 13.9, 4.4, 1.7 Hz), 4.19 (d, 1H, *J* = 9.0 Hz), 4.01 (d, 1H, *J* = 15.6 Hz), 3.94 (d, 1H, *J* = 15.4 Hz), 2.92 (s, 6H), 1.14 (m, 1H), 0.62 (m, 3H), 0.40 (m, 1H). LRMS *m/z* (M+H) 440.5 found, 440.2 required.


The following compounds were prepared by simple modifications of the above procedure. For many of the examples below, Boc-protected amino acids were utilized. In these cases the coupling products required a deprotection step (TFA/dichloromethane) to afford the depicted final products. Unless otherwise noted, the compounds were isolated as the free base. Additionally, the amine n-oxides below were prepared by treating the corresponding amines with hydrogen peroxide in isopropanol at 23°C.

Cmpd	Structure	Name	LRMS $m/z$ (M+H)
24-2		N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2-methylglycinamide; isolated as the TFA salt	$m/z$ (M+H) 426.4 found, 426.2 required.
24-3		N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)glycinamide; isolated as the TFA salt	$m/z$ (M+H) 412.4 found, 412.2 required
24-4		N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-methylalaninamide; isolated as the TFA salt	$m/z$ (M+H) 440.4 found, 440.2 required

24-5		N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)glycinamide	<i>m/z</i> (M+H) 428.5 found, 428.2 required
24-6		N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2,N-2-dimethylglycinamide	<i>m/z</i> (M+H) 456.6 found, 456.2 required
24-7		N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2,N-2-dimethylglycinamide, N-oxide	<i>m/z</i> (M+H) 472.5 found, 472.2 required

24-8		N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-methylalaninamide	<i>m/z</i> (M+H) 456.5 found, 456.2 required
24-9		N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2,N-2-dimethylglycinamide n-oxide	LRMS <i>m/z</i> (M+H) 456.5 found, 456.2 required

## SCHEME 25



- 5 **Step 1:** 2-bromo-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}acetamide (25-1)
- Bromoacetyl bromide (49  $\mu\text{L}$ , 0.56 mmol, 1.1 equiv) was added to a solution of (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine (16-7, 181 mg, 0.511 mmol, 1 equiv) and N,N-diisopropylethylamine (107  $\mu\text{L}$ , 0.613 mmol, 1.20 equiv) in dichloromethane (10 mL) at 23°C, and the resulting mixture was allowed to stir for 30 minutes. The reaction mixture was partitioned between water (70 mL) and dichloromethane (2 x 60 mL). The combined organic layers were dried over sodium sulfate and concentrated to give 2-bromo-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}acetamide (25-1) as a pink foam. LRMS  $m/z$  (M+H) 477.3 found, 477.1 required.
- 10
- 15







Step 2: N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-pyrrolidin-1-ylacetamide (25-2)




- A solution of 2-bromo-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}acetamide (25-1, 40 mg, 0.084 mmol, 1 equiv), N,N-diisopropylethylamine (19  $\mu$ L, 0.11 mmol, 1.30 equiv), and pyrrolidine (8  $\mu$ L, 0.093 mmol, 1.1 equiv) in a 1:1 mixture of dioxane and isopropanol (4 mL) was stirred at 23°C for 20 hours. The reaction mixture was partitioned between water (50 mL) and ethyl acetate (50 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-pyrrolidin-1-ylacetamide (25-2) as a colorless oil (TFA salt). LRMS *m/z* (M+H) 466.2 found, 466.2 required.




15





The following compounds were prepared by simple modifications of the above procedure. Unless otherwise noted, the compounds were isolated as the free base.

Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
25-3		2-azetidin-1-yl-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}acetamide; isolated as the TFA salt	<i>m/z</i> (M+H) 452.2 found, 452.2 required


25-4		N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-morpholin-4-ylacetamide; isolated as the TFA salt	<i>m/z</i> (M+H) 482.2 found, 482.2 required
25-5		N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-piperazin-1-ylacetamide; isolated as the TFA salt	<i>m/z</i> (M+H) 481.2 found, 481.2 required
25-6		N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-(4-methylpiperazin-1-yl)acetamide; isolated as the TFA salt	<i>m/z</i> (M+H) 495.2 found, 495.2 required

25-7		2-azetidin-1-yl-N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}acetamide	<i>m/z</i> (M+H) 468.6 found, 468.2 required
25-8		N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-pyrrolidin-1-ylacetamide	<i>m/z</i> (M+H) 482.6 found, 482.3 required
25-9		N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-piperidin-1-ylacetamide	<i>m/z</i> (M+H) 496.6 found, 496.3 required

25-10		N-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-morpholin-4-ylacetamide	<i>m/z</i> (M+H) 498.5 found, 498.3 required
25-11		N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2-(2-hydroxyethyl)glycinamide	<i>m/z</i> (M+H) 472.6 found, 472.2 required
25-12		N-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-(4-methylpiperazin-1-yl)acetamide	<i>m/z</i> (M+H) 511.7 found, 511.3 required

25-13		N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2-isopropylglycinamide; isolated as the TFA salt	<i>m/z</i> (M+H) 454.2 found, 454.2 required
25-14		N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)acetamide	<i>m/z</i> (M+H) 397.3 found, 397.2 required
25-15		N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2-ethylglycinamide	<i>m/z</i> (M+H) 456.6 found, 456.2 required
25-16		N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-hydroxyacetamide	<i>m/z</i> (M+H) 413.4 found, 413.2 required

## SCHEME 26




- Step 1:** 4-nitrophenyl (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate (26-1)
- 4-nitrophenyl chloroformate (171 mg, 0.85 mmol, 1.5 equiv) was added to a solution of (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine (16-7, 200 mg, 0.564 mmol, 1 equiv) and N,N-diisopropylethylamine (172  $\mu$ L, 0.988 mmol, 1.75 equiv) in dichloromethane (5 mL) at 0°C, and the resulting mixture was stirred for 15 minutes, then warmed to 23°C and stirred for an additional 30 minutes. Additional 4-nitrophenyl chloroformate (1.0 equiv) and N,N-diisopropylethylamine (1.0 equiv) were added and the mixture was stirred for 2.5 hours, then partitioned between water (55 mL) and dichloromethane (2 x 45 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide 4-nitrophenyl (1S)-1-cyclopropyl-2-[(2S)-




4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate (26-1) as a light yellow oil. LRMS  $m/z$  (M+H) 520.4 found, 520.2 required.

Step 2: N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}piperazine-1-carboxamide (26-2)


A solution of 4-nitrophenyl (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate (26-1, 40 mg, 0.077 mmol, 1 equiv), N,N-diisopropylethylamine (27  $\mu$ L, 0.15 mmol, 2.0 equiv), and tert-butyl piperazine-1-carboxylate (22 mg, 0.11 mmol, 1.5 equiv) in dioxane (5 mL) was heated at 70°C for 3 hours. The reaction mixture was partitioned between half-saturated aqueous sodium chloride solution (75 mL) and ethyl acetate (55 mL). The organic layer was dried over sodium sulfate and concentrated. A solution of the residue in a 1:1 mixture of TFA and dichloromethane (3 mL) was stirred for 1 hour, then concentrated. The residue was purified by reverse-phase LC ( $H_2O/CH_3CN$  gradient w/ 0.1 % TFA present) to provide N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}piperazine-1-carboxamide (26-2) as a colorless oil (TFA salt). LRMS  $m/z$  (M+H) 467.4 found, 467.2 required.

The following compounds were prepared by simple modifications of the above procedure. All of the compounds in the table below were isolated as the TFA salt.


Cmpd	Structure	Name	LRMS $m/z$ (M+H)
26-3		N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-piperidin-4-ylurea	$m/z$ (M+H) 481.4 found, 481.2 required

26-4		4-amino-N-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}piperidine-1-carboxamide	<i>m/z</i> (M+H) 481.4 found, 481.2 required
26-5		N-(2-aminoethyl)-N'-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}urea	<i>m/z</i> (M+H) 441.4 found, 441.2 required
26-6		N-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-(3-morpholin-4-ylpropyl)urea	<i>m/z</i> (M+H) 525.4 found, 525.3 required



26-7		N-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-N'-[2-(dimethylamino)ethyl]urea	<i>m/z</i> (M+H) 469.4 found, 469.2 required
------	---	---	--

## SCHEME 27






Step 1: 2-chloro-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}ethanesulfonamide (27-1)



2-chloroethanesulfonyl chloride (81  $\mu$ L, 0.78 mmol, 1.1 equiv) was added to a solution of (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine (16-7, 250 mg, 0.705 mmol, 1 equiv) and N,N-diisopropylethylamine (160  $\mu$ L, 0.917 mmol, 1.30 equiv) in dichloromethane (5 mL) at 0°C, and the resulting mixture was stirred for 15 minutes, then warmed to 23°C and stirred for an additional 15 minutes. Additional 2-chloroethanesulfonyl chloride (1.2 equiv) and N,N-diisopropylethylamine (1.3 equiv) were added and the mixture was stirred for 30 minutes, then partitioned between water (75 mL) and dichloromethane (2 x 60 mL). The combined organic layers were dried over sodium sulfate and concentrated to give 2-chloro-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}ethanesulfonamide (27-1) as a colorless oil. LRMS  $m/z$  (M+H-Cl) 445.4 found, 445.1 required.

Step 2: 2-azetidin-1-yl-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}ethanesulfonamide (27-2)


A solution of 2-chloro-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}ethanesulfonamide (27-1, 55 mg, 0.11 mmol, 1 equiv), N,N-diisopropylethylamine (80  $\mu$ L, 0.46 mmol, 4.0 equiv), and azetidine (15  $\mu$ L, 0.23 mmol, 2.0 equiv) in dioxane (2 mL) was heated at 60°C for 20 hours. The reaction mixture was partitioned between water (40 mL) and ethyl acetate (40 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC ( $H_2O/CH_3CN$  gradient w/ 0.1 % TFA present) to provide 2-azetidin-1-yl-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}ethanesulfonamide (27-2) as a light yellow oil (TFA salt). LRMS  $m/z$  (M+H) 502.4 found, 502.2 required.

The following compounds were prepared by simple modifications of the above procedure. All of the compounds in the table below were isolated as the TFA salt.

Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
27-3		N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-(isopropylamino)ethanesulfonamide	<i>m/z</i> (M+H) 504.4 found, 504.2 required.
27-4		N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-pyrrolidin-1-ylethanesulfonamide	LRMS <i>m/z</i> (M+H) 516.4 found, 516.2 required
27-5		N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-morpholin-4-ylethanesulfonamide	<i>m/z</i> (M+H) 532.3 found, 532.2 required

27-6		N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-piperazin-1-ylethanesulfonamide	<i>m/z</i> (M+H) 531.4 found, 531.2 required
27-7		N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-(4-methylpiperazin-1-yl)ethanesulfonamide	<i>m/z</i> (M+H) 545.4 found, 545.2 required

SCHEME 28




Step 1: [(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetic acid (28-1)





- 5 A solution of tert-butyl (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (5-5, 50 mg, 0.14 mmol, 1 equiv) in a 1:1 mixture of dichloromethane and trifluoroacetic acid (10 mL) was stirred for 30 minutes, then concentrated. The residue was partitioned between aqueous saturated sodium bicarbonate solution (60 mL) and ethyl acetate (50 mL). The organic layer was dried
- 10 over sodium sulfate and concentrated. A solution of the residue, ethyl bromoacetate (16  $\mu\text{L}$ , 0.14 mmol, 1.0 equiv) and N,N-diisopropylethylamine (29  $\mu\text{L}$ , 0.17 mmol, 1.2 equiv) in DMF (5 mL) was heated at 70°C for 3 hours. The reaction mixture was partitioned between water (125 mL) and ethyl acetate (2 x 100 mL). The combined organic layers were dried over sodium sulfate and concentrated. A solution of the
- 15 residue and sodium hydroxide (1N, 280  $\mu\text{L}$ , 0.28 mmol, 2.0 equiv) in *tert*-butanol (10 mL) was heated at 45°C for 20 hours. The reaction mixture was then partitioned between water and ethyl ether. The aqueous layer was acidified to pH 4 with aqueous HCl solution, then extracted with ethyl acetate (2 x 40 mL). The combined organic layers were dried over sodium sulfate and concentrated to provide [(2S)-4-(2,5-
- 20 difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetic acid (28-1) as a colorless oil. LRMS  $m/z$  (M+H) 316.3 found, 316.1 required.





Step 2:      N-(tert-butyl)-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetamide (28-2)

A solution of [(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetic acid (28-1, 16 mg, 0.051 mmol, 1 equiv), *tert*-butylamine (6  $\mu$ L, 0.06 mmol, 1.2 equiv), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC, 12 mg, 0.061 mmol, 1.2 equiv), 1-hydroxy-7-azabenzotriazole (8 mg, 0.06 mmol, 1.2 equiv) and triethylamine (18  $\mu$ L, 0.13 mmol, 2.5 equiv) in DMF (2 mL) was stirred at 23°C for 20 hours. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution (50 mL) and ethyl acetate (50 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide N-(tert-butyl)-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetamide (28-2) as a colorless oil (TFA salt). LRMS *m/z* (M+H) 371.4 found, 371.2 required.





The following compounds were prepared by simple modifications of the above procedure. All of the compounds in the table below were isolated as the TFA salt.




Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
28-3		2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylacetamide	<i>m/z</i> (M+H) 357.2 found, 357.2 required

28-4		(2S)-1-(2-azetidin-1-yl-2-oxoethyl)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole	<i>m/z</i> (M+H) 355.1 found, 355.2 required
28-5		(2S)-4-(2,5-difluorophenyl)-1-(2-oxo-2-pyrrolidin-1-ylethyl)-2-phenyl-2,5-dihydro-1H-pyrrole	<i>m/z</i> (M+H) 369.1 found, 369.2 required
28-6		4-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl)morpholine	<i>m/z</i> (M+H) 385.1 found, 385.2 required
28-7		1-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl)piperazine	<i>m/z</i> (M+H) 384.2 found, 384.2 required


28-8		1-[[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl]-4-methylpiperazine	<i>m/z</i> (M+H) 398.2 found, 398.2 required
28-9		2-[[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylbutanamide	<i>m/z</i> (M+H) 385.3 found, 385.2 required
28-10		4-{2-[[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]butanoyl]morpholine	<i>m/z</i> (M+H) 413.4 found, 413.2 required
28-11		2-[[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethylacetamide	<i>m/z</i> (M+H) 343.3 found, 343.2 required



28-12		N-cyclobutyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetamide	<i>m/z</i> (M+H) 369.3 found, 369.2 required
28-13		2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethylpropanamide	<i>m/z</i> (M+H) 357.3 found, 357.2 required
28-14		N-cyclobutyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanamide	<i>m/z</i> (M+H) 383.4 found, 383.2 required
28-15		2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methylpropanamide	<i>m/z</i> (M+H) 343.2 found, 343.2 required

28-16		2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylpropanamide	<i>m/z</i> (M+H) 371.4 found, 371.2 required
28-17		N-(tert-butyl)-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanamide	<i>m/z</i> (M+H) 385.3 found, 385.2 required
28-18		4-{2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoyl}morpholine	<i>m/z</i> (M+H) 399.3 found, 399.2 required

## SCHEME 29

5 Step 1: methyl 2,2-dimethyl-3-oxopropanoate (29-1)

DMSO (22.0 mL, 303 mmol, 4.00 equiv) was added to a pre-cooled solution of oxalyl chloride (13.7 mL, 151 mmol, 2.00 equiv) in dichloromethane (40 mL) at  $-78^\circ\text{C}$ , and the resulting mixture was stirred for 15 minutes before a solution of methyl 3-hydroxy-2,2-dimethylpropanoate (10.0 g, 75.7 mmol, 1 equiv) in dichloromethane (100 mL) was added. The reaction mixture was stirred for 1 hour,

then triethylamine (52.7 mL, 378 mmol, 5.00 equiv) was added. The mixture was warmed to 0°C and held for 30 minutes, then concentrated to a volume of ca 50 mL. The residue was partitioned between aqueous saturated sodium bicarbonate solution (200 mL) and ethyl acetate (200 mL). The organic layer was dried over sodium sulfate and concentrated on a rotatory evaporator without using heat (product is volatile) to provide methyl 2,2-dimethyl-3-oxopropanoate (29-1) as a colorless oil. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 9.66 (s, 1H), 3.76 (s, 3H), 1.36 (s, 6H).

Step 2: methyl 3-amino-3-cyano-2,2-dimethylpropanoate (29-2)

A solution of methyl 2,2-dimethyl-3-oxopropanoate (29-1, 6.50 g, 49.9 mmol, 1 equiv), sodium bisulfite (10.4 g, 99.9 mmol, 2.00 equiv), and concentrated ammonium hydroxide (15 M, 33.3 mL, 499 mmol, 10.0 equiv) in water (200 mL) was stirred at 23°C for 20 minutes, then potassium cyanide (6.51 g, 99.9 mmol, 2.00 equiv) was added. The resulting mixture was stirred at 23°C for 20 hours, then extracted with ethyl ether (3 x 100 mL). The combined organic layers were washed with brine, then dried over sodium sulfate and concentrated to provide methyl 3-amino-3-cyano-2,2-dimethylpropanoate (29-2) as a colorless oil. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 3.93 (br m, 1H), 3.93 (s, 3H), 1.76 (br s, 2H), 1.36 (s, 3H), 1.35 (s, 3H).

Step 3: N-(tert-butoxycarbonyl)-3,3-dimethylaspartic acid (29-3)

A solution of methyl 3-amino-3-cyano-2,2-dimethylpropanoate (29-2, 6.5 g, 41.6 mmol, 1 equiv) in aqueous 3 N HCl solution (100 mL) was heated at reflux for 6 h, then concentrated. Di-*tert*-butyl dicarbonate (13.6 g, 62.4 mmol, 1.50 equiv) was added to a solution of the residue in a mixture of aqueous 1 N NaOH solution (125 mL, 125 mmol, 3.00 equiv) and dioxane (125 mL) at 23°C. The resulting mixture was stirred for 5 hours, then extracted with ethyl ether (2 x 100 mL, discarded). The aqueous layer was acidified to pH 3 with aqueous HCl, then extracted with ethyl ether (3 x 150 mL). The combined organic layers were dried over sodium sulfate and concentrated to provide N-(tert-butoxycarbonyl)-3,3-dimethylaspartic acid (29-3) as a colorless oil. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 5.43 (d, 1H, *J* = 9.8 Hz), 4.57 (d, 1H, *J* = 10.1 Hz), 1.76 (br s, 2H), 1.46 (s, 9H), 1.42 (s, 3H), 1.26 (s, 3H).

Step 4: (3S)-3-[(tert-butoxycarbonyl)amino]-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid (29-4)

A solution of tert-butyl (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (5-5, 1.00 g, 2.80 mmol, 1 equiv) in a 4:1 mixture of dichloromethane and trifluoroacetic acid (20 mL) was stirred for 30 minutes, then concentrated. The residue was dried via benzene azeotrope (1 x 50 mL), then dissolved in DMF (30 mL). N-(tert-butoxycarbonyl)-3,3-dimethylaspartic acid (29-3, 1.08 g, 4.20 mmol, 1.50 equiv), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC, 0.805 g, 4.20 mmol, 1.50 equiv), 1-hydroxy-7-azabenzotriazole (0.571 g, 4.20 mmol, 1.50 equiv) and triethylamine (2.73 mL, 19.6 mmol, 7.00 equiv) were added, and the resulting mixture was stirred at 23°C for 20 hours. The reaction mixture was partitioned between water, acidified to pH 2, and ethyl acetate (2 x 100 mL). The combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by flash column chromatography through silica gel (hexane initially, grading to 100% EtOAc) followed by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide (3S)-3-[(tert-butoxycarbonyl)amino]-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid (29-4) as a white foam. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) major rotamer: δ 7.28 (m, 5H), 7.05 (m, 3H), 6.40 (s, 1H), 5.89 (br s, 1H), 5.74 (d, 1H, *J* = 10.2 Hz), 5.17 (d, 1H, *J* = 14.2 Hz), 4.93 (ddd, 1H, *J* = 13.9, 4.4, 1.7 Hz), 4.62 (d, 1H, *J* = 10.2 Hz), 1.45 (s, 9H), 1.30 (s, 3H), 1.05 (s, 3H). LRMS *m/z* (M+H) 501.5 found, 501.2 required.

Step 5: (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethyl-2,2-dimethyl-4-oxobutanamide (29-5)





A solution of (3S)-3-[(tert-butoxycarbonyl)amino]-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid (29-4, 50 mg, 0.10 mmol, 1.0 equiv), ethylamine hydrochloride (81 mg, 1.0 mmol, 10 equiv), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC, 29 mg, 0.15 mmol, 1.5 equiv), 1-hydroxy-7-azabenzotriazole (20 mg, 0.15 mmol, 1.5 equiv) and triethylamine (209 µL, 1.5 mmol, 15 equiv) in DMF (4 mL) was stirred at 23°C for 20 hours. The reaction mixture was partitioned between aqueous saturated sodium bicarbonate solution (50 mL) and ethyl acetate (2 x 50 mL). The combined organic layers were dried over sodium sulfate and concentrated. There




residue was purified by flash column chromatography (hexanes initially, grading to 100% EtOAc). A solution of the product was stirred in a 1:1 mixture of TFA and dichloromethane (2 mL) for 1 hour, then concentrated to provide (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethyl-2,2-dimethyl-4-oxobutanamide (29-5) as a white foam (TFA salt). LRMS  $m/z$  (M+H) 428.5 found, 428.2 required.

The following compounds were prepared by simple modifications of the above procedure. All of the compounds in the table below were isolated as the TFA salt.

10


Cmpd	Structure	Name	LRMS $m/z$ (M+H)
29-6		(3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxo-N-piperidin-4-ylbutanamide	$m/z$ (M+H) 483.6 found, 483.3 required
29-7		(3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid	$m/z$ (M+H) 401.4 found, 401.2 required

29-8		(3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2,2-tetramethyl-4-oxobutanamide	<i>m/z</i> (M+H) 428.5 found, 428.2 required
29-9		(1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethyl-3-oxo-3-piperazin-1-ylpropylamine	LRMS <i>m/z</i> (M+H) 469.5 found, 469.2 required
29-10		(3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropyl-2,2-dimethyl-4-oxobutanamide	<i>m/z</i> (M+H) 442.4 found, 442.2 required
29-11		(3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,2,2-trimethyl-4-oxobutanamide	<i>m/z</i> (M+H) 414.5 found, 414.2 required

29-12		(3R)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2,2-tetramethyl-4-oxobutanamide	<i>m/z</i> (M+H) 428.5 found, 428.2 required
29-13		(3R)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid	<i>m/z</i> (M+H) 401.4 found, 401.2 required
29-14		(1R)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethyl-3-oxo-3-piperazin-1-ylpropylamine	<i>m/z</i> (M+H) 469.5 found, 469.2 required



SCHEME 30



**Step 1:** ethyl ((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)acetate (30-1)


A solution of (1S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-2,2-dimethylpropylamine (16-1, 325 mg, 0.877 mmol, 1 equiv), ethyl bromoacetate (0.195 mL, 1.76 mmol, 2.00 equiv), and N,N-diisopropylethylamine (0.613 mL, 3.51 mmol, 4.00 equiv) in *n*-BuOH (4.0 mL) was heated under microwave irradiation at 150°C for 15 minutes. The reaction mixture was concentrated and the residue was partitioned between a ethyl acetate (100 mL) and a 1:1 mixture of saturated aqueous potassium bisulfate solution and brine (100 mL). The organic layer was then washed with saturated aqueous sodium bicarbonate solution (100 mL) followed by brine (100 mL), then dried over magnesium sulfate and concentrated to give ethyl ((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)acetate (30-1) as an off-white solid. LRMS  $m/z$  (M+H) 457.5 found, 457.2 required.




Step 2: 2-(({(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-ethylacetamide (30-2)




A solution of ethyl ({(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)acetate (30-1, 50 mg, 0.11 mmol) in a solution of ethylamine in THF (2 M, 2 mL) was heated in a sealable tube at 100°C for 16 hours. The reaction mixture was concentrated and the residue purified by flash column chromatography (EtOAc) to give 2-(({(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-ethylacetamide (30-2) as a white solid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) showed a 1:1 mixture of rotamers; one rotamer: δ 7.43-7.22 (m, 5H), 7.11-6.96 (m, 3H), 6.89 (br m, 1H), 6.38 (br s, 1H), 5.94 (m, 1H), 4.92 (ddd, 1H, *J* = 14.9, 4.4, 1.7 Hz), 4.87 (br d, 1H, *J* = 15.3 Hz), 3.42 (d, 1H, *J* = 16.8 Hz), 3.35 (m, 2H), 3.03 (s, 1H), 2.88 (d, 1H, *J* = 16.8 Hz), 2.34 (br s, 1H), 1.18 (t, 3H, *J* = 7.3 Hz), 1.06 (s, 9H). LRMS *m/z* (M+H) 456.5 found, 456.2 required.




15




The following compounds were prepared by simple modifications of the above procedure.




Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
30-3		2-(({(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-methylacetamide	LRMS <i>m/z</i> (M+H) 442.5 found, 442.2 required.




30-4		2-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)-N,N-dimethylacetamide	LRMS $m/z$ (M+H) 456.5 found, 456.2 required
30-5		2-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)-N-methyl-N-ethylacetamide	LRMS $m/z$ (M+H) 470.5 found, 470.2 required
30-6		2-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)-N-methylacetamide	LRMS $m/z$ (M+H) 426.4 found, 426.2 required

30-7		2-(((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)-N-ethylacetamide	LRMS $m/z$ (M+H) 440.4 found, 440.2 required
30-8		2-(((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)-N,N-dimethylacetamide	LRMS $m/z$ (M+H) 440.5 found, 440.2 required
30-9		2-(((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)-N-isopropylacetamide	LRMS $m/z$ (M+H) 454.5 found, 454.2 required

30-10		2-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)-N-ethyl-N-methylacetamide	LRMS <i>m/z</i> (M+H) 454.4 found, 454.2 required
30-11		2-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)-N,N-diethylacetamide	LRMS <i>m/z</i> (M+H) 468.5 found, 468.2 required
30-12		(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-N-(2-oxo-2-pyrrolidin-1-ylethyl)ethanamine	LRMS <i>m/z</i> (M+H) 466.5 found, 466.2 required


30-13		(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-(2-morpholin-4-yl-2-oxoethyl)-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 482.4 found, 482.2 required
30-14		1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]amino)acetyl]piperidin-4-ol	LRMS <i>m/z</i> (M+H) 496.6 found, 496.2 required
30-15		(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 495.5 found, 495.2 required

30-16		(1S)-N-(2-azetidin-1-yl)-2-oxoethyl-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 452.5 found, 452.2 required
30-17		(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-[2-(1,1-dioxidothiomorpholin-4-yl)-2-oxoethyl]-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 530.5 found, 530.2 required
30-18		(1S)-N-[2-(4-acetylpiperazin-1-yl)-2-oxoethyl]-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine	LRMS <i>m/z</i> (M+H) 423.6 found, 523.2 required

30-19		(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-(2-morpholin-4-yl-2-oxoethyl)-2-oxoethanamine	LRMS $m/z$ (M+H) 498.6 found, 498.2 required
30-20		(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-N-(2-oxo-2-pyrrolidin-1-ylethyl)ethanamine	LRMS $m/z$ (M+H) 482.6 found, 482.2 required
30-21		2-(({(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-isopropylacetamide	LRMS $m/z$ (M+H) 470.5 found, 470.2 required



SCHEME 31




**Step 1:** 4-nitrophenyl (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate (31-1)

- 5 A solution of 4-nitrophenyl chloroformate (188 mg, 0.931 mmol, 1.10 equiv) in dichloromethane (5 mL) was added dropwise to a solution of (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine (16-7, 300 mg, 0.847 mmol, 1 equiv) and triethylamine (0.354 mL, 2.54 mmol, 3.00 equiv) in dichloromethane (10 mL) at 0°C. The reaction mixture was warmed to 23°C and stirred for 20 hours. The reaction mixture was then partitioned between saturated aqueous sodium bicarbonate solution (50 mL) and ethyl acetate (75 mL). The organic layer was washed with brine, then dried over magnesium sulfate and concentrated to provide 4-nitrophenyl (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate (31-1) as a pale yellow solid. LRMS *m/z* (M+H) 520.5 found, 520.2 required.
- 10
- 15


Step 2: 2-(dimethylamino)ethyl (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate (31-2)

5 A solution of (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate (31-1, 50 mg, 0.096 mmol) in 2-(dimethylamino)ethanol (1 mL) was heated at 100°C for 1 hour. The reaction mixture was cooled, then partitioned between ethyl acetate (40 mL) and brine (3 x 40 mL). The organic layer was dried over magnesium sulfate and concentrated. The  
 10 residue was purified by flash column chromatography (EtOAc initially, grading to 10% saturated NH<sub>3</sub>/ethanol in EtOAc) to give 2-(dimethylamino)ethyl (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate (31-2) as a pale yellow solid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) showed a 2:1 mixture of rotamers; major rotamer: δ 7.42-7.24 (m, 5H), 7.11-6.95 (m, 3H),  
 15 6.41 (br s, 1H), 5.93 (m, 1H), 5.58 (d, 1H, *J* = 7.6 Hz), 5.04 (br d, 1H, *J* = 13.9 Hz), 4.91 (ddd, 1H, *J* = 14.7, 4.4, 1.7 Hz), 4.30 (t, 1H, *J* = 7.8 Hz), 4.15 (m, 2H), 2.54 (m, 2H), 2.27 (s, 6H), 1.14 (m, 1H), 0.57-0.33 (m, 4H). LRMS *m/z* (M+H) 470.5 found, 470.2 required.

20 The following compound was prepared by simple modifications of the above procedure.

Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
31-3		1-methylpiperidin-4-yl (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate	LRMS <i>m/z</i> (M+H) 496.6 found, 496.2 required.

SCHEME 32




5

Tert-butyl (2S)-2-phenyl-4-[[trifluoromethyl)sulfonyl]oxy}-2,5-dihydro-1H-pyrrole-1-carboxylate (5-4, 100 mg, 0.254 mmol, 1 equiv) was added to a solution of dry THF (5 mL) containing copper iodide (5 mg, 0.025 mmol, 0.2 equiv) at  $-10^{\circ}\text{C}$ . Cyclopropyl magnesium bromide (0.508 mL, 0.508 mmol, 2 equiv) was added drop wise and the resulting solution was stirred at  $-10^{\circ}\text{C}$  for 15 minutes. The reaction mixture was partitioned between aqueous ammonium chloride solution (15 mL) and ethyl acetate (3 x 10 mL). The combined organic layers were then dried over sodium sulfate, filtered, and concentrated to give tert-butyl (2S)-4-cyclopropyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (32-1), which is converted to (2S)-4-cyclopropyl-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (32-2) in the experimental mentioned above.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.22 (m, 5H), 5.82 (m, 1H), 5.35 (m, 1H), 4.43 (dd,  $J = 14.5, 5.5$  Hz, 1H), 4.00 (d,  $J = 14.5$  Hz, 1H), 2.82 (s, 6H), 1.40 (m, 1H), 0.76 (m, 2H), 0.54 (m, 2H). LRMS  $m/z$  ( $\text{M}+\text{H}$ ) 257.2 found, 257.3 required.


20

The following compound was prepared by simple modifications of the above procedure.

compd	Structure	Name	LRMS $m/z$ (M+H)
32-3		(2S)-4-cyclopentyl-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 285.3 found, 285.4 required.

5

SCHEME 33







10 To a solution of 1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanol (22-9) in dichloromethane at 0°C was added DMAP (1.0 mg, 0.006 mmol, 0.1 equiv), triethylamine (16  $\mu\text{L}$ , 0.113 mmol, 2 equiv), and para-nitrophenyl chloroformate (23 mg, 0.113 mmol, 2 equiv). The resulting mixture was warmed to 23 °C and stirred for 2 h. N-methyl piperazine  
 15 (23 mg, 0.225 mmol, 4.0 equiv) was added, and the resulting solution stirred another hour at 23 °C. Upon reaction completion, the reaction mixture was concentrated and purified by reverse phase liquid chromatography to yield (1S)-1-cyclopropyl-2-[(2S)-


4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 4-methylpiperazine-1-carboxylate (33-1) as the TFA salt. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  
 δ 7.35 (m, 5H), 7.05 (m, 3H), 6.40 (s, 1H), 5.98 (s, 1H), 5.17 (d, *J* = 13.4 Hz, 1H),  
 4.85 (m, 1H), 4.62 (m, 1H), 4.28 (m, 2H), 3.50 (m, 4H), 3.04 (m, 1H), 2.83 (s, 3H),  
 5 2.82 (m, 1H), 1.21 (m, 1H), 0.60 (m, 4H)

The following compounds were prepared by simple modifications of the above procedure. Unless otherwise indicated, the trifluoroacetate salt of the compound was isolated,


10

Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
33-2		1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 2-morpholin-4-ylethylcarbamate, isolated as the free base	LRMS <i>m/z</i> (M+H) 512.6 found, 512.7 required
33-3		N-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]oxy-carbonyl]glycine, isolated as the free base	LRMS <i>m/z</i> (M+H) 457.4 found, 457.4 required

33-4		(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 1-methylpiperidin-4-ylcarbamate	LRMS <i>m/z</i> (M+H) 496.5 found, 496.6 required
33-5		(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylmethyl(1-methylpiperidin-4-yl)carbamate	LRMS <i>m/z</i> (M+H) 510.5 found, 510.6 required

33-6		(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 4-(dimethylamino)piperidine-1-carboxylate	LRMS <i>m/z</i> (M+H) 510.6 found, 510.6 required
------	---	---	--

SCHEME 34



Tert-butyl (2S)-2-phenyl-4-[[[(trifluoromethyl)sulfonyl]oxy}-2,5-dihydro-1H-pyrrole-1-carboxylate (5-4, 1.6 g, 4.05 mmol) was dissolved in anhydrous dioxane (5 mL) and added to a degassed and N<sub>2</sub> purged solution of pinacol diborane (1.13 g, 4.46 mmol), potassium acetate (1.19 g, 12.2 mmol), bisdiphenylphosphine ferrocene (0.11 g, 0.2 mmol) and Pd(Cl)<sub>2</sub>(dppf) (0.15g, 0.2 mmol) in dioxane (50 mL). The resulting mixture was heated under N<sub>2</sub> at 80°C for 12 h. The reaction mixture was partitioned between water (100 mL) and ethyl acetate (3 x 100 mL). The combined organic layers were then washed with sat. aqueous NaCl, dried over sodium sulfate, filtered, and concentrated to give tert-butyl (2S)-2-phenyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2,5-dihydro-1H-pyrrole-1-carboxylate (34-1) which is used crude in the subsequent Suzuki coupling.



Crude 34-1 (0.364 g, 0.98 mmol) was dissolved in anhydrous DMF (10mL), degassed and filled with an N<sub>2</sub> atmosphere. 2,4-dichloro-5-fluoropyrimidine (0.25 g, 1.47 mmol), potassium carbonate (0.40 g, 2.94 mmol) and PdCl<sub>2</sub> (dppf) (36










mg, 0.05 mmol) were added to the solution and the mixture was stirred at 80°C for 2.5 h. The reaction mixture was partitioned between water (3 x 100 mL) and ethyl acetate (100 mL). The organic layer was then washed with sat. aqueous NaCl, dried over sodium sulfate, filtered, and concentrated. The crude oil was purified using reverse  
 5 phase HPLC (CH<sub>3</sub>CN/H<sub>2</sub>O/TFA 0-90% gradient) to provide tert-butyl (2S)-4-(2-chloro-5-fluoropyrimidin-4-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (34-2). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.47 (d, *J* = 2.4 Hz, 1H), 7.35 (m, 5H), 6.96 (m, 1H), 5.82 and 5.66 (m, 1H), 4.82 (m, 2H), 1.50 and 1.22 (s, 9H). LRMS: *m/z* (M-CH<sub>3</sub>) 361.3 found, 361.3 required.

10


The following compounds were prepared by simple modifications of the above procedure. Unless otherwise indicated, the compounds were isolated as the free base.

cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
34-3		(2S)-4-(5-fluoro-2-methylpyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 327.3 found, 327.4 required.
34-4		(2S)-4-(2-chloro-5-fluoropyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 347.7 found, 347.8 required.

34-5		(2S)-4-(4-chloro-5-methylpyrimidin-2-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 343.7 found, 343.8 required.
34-6		(2S)-4-(6-chloropyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 329.7 found, 329.8 required.
34-7		(2S)-4-(2-chloropyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 329.7 found, 329.8 required.
34-8		(2S)-N,N-dimethyl-4-(4-methylpyridin-3-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide, isolated as the TFA salt	LRMS $m/z$ (M+H) 308.3 found, 308.4 required.

34-9		(2S)-N,N-dimethyl-2-phenyl-4-(1,3-thiazol-2-yl)-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 300.3 found, 300.4 required.
34-10		(2S)-N,N-dimethyl-2-phenyl-4-(1,3-thiazol-4-yl)-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 300.3 found, 300.4 required.
34-11		(2S)-N,N-dimethyl-2-phenyl-4-(1,2-thiazol-5-yl)-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 300.3 found, 300.4 required.

SCHEME 35




- tert-Butyl 2-(3-([tert-butyl(dimethyl)silyl]oxy)phenyl)-4-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (35-1 [an intermediate in the preparation of
- 5 compound 17-6], 1.48 g, 3.82 mmol) was dissolved in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (25 mL) and treated with TFA (5 mL) for 1 hour. The reaction mixture was partitioned between sat. aq. NaHCO<sub>3</sub> (100 mL) and ethyl acetate (3 x 20 mL). The combined organic layers were then washed with sat. aqueous NaCl, dried over sodium sulfate, filtered, and concentrated. The crude amine was then dissolved in CH<sub>2</sub>Cl<sub>2</sub>, and treated
- 10 successively with Et<sub>3</sub>N (1.6 mL, 11.5 mmol) and dimethylcarbamoyl chloride (0.37 mL, 4.01 mmol), and the resulting solution was stirred 3.5 h at 25°C. Upon completion as judged by mass spectrometry, the reaction was concentrated, redissolved in CH<sub>2</sub>Cl<sub>2</sub> (30 mL) and triethylamine trihydrofluoride complex (0.6 mL, 3.82 mmol) was added and the reaction stirred an additional 12 h at 25°C. Upon
- 15 completion, the reaction was partitioned between NH<sub>4</sub>Cl (10 mL) and CH<sub>2</sub>Cl<sub>2</sub> (3 x 10

mL), and the combined organic layers were concentrated. The desired compound crystallized from CH<sub>2</sub>Cl<sub>2</sub> and as filtered to provide 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide 35-1a as a white solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.18 (t, *J* = 7.8 Hz, 2H), 7.00 (m, 3H),  
5 6.90 (s, 1H), 6.84 (d, *J* = 7.7 Hz, 1H), 6.74 (dd, *J* = 7.9, 2.4 Hz, 1H), 4.90 (dd, *J* = 13.4, 3.7 Hz, 1H), 4.50 (d, *J* = 13.4 Hz, 1H), 2.91 (s, 6H). LRMS: *m/z* (M-CH<sub>3</sub>) 345.3 found, 345.4 required.

Racemic 35-1a was subjected to chiral column chromatography  
10 (ChiralPak AD column, 4.6 mm x 265 mm, 10 micron, 85% hexanes/ 10% MeOH/ 5% EtOH, 60 mL/min) to resolve the enantiomers and yield (2*S*)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide 35-2.

## SCHEME 35A



5

2-2 (0.5 g, 1.78 mmol) was added to a microwave vessel and combined with tert-butyldimethylsilyl-3-iodophenol (1.78 g, 5.33 mmol, 3.0 equiv), tributylamine (1.0g, 5.33 mmol, 3.0 equiv), triphenylarsine (0.22 g, 0.71 mmol, 0.4




equiv), and palladium (II) acetate (0.08 g, 0.35 mmol, 0.2 equiv). The vessel was sealed, evacuated and charged with N<sub>2</sub>. Anhydrous 1,4-dioxane (1.5 mL) was added and the heterogenous mixture was again degassed and charged with N<sub>2</sub>. The vessel was placed in a Personal Chemistry microwave and heated at 150°C for 15 minutes.

- 5 Following completion, the reaction was diluted with EtOAc (50 mL) and washed sequentially with sat. aq. NH<sub>4</sub>Cl (50mL), water (50 mL) and brine (50mL). The combined organic extracts were dried with MgSO<sub>4</sub>, concentrated and subjected to flash column chromatography (SiO<sub>2</sub>, 100g, 0-15% EtOAc/hexanes gradient) to provide tert-Butyl 2-(3-{[tert-butyl(dimethyl)silyl]oxy}phenyl)-4-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate 35-1 as a clear oil: <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) major rotamer δ 7.17 (t, *J* = 7.6 Hz, 1H), 6.95 (m, 3H), 6.83 (d, *J* = 7.6 Hz, 1H), 6.73 (m, 2H), 6.31 (s, 1H), 5.50 (s, 1H), 4.66 (d, 2H), 1.27 (s, 9H), 0.97 (s, 9H), 0.18 (s, 6H). LRMS: *m/z* (M+H<sup>+</sup>) 488.5 found, 488.7 required.
- 10





- tert-Butyl 2-(3-{[tert-butyl(dimethyl)silyl]oxy}phenyl)-4-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate 35-1 (1.48 g, 3.82 mmol) was reacted as described in Scheme 35 to provide the racemic 35-1a, which was subjected to chiral column chromatography (ChiralPak AD column, 4.6 mm x 265 mm, 10 micron, 85% hexanes/10% MeOH/ 5% EtOH, 60 mL/min) to resolve the enantiomers and yield (2*S*)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide 35-2.
- 15
- 20





The following compounds were prepared by simple modifications of the above procedure or those earlier listed. Unless otherwise indicated, the compounds in the table were isolated as the free base.





25

Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
35-3		(2S)-4-(2,5-difluorophenyl)-N-(2-hydroxyethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 359.3 found, 359.4 required.
35-4		N-([4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-N-methyl-beta-alanine	LRMS <i>m/z</i> (M+H) 403.3 found, 403.4 required.
35-5		methyl N-([4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-N-methyl-beta-alaninate	LRMS <i>m/z</i> (M+H) 417.3 found, 417.4 required.



35-6		4-([4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]acetyl)-morpholin-4-ium; isolated as the trifluoroacetate salt	LRMS <i>m/z</i> (M+H) 401.3 found, 401.4 required.
35-7		3-[(2S)-4-(2,5-difluorophenyl)-1-(2-hydroxy-2-methylpropanoyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol	LRMS <i>m/z</i> (M+H) 360.3 found, 360.4 required.
35-8		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-sulfonamide	LRMS <i>m/z</i> (M+H) 381.3 found, 381.4 required.
35-9		3-[4-(2,5-difluorophenyl)-1-(methylsulfonyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol	LRMS <i>m/z</i> (M+H) 352.3 found, 352.4 required.




35-10		3-[4-(2,5-difluorophenyl)-1-(morpholin-4-ylcarbonyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol	LRMS $m/z$ (M+H) 387.3 found, 387.4 required.
35-11		3-[4-(2,5-difluorophenyl)-1-(2,2-dimethylpropanoyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol	LRMS $m/z$ (M+H) 358.3 found, 358.4 required.
35-12		(2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-tetrahydrofuran-3-yl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 401.5 found, 401.4 required.
35-13		(2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(2-methoxyethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 389.4 found, 389.4 required.




35-14		(2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-tetrahydro-2H-pyran-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 415.4 found, 415.5 required.
35-15		3-[(2S)-4-(2,5-difluorophenyl)-1-(piperidin-1-ylcarbonyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol	LRMS $m/z$ (M+H) 385.3 found, 385.4 required.
35-16		4-(2,5-difluorophenyl)-N-[1-(2-fluoroethyl)piperidin-4-yl]-2-(3-hydroxyphenyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 460.4 found, 460.6 required.
35-17		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 415.2 found, 415.5 required.

35-18		<p>2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}(methyl)amino]ethyl-4-methylpiperazine-1-carboxylate</p>	<p>LRMS <math>m/z</math> (M+H) 501.5 found, 501.6 required.</p>
35-19		<p>3-{4-(2,5-difluorophenyl)-1-[(4-methylpiperazin-1-yl)carbonyl]-2,5-dihydro-1H-pyrrol-2-yl}phenol</p>	<p>LRMS <math>m/z</math> (M+H) 400.4 found, 400.4 required.</p>
35-20		<p>2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}(methyl)amino]ethyl morpholine-4-carboxylate</p>	<p>LRMS <math>m/z</math> (M+H) 488.5 found, 488.5 required.</p>




35-21		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(1-methyl-1H-pyrazol-4-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 425.5 found, 425.4 required.
35-22		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(isoxazol-5-ylmethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 412.4 found, 412.4 required.
35-23		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(4-morpholinyl)-2-oxoethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 458.5 found, 458.5 required.




35-24		4-(2,5-difluorophenyl)-2,5-dihydro-2-(3-hydroxyphenyl)-N-methyl-N-[2-(4-methyl-1-piperazinyl)-2-oxoethyl]-H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 471.6 found, 471.5 required.
35-25		2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-(methyl)amino]ethyl dimethylamino-carboxylate	LRMS $m/z$ (M+H) 416.4 found, 416.4 required.
35-26		2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-(methyl)amino]ethyl piperidine-1-carboxylate	LRMS $m/z$ (M+H) 486.6 found, 486.5 required.




35-27		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 442.5 found, 442.5 required.
35-28		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(5-oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 428.4 found, 428.4 required.
35-29		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(tetrahydro-2H-pyran-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 429.5 found, 429.5 required.



35-30		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-[[5-(methoxymethyl)-1H-pyrazol-3-yl]methyl]-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 455.5 found, 455.5 required.
35-31		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 428.4 found, 428.5 required.
35-32		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(4-methyl-1,2,5-oxadiazol-3-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS <i>m/z</i> (M+H) 427.4 found, 427.4 required.




35-33		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-2-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 428.4 found, 428.5 required.
35-34		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(isoxazol-3-ylmethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 412.4 found, 412.4 required.
35-35		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-1,2,4-triazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 426.4 found, 426.4 required.

35-36		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-pyrazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 425.4 found, 425.5 required.
35-37		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(1-methyl-5-oxopyrrolidin-2-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 442.4 found, 442.5 required.
35-38		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(1-isoxazol-3-ylethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 426.4 found, 426.4 required.

35-39		4-(2,5-difluorophenyl)-N-(1,3-dioxolan-2-ylmethyl)-2-(3-hydroxyphenyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 417.4 found, 417.4 required.
35-40		4-(2,5-difluorophenyl)-N-(1,4-dioxan-2-ylmethyl)-2-(3-hydroxyphenyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 431.5 found, 431.5 required.
35-41		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(5-methyl-1,3,4-oxadiazol-2-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 427.4 found, 427.4 required.

35-42		4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(methylsulfonyl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide	LRMS $m/z$ (M+H) 437.4 found, 437.5 required.
35-43		2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]-carbonyl]-(methyl)amino]-ethane sulfonic acid	LRMS $m/z$ (M+H) 439.4 found, 439.5 required.

## SCHEME 36



5 Step 1: 4-nitrophenyl (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylcarbamate (36-1)




A solution of 4-nitrophenyl chloroformate (300 mg, 1.49 mmol, 1.2 equiv) in THF (10 mL) was added drop wise to a stirred solution of (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-carbonyl}-2,2-dimethylpropylamine (16-1, 460 mg, 1.24 mmol, 1.0 equiv) and triethylamine (151 mg, 1.49 mmol, 1.2 equiv) in THF (10 mL) at  $23^\circ\text{C}$ . After 1 hr the reaction was


filtered and the filtrate concentrated to a pale yellow gum which was purified by flash chromatography on silica gel. Elution with dichloromethane to 3 % ethyl acetate/dichloromethane provided 2-hydroxyethyl (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylcarbamate (36-1) as a colorless foam. LRMS  $m/z$  (M+H) 536.5 found, 535.2 required.

Step 2: 2-hydroxyethyl (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylcarbamate (36-2)

10 A mixture of 4-nitrophenyl (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylcarbamate (36-1, 500 mg, 0.93 mmol, 1.0 equiv) and ethylene glycol (3 mL, excess) was heated at 100 °C for 1 h. The reaction mixture was cooled and partitioned between ethyl acetate and water. The organic layer was washed with brine, dried over magnesium sulfate and  
15 concentrated. The residue was purified by flash chromatography on silica gel. Elution with 20 % ethyl acetate/dichloromethane to 50 % ethyl acetate/dichloromethane gave 2-hydroxyethyl (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylcarbamate (36-2) as a colorless foam. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) major rotamer: δ 7.38-7.21 (m, 5H), 7.11-6.94 (m, 3H), 6.40 (s, 1H), 5.92 (m, 1H), 5.51 (m, 1H), 5.07 (d, 1H,  $J$  = 13 Hz), 4.91 (m, 1H), 4.37 (d, 1H,  $J$  = 10 Hz), 4.20 (m, 2H), 3.81 (t, 2H,  $J$  = 4.5 Hz), 3.64 (t, 1H,  $J$  = 4 Hz), ) 0.91 (s, 9H). LRMS  $m/z$  (M+H) 459.3 found, 459.2 required.


The following compounds were prepared by simple modifications of the above  
25 procedures.

Cmpd	Structure	Name	LRMS $m/z$ (M+H)
36-3		3-hydroxypropyl (1 <i>S</i> )-1-[[ <i>(2S)</i> -4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]carbonyl]-2,2-dimethylpropyl-carbamate	LRMS $m/z$ (M+H) 473.5 found, 472.2 required.
36-4		2-hydroxyethyl {(1 <i>S</i> )-1-isopropyl-2-[( <i>2S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]-2-oxoethyl} carbamate	LRMS $m/z$ (M+H) 445.4 found, 445.2 required
36-5		2-hydroxyethyl {(1 <i>S</i> )-1-cyclopropyl-2-[( <i>2S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]-2-oxoethyl} carbamate	LRMS $m/z$ (M+H) 443.4 found, 443.2 required

36-6		4-hydroxybutyl (1S)-1- {[(2S)-4-(2,5- difluorophenyl)-2- phenyl-2,5-dihydro-1H- pyrrol-1-yl]carbonyl}- 2,2- dimethylpropylcarbamate	LRMS <i>m/z</i> (M+H) 487.4 found, 486.2 required
------	---	---	--

SCHEME 37

5



10 (2S)-4-(2,5-difluorophenyl)-1-[2-(methylsulfonyl)ethyl]-2-phenyl-2,5-dihydro-1H-pyrrole (37-1)

15 A solution of tert-butyl (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate (5-5, 100 mg, 0.280 mmol, 1 equiv) in a 4:1 mixture of dichloromethane and trifluoroacetic acid (20 mL) was stirred for 30 min, then concentrated. The residue was partitioned between saturated aqueous sodium bicarbonate solution and ethyl acetate (2 x 50 mL). The combined organic layers were dried over sodium sulfate and were concentrated. A solution of the residue and methyl vinyl sulfone (0.025 mL, 0.28 mmol, 1.0 equiv) in a 1:1 mixture of dioxane and ethanol (5 mL) was stirred at 23°C for 16 h. Additional methyl vinyl sulfone








- (0.025 mL, 0.28 mmol, 1.0 equiv) was added and stirring was continued for 4 h. The reaction mixture was concentrated and the residue was purified by flash column chromatography (hexanes initially, grading to 50% EtOAc in hexanes) to give (2*S*)-4-(2,5-difluorophenyl)-1-[2-(methylsulfonyl)ethyl]-2-phenyl-2,5-dihydro-1*H*-pyrrole (37-1) as a white solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.38-7.28 (m, 5H), 7.09-6.89 (m, 3H), 6.28 (br s, 1H), 4.69 (td, 1H, *J* = 4.9, 1.8 Hz), 4.34 (dd, 1H, *J* = 12.5, 4.9 Hz), 3.84 (ddd, 1H, *J* = 13.0, 4.8, 1.7 Hz), 3.36-3.02 (m, 4H), 2.79 (s, 3H). LRMS *m/z* (M+H) 364.3 found, 364.1 required.

10


The following compounds were prepared by simple modifications of the above procedure. Compounds 37-4 and 37-5 were isolated as TFA salts following purification by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present).

Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
37-2		(2 <i>S</i> )-4-(2,5-difluorophenyl)-1-[2-(ethylsulfonyl)ethyl]-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrole	LRMS <i>m/z</i> (M+H) 378.2 found, 378.1 required.
37-3		1-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]pentan-3-one	LRMS <i>m/z</i> (M+H) 342.4 found, 342.2 required.

37-4		4-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]butan-2-one	LRMS <i>m/z</i> (M+H) 328.2 found, 328.1 required.
37-5		4-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]-3-methylbutan-2-one	LRMS <i>m/z</i> (M+H) 342.3 found, 342.2 required.
37-6		2-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]- <i>N,N</i> -dimethylethanesulfonamide	LRMS <i>m/z</i> (M+H) 393.3 found, 393.1 required.
37-7		3-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-1-[2-(methylsulfonyl)ethyl]-2,5-dihydro-1 <i>H</i> -pyrrol-2-yl]phenol	LRMS <i>m/z</i> (M+H) 380.2 found, 380.4 required.

37-8		methyl 3-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]propanoate	LRMS <i>m/z</i> (M+H) 344.3 found, 344.1 required.
------	---	--	---

SCHEME 38



5

(2*S*)-4-(2,5-difluorophenyl)-1-[2-(ethylsulfonyl)propyl]-2-phenyl-2,5-dihydro-1*H*-pyrrole (38-1)

10 A solution of lithium bis(trimethylsilyl)amide in THF (1.0 M, 0.74 mL, 0.74 mmol, 4.0 equiv) was added to a solution of (2*S*)-4-(2,5-difluorophenyl)-1-[2-(ethylsulfonyl)ethyl]-2-phenyl-2,5-dihydro-1*H*-pyrrole (70 mg, 0.185 mmol, 1 equiv) in THF (5 mL) at  $-78^{\circ}\text{C}$ , then stirred for 1 h. Iodomethane (0.035 mL, 0.56 mmol, 3.0 equiv) was added and the resulting mixture was warmed to  $0^{\circ}\text{C}$  and stirred for 30


15 minutes. The mixture was then partitioned between saturated potassium bisulfate solution and ethyl acetate (2 x 40 mL). The combined organic layers were dried over magnesium sulfate and were concentrated. The residue was purified by flash column chromatography ( $\text{CH}_2\text{Cl}_2$  initially, grading to 4% EtOAc in  $\text{CH}_2\text{Cl}_2$ ) to give (2*S*)-4-(2,5-difluorophenyl)-1-[2-(ethylsulfonyl)propyl]-2-phenyl-2,5-dihydro-1*H*-pyrrole

20 (38-1) as a single diastereomer (colorless oil).  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  7.38-

7.28 (m, 5H), 7.08-6.92 (m, 3H), 6.29 (br s, 1H), 4.66 (br s), 4.37 (dd, 1H,  $J = 12.5$ , 4.9 Hz), 3.84 (br d, 1H,  $J = 13.0$  Hz), 3.36 (dd, 1H,  $J = 13.7$ , 7.1 Hz), 2.99 (m, 1H), 2.84 (m, 2H), 2.77 (dd, 1H,  $J = 13.9$ , 7.3 Hz), 1.31 (d, 3H,  $J = 7.1$  Hz), 1.19 (t, 3H,  $J = 7.6$  Hz). LRMS  $m/z$  (M+H) 392.4 found, 392.1 required.

5

## SCHEME 39



10 Step 1: 3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoic acid (39-1)


A solution of methyl 3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoate (37-8, 200 mg, 0.582 mmol, 1 equiv) and sodium hydroxide (1N, 1.16 mL, 1.16 mmol, 2.00 equiv) in t-BuOH (5 mL) was heated at 50 °C for 2 h. The reaction mixture was concentrated and the residue was partitioned between brine made acidic with aqueous 1 N HCL solution (1.2 mL) and ethyl acetate (2 x 40 mL). The combined organic layers were dried over sodium sulfate and concentrated to give 3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoic acid (39-1) as a white foam. LRMS  $m/z$  (M+H) 330.1 found, 330.3 required.




20



**Step 2:** 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-*N*-methylpropanamide (39-2)

A solution of 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoic acid (39-1, 50 mg, 0.15 mmol, 1 equiv), methylamine (2*M*, 1.0 mL, 2.0 mmol, 13 equiv), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC, 90 mg, 0.47 mmol, 3.0 equiv), 1-hydroxy-7-azabenzotriazole (62 mg, 0.47 mmol, 3.0 equiv) and triethylamine (0.32 mL, 2.3 mmol, 15 equiv) in DMF (3 mL) was heated in a sealable tube at 60 °C for 20 h. The reaction mixture was partitioned between saturated aqueous sodium bicarbonate solution (50 mL) and ethyl acetate (2 x 50 mL). The organic layer was dried over sodium sulfate and concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) to provide 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-*N*-methylpropanamide (39-2) as a colorless oil (TFA salt). <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.58-7.52 (m, 5H), 7.44-7.18 (m, 3H), 6.60 (br s, 1H), 5.72 (br s, 1H), 4.86 (br m, 1H), 4.64 (br m), 3.70 (br m, 2H), 2.72 (m, 2H), 2.68 (s, 3H). LRMS *m/z* (M+H) 343.3 found, 343.2 required.


The following compounds were prepared by simple modifications of the above procedure. All compounds were isolated as TFA salts.

Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
39-3		3-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]- <i>N,N</i> -dimethylpropanamide	LRMS <i>m/z</i> (M+H) 357.4 found, 357.2 required.

39-4		3-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]- <i>N,N</i> ,2-trimethylpropanamide	LRMS <i>m/z</i> (M+H) 371.4 found, 371.2 required.
39-5		4-{3-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]propanoyl}morpholine	LRMS <i>m/z</i> (M+H) 399.3 found, 399.2 required.
39-6		1-{3-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]propanoyl}-4-(methylsulfonyl)piperazine	LRMS <i>m/z</i> (M+H) 476.5 found, 476.2 required.

39-7		1-{3-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]propanoyl}piperidin-4-ol	LRMS <i>m/z</i> (M+H) 413.5 found, 413.2 required.
39-8		methyl 3-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]propanoate	LRMS <i>m/z</i> (M+H) 439.6 found, 439.2 required.

SCHEME 40



5 Step 1: ethyl ((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}oxy)acetate (40-1)

A solution of (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanol (22-9, 150 mg, 0.422 mmol, 1 equiv) in DMF (1 mL) was cooled to 0°C and treated with a solution of NaH (12 mg, 0.506 mmol, 1.2 eq) in DMF (0.5 ml), then stirred for 10 min. Ethyl bromoacetate (0.078 mL, 0.46 mmol, 1.1 eq) was added to the maroon-colored reaction mixture and the resulting mixture was stirred at 0 °C for 5 min, then warmed to 23 °C and stirred for 30 min. The yellow reaction mixture was partitioned between water (65 mL) and EtOAc (2 x 60 mL) and the combined organic layers were dried over sodium sulfate and concentrated. The residue was purified by reverse-phase liquid chromatography (acetonitrile/ H<sub>2</sub>O with 0.1% TFA present) to afford ethyl ((1S)-1-

10

15







cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}oxy)acetate (40-1) as a light yellow oil. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) major rotamer: δ 7.22-7.39 (m, 5H), 6.93-7.12 (m, 3H), 6.42 (br s), 5.87 (d, 1H, *J* = 2.4 Hz), 4.93 (br s, 2H), 4.29 (m, 2H), 3.92 (d, 1H, *J* = 7.6 Hz), 3.66 (d, 1H, *J* = 8.8 Hz), 3.41 (d, 1H, *J* = 8.8 Hz), 1.18 (t, 3H, *J* = 7.3 Hz), 0.61 (m, 4H). LRMS *m/z* (M+H) 442.4 found, 442.5 required.



**Step 2:** 2-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}oxy)-N-ethylacetamide (40-2)

10 A solution of ethyl ((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}oxy)acetate (40-1, 16 mg, 0.036 mmol) and ethylamine (10 equiv) in methanol (1 mL) was heated in a sealable tube at 75 deg C for 20 h. The reaction mixture was concentrated, then purified by reverse-phase liquid chromatography (acetonitrile/ H<sub>2</sub>O with 0.1% TFA present) to provide 2-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}oxy)-N-ethylacetamide (40-2) as a light yellow oil. LRMS *m/z* (M+H) 441.5 found, 441.5 required.


The following compounds were prepared by simple modifications of the above procedure. All compounds were isolated as TFA salts.

Cmpd	Structure	Name	LRMS <i>m/z</i> (M+H)
40-3		4-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethoxy)acetyl)morpholine	LRMS <i>m/z</i> (M+H) 483.5 found, 483.2 required.

40-4		2-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethoxy}-N-(2-hydroxyethyl)acetamide	LRMS $m/z$ (M+H) 457.4 found, 457.2 required.
40-5		1-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethoxy}acetyl)-4-methylpiperazine	LRMS $m/z$ (M+H) 496.6 found, 496.2 required.
40-6		1-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethoxy}-acetyl)piperazine	LRMS $m/z$ (M+H) 482.6 found, 482.2 required.

40-7		2-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethoxy}-N-piperidin-4-yl)acetamide	LRMS $m/z$ (M+H) 496.6 found, 496.2 required.
40-8		1-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethoxy}acetyl)piperidin-4-amine	LRMS $m/z$ (M+H) 496.6 found, 496.2 required.

SCHEME 41





- 5 Step 1: methyl *N*-{[(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl]}-β-alaninate (41-1)
- A solution of {[(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl}amine (16-7, 100 mg, 0.282 mmol, 1 equiv) and methyl acrylate (0.076 mL, 0.85 mmol, 3.0 equiv) in a 1:1 mixture of ethanol and dioxane (10 mL) was heated at 85 °C for 20 h. The reaction mixture was concentrated and the residue was purified by flash column chromatography (100% EtOAc) to provide methyl *N*-{[(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl]}-β-alaninate (41-1) as a colorless oil that solidified to a white solid over time. LRMS *m/z* (M+H) 441.4 found, 441.2
- 15 required.

Step 2:        *N*-{(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl}-3-morpholin-4-yl-3-oxopropan-1-amine (41-2)

---

5                A solution of methyl *N*-{(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl}-β-alaninate (40-1, 84 mg, 0.19 mmol, 1 equiv) and sodium hydroxide (1*N*, 0.57 mL, 0.57 mmol, 3.00 equiv) in *t*-BuOH (4 mL) was stirred at 23 deg C for 4 h. The reaction mixture was concentrated and the residue was partitioned between brine adjusted to pH 4 with  
10        aqueous 1 *N* HCL solution and ethyl acetate (2 x 40 mL). The combined organic layers were dried over sodium sulfate and concentrated. A solution of the residue, morpholine (0.083 mL, 0.95 mmol, 5.0 equiv), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC, 73 mg, 0.38 mmol, 2.0 equiv), and 1-hydroxy-7-azabenzotriazole (52 mg, 0.38 mmol, 2.0 equiv) in DMF (2 mL) was  
15        heated at 50 °C for 20 h. The reaction mixture was partitioned between aqueous saturated sodium bicarbonate solution and ethyl acetate (2 x 30 mL). The combined organic layers were dried over sodium sulfate and were concentrated. The residue was purified by reverse-phase LC (H<sub>2</sub>O/CH<sub>3</sub>CN gradient w/ 0.1 % TFA present) and the desired fractions partitioned between saturated sodium bicarbonate solution and  
20        ethyl acetate. The organic layer was dried over sodium sulfate and concentrated to provide *N*-{(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl}-3-morpholin-4-yl-3-oxopropan-1-amine (41-2) as a white foam. LRMS *m/z* (M+H) 496.5 found, 496.2 required.

The following compound was prepared by simple modification of the above procedure.

Cmpd	Structure	Name	LRMS $m/z$ (M+H)
41-3		$N^3$ -{((1 <i>S</i> )-1-cyclopropyl-2-[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]-2-oxoethyl)- $N^1,N^1$ -dimethyl- $\beta$ -alaninamide	LRMS $m/z$ (M+H) 454.5 found, 454.2 required.
41-4		((1 <i>S</i> )-1-{[(2 <i>S</i> )-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1 <i>H</i> -pyrrol-1-yl]carbonyl}-2,2-dimethylpropyl)(3-morpholin-4-yl-3-oxopropyl)amine	LRMS $m/z$ (M+H) 512.5 found, 512.3 required.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/18482

**Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☒ Claim Nos.: 1-10, 13-15 and parts of 11 and parts of 16-42  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
Please See Continuation Sheet
  
3. ☐ Claim Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest


☐  
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

## WHAT IS CLAIMED IS:

1. A compound of Formula I:



- 5 or a pharmaceutically acceptable salt or stereoisomer thereof, wherein

- a is 0 or 1;  
 b is 0 or 1;  
 m is 0, 1, or 2;  
 10 n is 0 or 1;  
 r is 0 or 1;  
 s is 0 or 1;  
 u is 2, 3, 4 or 5;
- 15 a dashed line represents an optional double bond, provided that one and only one double bond is present in the ring;

R¹ is selected from:

- 1) (C₁-C₆-alkylene)<sub>n</sub>(C=X)C₁-C₁₀ alkyl,  
 20 2) (C₁-C₆-alkylene)<sub>n</sub>(C=X)aryl,  
 3) (C₁-C₆-alkylene)<sub>n</sub>(C=X)C₂-C₁₀ alkenyl,  
 4) (C₁-C₆-alkylene)<sub>n</sub>(C=X)C₂-C₁₀ alkynyl,  
 5) (C₁-C₆-alkylene)<sub>n</sub>(C=X)C₃-C₈ cycloalkyl,  
 6) (C₁-C₆-alkylene)<sub>n</sub>(C=X)heterocyclyl,  
 25 7) (C₁-C₆-alkylene)<sub>n</sub>(C=X)NRᶜRᶜ',  
 8) (C₁-C₆-alkylene)<sub>n</sub>SO₂NRᶜRᶜ',  
 9) (C₁-C₆-alkylene)<sub>n</sub>SO₂C₁-C₁₀ alkyl,



- 10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkenyl,  
 11) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkynyl,  
 12) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-aryl,  
 13) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-heterocyclyl,  
 5 14) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
 15) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>P(=O)R<sup>d</sup>R<sup>d'</sup>,  
 16) aryl;  
 17) heterocyclyl; and  
 18) C<sub>1</sub>-C<sub>10</sub> alkyl;  
 10 said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, alkylene, heteroaryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>2</sup> and R<sup>6</sup> are independently selected from:

- 1) aryl,  
 15 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,  
 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and  
 4) heterocyclyl,

said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

20

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>8</sup>, and R<sup>9</sup> are independently selected from:

- 1) H,  
 2) C<sub>1</sub>-C<sub>10</sub> alkyl,  
 3) aryl,  
 25 4) C<sub>2</sub>-C<sub>10</sub> alkenyl,  
 5) C<sub>2</sub>-C<sub>10</sub> alkynyl,  
 7) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,  
 8) C<sub>1</sub>-C<sub>6</sub> aralkyl,  
 9) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and  
 30 10) heterocyclyl,

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>; or

R<sup>4</sup> and R<sup>5</sup>, or R<sup>8</sup> and R<sup>9</sup>, attached to the same carbon atom are combined to form

-(CH<sub>2</sub>)<sub>u</sub>- wherein one of the carbon atoms is optionally replaced by a moiety selected from O, S(O)<sub>m</sub>, -N(R<sup>a</sup>)C(O)-, -N(R<sup>b</sup>)- and -N(COR<sup>a</sup>)-;

R<sup>10</sup> is independently selected from:

- |    |  |
|----|--|
| 5  | 1) (C=O) <sub>a</sub> O <sub>b</sub> C <sub>1</sub> -C <sub>10</sub> alkyl,      |
|    | 2) (C=O) <sub>a</sub> O <sub>b</sub> aryl,                                       |
|    | 3) C <sub>2</sub> -C <sub>10</sub> alkenyl,                                      |
|    | 4) C <sub>2</sub> -C <sub>10</sub> alkynyl,                                      |
|    | 5) (C=O) <sub>a</sub> O <sub>b</sub> heterocyclyl,                               |
| 10 | 6) CO <sub>2</sub> H,  |
|    | 7) halo,   |
|    | 8) CN,   |
|    | 9) OH,   |
|    | 10) O <sub>b</sub> C <sub>1</sub> -C <sub>6</sub> perfluoroalkyl,                |
| 15 | 11) O <sub>a</sub> (C=O) <sub>b</sub> NR <sup>12</sup> R <sup>13</sup> ,         |
|    | 12) S(O) <sub>m</sub> R <sup>a</sup> ,   |
|    | 13) S(O) <sub>2</sub> NR <sup>12</sup> R <sup>13</sup> ,                         |
|    | 14) oxo,   |
|    | 15) CHO,   |
| 20 | 16) (N=O)R <sup>12</sup> R <sup>13</sup> , or                                    |
|    | 17) (C=O) <sub>a</sub> O <sub>b</sub> C <sub>3</sub> -C <sub>8</sub> cycloalkyl, |

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R<sup>11</sup>;

25 R<sup>11</sup> is selected from:

- |    |   |
|----|---|
| 30 | 1) (C=O) <sub>r</sub> O <sub>s</sub> (C <sub>1</sub> -C <sub>10</sub> )alkyl,   |
|    | 2) O <sub>r</sub> (C <sub>1</sub> -C <sub>3</sub> )perfluoroalkyl,              |
|    | 3) (C <sub>0</sub> -C <sub>6</sub> )alkylene-S(O) <sub>m</sub> R <sup>a</sup> , |
|    | 4) oxo,   |
|    | 5) OH,  |
|    | 6) halo,  |
|    | 7) CN,  |
|    | 8) (C=O) <sub>r</sub> O <sub>s</sub> (C <sub>2</sub> -C <sub>10</sub> )alkenyl, |
|    | 9) (C=O) <sub>r</sub> O <sub>s</sub> (C <sub>2</sub> -C <sub>10</sub> )alkynyl, |

- 10)  $(C=O)_rO_s(C_3-C_6)cycloalkyl$ ,
- 11)  $(C=O)_rO_s(C_0-C_6)alkylene-aryl$ ,
- 12)  $(C=O)_rO_s(C_0-C_6)alkylene-heterocyclyl$ ,
- 13)  $(C=O)_rO_s(C_0-C_6)alkylene-N(R^b)_2$ ,
- 5 14)  $C(O)R^a$ ,
- 15)  $(C_0-C_6)alkylene-CO_2R^a$ ,
- 16)  $C(O)H$ ,
- 17)  $(C_0-C_6)alkylene-CO_2H$ ,
- 18)  $C(O)N(R^b)_2$ ,
- 10 19)  $S(O)_mR^a$ , and
- 20)  $S(O)_2N(R^b)_2$

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $R^b$ , OH,  $(C_1-C_6)alkoxy$ , halogen,  $CO_2H$ , CN,  $O(C=O)C_1-C_6$  alkyl, oxo, and  $N(R^b)_2$ ;

15

$R^{12}$  and  $R^{13}$  are independently selected from:

- 1) H,
- 2)  $(C=O)O_bC_1-C_{10}$  alkyl,
- 3)  $(C=O)O_bC_3-C_8$  cycloalkyl,
- 20 4)  $(C=O)O_baryl$ ,
- 5)  $(C=O)O_bheterocyclyl$ ,
- 6)  $C_1-C_{10}$  alkyl,
- 7) aryl,
- 8)  $C_2-C_{10}$  alkenyl,
- 25 9)  $C_2-C_{10}$  alkynyl,
- 10) heterocyclyl,
- 11)  $C_3-C_8$  cycloalkyl,
- 12)  $SO_2R^a$ , and
- 13)  $(C=O)NR^b_2$ ,

30 said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from  $R^{11}$ , or

$R^{12}$  and  $R^{13}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and

optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from R<sup>11</sup>;

5 R<sup>14</sup> is independently selected from:

- 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 10 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 6) CO<sub>2</sub>H,
- 7) halo,
- 8) CN,
- 9) OH,
- 15 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
- 12) S(O)<sub>m</sub>R<sup>a</sup>,
- 13) S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
- 14) oxo,
- 20 15) CHO,
- 16) (N=O)R<sup>12</sup>R<sup>13</sup>, or
- 17) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R<sup>11</sup>;

25

R<sup>a</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl, optionally substituted with one to three substituents selected from R<sup>14</sup>;

R<sup>b</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl,  
 30 (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>, optionally substituted with one to three substituents selected from R<sup>14</sup>;

$R^c$  and  $R^{c'}$  are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from  $R^{10}$ , or

$R^c$  and  $R^{c'}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 3-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^{11}$ ;


$R^d$  and  $R^{d'}$  are independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy and  $NR^{b2}$ , or

$R^d$  and  $R^{d'}$  can be taken together with the phosphorous to which they are attached to form a monocyclic heterocycle with 5-7 members the ring and optionally containing, in addition to the phosphorous, one or two additional heteroatoms selected from  $NR^e$ , O and S, said monocyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^{11}$ ;

$R^e$  is selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl; and

X is selected from O,  $NR^e$  and S.

2. The compound according to Claim 1 of the Formula I:



or a pharmaceutically acceptable salt or stereoisomer thereof, wherein

a is 0 or 1;

- b is 0 or 1;  
 m is 0, 1, or 2;  
 n is 0 or 1;  
 r is 0 or 1;  
 5 s is 0 or 1;  
 u is 2, 3, 4 or 5;

a dashed line represents an optional double bond, provided that one and only one double bond is present in the ring;

10

R<sup>1</sup> is selected from:

- 1) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 2) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)aryl,
  - 3) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>2</sub>-C<sub>10</sub> alkenyl,
  - 15 4) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>2</sub>-C<sub>10</sub> alkynyl,
  - 5) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
  - 6) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)heterocyclyl,
  - 7) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)NR<sup>c</sup>R<sup>c'</sup>,
  - 8) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>,
  - 20 9) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkenyl,
  - 11) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkynyl,
  - 12) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-aryl,
  - 13) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-heterocyclyl,
  - 25 14) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
  - 15) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>P(=O)R<sup>d</sup>R<sup>d'</sup>,
  - 16) aryl;
  - 17) heterocyclyl; and
  - 18) C<sub>1</sub>-C<sub>10</sub> alkyl;
- 30 said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, alkylene, heteroaryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>2</sup> and R<sup>6</sup> are independently selected from:

- 1) aryl,
- 35 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,

3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and

4) heterocyclyl,

said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

5

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>8</sup>, and R<sup>9</sup> are independently selected from:

1) H,

2) C<sub>1</sub>-C<sub>10</sub> alkyl,

3) aryl,

10 4) C<sub>2</sub>-C<sub>10</sub> alkenyl,

5) C<sub>2</sub>-C<sub>10</sub> alkynyl,

6) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

7) C<sub>1</sub>-C<sub>6</sub> aralkyl,

8) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and

15 9) heterocyclyl,

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>; or

R<sup>4</sup> and R<sup>5</sup>, or R<sup>8</sup> and R<sup>9</sup>, attached to the same carbon atom are combined to form

20 -(CH<sub>2</sub>)<sub>u</sub>- wherein one of the carbon atoms is optionally replaced by a moiety selected from O, S(O)<sub>m</sub>, -N(R<sup>a</sup>)C(O)-, -N(R<sup>b</sup>)- and -N(COR<sup>a</sup>)-;

R<sup>10</sup> is independently selected from:

1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,

25 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,

3) C<sub>2</sub>-C<sub>10</sub> alkenyl,

4) C<sub>2</sub>-C<sub>10</sub> alkynyl,

5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,

6) CO<sub>2</sub>H,

30 7) halo,

8) CN,

9) OH,

10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,

- 12)  $S(O)_m R^a$ ,  
 13)  $S(O)_2 N R^{12} R^{13}$ ,  
 14) oxo,  
 15) CHO,  
 5 16)  $(N=O) R^{12} R^{13}$ , or  
 17)  $(C=O)_a O_b C_3-C_8$  cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from  $R^{11}$ ;

10  $R^{11}$  is selected from:

- 1)  $(C=O)_r O_s (C_1-C_{10})$  alkyl,  
 2)  $O_r (C_1-C_3)$  perfluoroalkyl,  
 3)  $(C_0-C_6)$  alkylene- $S(O)_m R^a$ ,  
 4) oxo,  
 15 5) OH,  
 6) halo,  
 7) CN,  
 8)  $(C=O)_r O_s (C_2-C_{10})$  alkenyl,  
 9)  $(C=O)_r O_s (C_2-C_{10})$  alkynyl,  
 20 10)  $(C=O)_r O_s (C_3-C_6)$  cycloalkyl,  
 11)  $(C=O)_r O_s (C_0-C_6)$  alkylene-aryl,  
 12)  $(C=O)_r O_s (C_0-C_6)$  alkylene-heterocyclyl,  
 13)  $(C=O)_r O_s (C_0-C_6)$  alkylene- $N(R^b)_2$ ,  
 14)  $C(O) R^a$ ,  
 25 15)  $(C_0-C_6)$  alkylene- $CO_2 R^a$ ,  
 16)  $C(O) H$ ,  
 17)  $(C_0-C_6)$  alkylene- $CO_2 H$ ,  
 18)  $C(O) N(R^b)_2$ ,  
 19)  $S(O)_m R^a$ , and  
 30 20)  $S(O)_2 N(R^b)_2$

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $R^b$ , OH,  $(C_1-C_6)$  alkoxy, halogen,  $CO_2 H$ , CN,  $O(C=O) C_1-C_6$  alkyl, oxo, and  $N(R^b)_2$ ;



R<sup>12</sup> and R<sup>13</sup> are independently selected from:

- 1) H,
  - 2) (C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 3) (C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
  - 5 4) (C=O)O<sub>b</sub>aryl,
  - 5) (C=O)O<sub>b</sub>heterocyclyl,
  - 6) C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 7) aryl,
  - 8) C<sub>2</sub>-C<sub>10</sub> alkenyl,
  - 10 9) C<sub>2</sub>-C<sub>10</sub> alkynyl,
  - 10) heterocyclyl,
  - 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
  - 12) SO<sub>2</sub>R<sup>a</sup>, and
  - 13) (C=O)NR<sup>b</sup><sub>2</sub>,
- 15 said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from R<sup>11</sup>, or

R<sup>12</sup> and R<sup>13</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and  
 20 optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from R<sup>11</sup>;

R<sup>a</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl;

25

R<sup>b</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>;

R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and  
 30 (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, optionally substituted with one, two or three substituents selected from R<sup>10</sup>, or

R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally

containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

- 5 R<sup>d</sup> and R<sup>d'</sup> are independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy and NR<sup>b</sup><sub>2</sub>, or


- R<sup>d</sup> and R<sup>d'</sup> can be taken together with the phosphorous to which they are attached to form a monocyclic heterocycle with 5-7 members the ring and optionally containing,  
 10 in addition to the phosphorous, one or two additional heteroatoms selected from NR<sup>e</sup>, O and S, said monocyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

R<sup>e</sup> is selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl; and

15

X is selected from O, NR<sup>e</sup> and S.

3. The compound according to Claim 2 of Formula I:



- 20 or a pharmaceutically acceptable salt or stereoisomer thereof, wherein

- a is 0 or 1;  
 b is 0 or 1;  
 m is 0, 1, or 2;  
 25 n is 0 or 1;  
 r is 0 or 1;  
 s is 0 or 1;

u is 2, 3, 4 or 5;

a dashed line represents an optional double bond, provided that one and only one double bond is present in the ring;

5

R<sup>1</sup> is selected from:

- 1) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 2) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)aryl,
  - 3) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>2</sub>-C<sub>10</sub> alkenyl,
  - 10 4) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>2</sub>-C<sub>10</sub> alkynyl,
  - 5) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
  - 6) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)heterocyclyl,
  - 7) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=X)NR<sup>c</sup>R<sup>c'</sup>,
  - 8) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>,
  - 15 9) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkenyl,
  - 11) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>2</sub>-C<sub>10</sub> alkynyl,
  - 12) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-aryl,
  - 13) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-heterocyclyl,
  - 20 14) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
  - 15) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>P(=O)R<sup>d</sup>R<sup>d'</sup>,
  - 16) aryl;
  - 17) heterocyclyl; and
  - 18) C<sub>1</sub>-C<sub>10</sub> alkyl;
- 25 said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, alkylene, heteroaryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>2</sup> and R<sup>6</sup> are independently selected from:

- 1) aryl,
- 30 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 4) heterocyclyl,

said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

35

R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>7</sup>, R<sup>8</sup>, and R<sup>9</sup> are independently selected from:

- 1) H,
- 2) C<sub>1</sub>-C<sub>10</sub> alkyl,
- 3) aryl,
- 5 4) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 5) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 6) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 7) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 8) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 10 9) heterocyclyl,

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>; or

- 15 R<sup>4</sup> and R<sup>5</sup>, or R<sup>8</sup> and R<sup>9</sup>, attached to the same carbon atom are combined to form -(CH<sub>2</sub>)<sub>u</sub>- wherein one of the carbon atoms is optionally replaced by a moiety selected from O, S(O)<sub>m</sub>, -N(R<sup>a</sup>)C(O)-, -N(R<sup>b</sup>)- and -N(COR<sup>a</sup>)-;

R<sup>10</sup> is independently selected from:

- 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 20 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 6) CO<sub>2</sub>H,
- 7) halo,
- 25 8) CN,
- 9) OH,
- 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
- 12) S(O)<sub>m</sub>R<sup>a</sup>,
- 30 13) S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
- 14) oxo,
- 15) CHO,
- 16) (N=O)R<sup>12</sup>R<sup>13</sup>, or
- 17) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R<sup>11</sup>;

R<sup>11</sup> is selected from:

- 5           1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl,
- 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl,
- 3) (C<sub>0</sub>-C<sub>6</sub>)alkylene-S(O)<sub>m</sub>R<sup>a</sup>,
- 4) oxo,
- 5) OH,
- 10          6) halo,
- 7) CN,
- 8) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>2</sub>-C<sub>10</sub>)alkenyl,
- 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>2</sub>-C<sub>10</sub>)alkynyl,
- 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,
- 15          11) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,
- 12) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-heterocyclyl,
- 13) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-N(R<sup>b</sup>)<sub>2</sub>,
- 14) C(O)R<sup>a</sup>,
- 15) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>R<sup>a</sup>,
- 20          16) C(O)H,
- 17) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>H,
- 18) C(O)N(R<sup>b</sup>)<sub>2</sub>,
- 19) S(O)<sub>m</sub>R<sup>a</sup>, and
- 20) S(O)<sub>2</sub>N(R<sup>b</sup>)<sub>2</sub>,

- 25   said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from R<sup>b</sup>, OH, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halogen, CO<sub>2</sub>H, CN, O(C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, oxo, and N(R<sup>b</sup>)<sub>2</sub>;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

- 30          1) H,
- 2) (C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 3) (C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 4) (C=O)O<sub>b</sub>aryl,
- 5) (C=O)O<sub>b</sub>heterocyclyl,

- 6) C<sub>1</sub>-C<sub>10</sub> alkyl,  
 7) aryl,  
 8) C<sub>2</sub>-C<sub>10</sub> alkenyl,  
 9) C<sub>2</sub>-C<sub>10</sub> alkynyl,  
 5 10) heterocyclyl,  
 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
 12) SO<sub>2</sub>R<sup>a</sup>, and  
 13) (C=O)NR<sup>b</sup><sub>2</sub>,

10 said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from R<sup>11</sup>, or

R<sup>12</sup> and R<sup>13</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from R<sup>11</sup>;

R<sup>a</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl;

20 R<sup>b</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>;

R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl or

25 R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with  
 30 one, two or three substituents selected from R<sup>11</sup>;


R<sup>d</sup> and R<sup>d'</sup> are independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy and NR<sup>b</sup><sub>2</sub>, or

$R^d$  and  $R^{d'}$  can be taken together with the phosphorous to which they are attached to form a monocyclic heterocycle with 5-7 members the ring and optionally containing, in addition to the phosphorous, one or two additional heteroatoms selected from  $NR^e$ , O and S, said monocyclic heterocycle optionally substituted with one, two or three substituents selected from  $R^{11}$ ;

$R^e$  is selected from: H and  $(C_1-C_6)alkyl$ ; and

X is selected from O,  $NR^e$  and S.

4. The compound according to Claim 2 of the Formula II,



or a pharmaceutically acceptable salt or stereoisomer thereof, wherein

wherein:

a is 0 or 1;

b is 0 or 1;

m is 0, 1, or 2;

r is 0 or 1;

s is 0 or 1;

a dashed line represents an optional double bond, provided that one and only one double bond is present in the ring;

$R^1$  is selected from:

- 1)  $(C_1-C_6-alkylene)_n(C=O)C_1-C_{10} alkyl$ ,
- 2)  $(C_1-C_6-alkylene)_n(C=O)aryl$ ,
- 3)  $(C_1-C_6-alkylene)_n(C=O)C_2-C_{10} alkenyl$ ,

- 4) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=O)C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=O)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 6) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=O)heterocyclyl,
- 7) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>(C=O)NR<sup>c</sup>R<sup>c'</sup>,
- 5 8) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>,
- 9) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 10) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-aryl,
- 11) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-heterocyclyl,
- 12) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>SO<sub>2</sub>-C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 10 13) (C<sub>1</sub>-C<sub>6</sub>-alkylene)<sub>n</sub>P(=O)R<sup>d</sup>R<sup>d'</sup>,
- 14) aryl,
- 15) heterocyclyl, and
- 16) C<sub>1</sub>-C<sub>10</sub> alkyl;

15 said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, alkylene, heteroaryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>2</sup> and R<sup>6</sup> are independently selected from:

- 1) aryl,
- 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 20 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 4) heterocyclyl,

said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

25 R<sup>3</sup>, R<sup>4</sup> and R<sup>8</sup> are independently selected from:

- 1) H,
- 2) C<sub>1</sub>-C<sub>10</sub> alkyl,
- 3) aryl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 30 5) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 6) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 7) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 8) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 9) heterocyclyl,



said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>10</sup> is independently selected from:

- 5           1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 10          6) CO<sub>2</sub>H,
- 7) halo,
- 8) CN,
- 9) OH,
- 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 15          11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
- 12) S(O)<sub>m</sub>R<sup>a</sup>,
- 13) S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
- 14) oxo,
- 15) CHO,
- 20          16) (N=O)R<sup>12</sup>R<sup>13</sup>, or
- 17) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

25 R<sup>11</sup> is selected from:

- 1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl,
- 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl,
- 3) oxo,
- 4) OH,
- 30          5) halo,
- 6) CN,
- 7) (C<sub>2</sub>-C<sub>10</sub>)alkenyl,
- 8) (C<sub>2</sub>-C<sub>10</sub>)alkynyl,
- 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,
- 35          10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,

- 11)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-heterocyclyl}$ ,  
 12)  $(\text{C}=\text{O})_r\text{O}_s(\text{C}_0\text{-C}_6)\text{alkylene-N(R}^b)_2$ ,  
 13)  $\text{C(O)R}^a$ ,  
 14)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{R}^a$ ,  
 5 15)  $\text{C(O)H}$ ,  
 16)  $(\text{C}_0\text{-C}_6)\text{alkylene-CO}_2\text{H}$ ,  
 17)  $\text{C(O)N(R}^b)_2$ ,  
 18)  $\text{S(O)}_m\text{R}^a$ , and  
 19)  $\text{S(O)}_2\text{N(R}^b)_2$ ;

10

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $\text{R}^b$ , OH,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ , halogen,  $\text{CO}_2\text{H}$ , CN,  $\text{O}(\text{C}=\text{O})\text{C}_1\text{-C}_6$  alkyl, oxo, and  $\text{N(R}^b)_2$ ;

15  $\text{R}^{12}$  and  $\text{R}^{13}$  are independently selected from:

- 1) H,  
 2)  $(\text{C}=\text{O})\text{O}_b\text{C}_1\text{-C}_{10}$  alkyl,  
 3)  $(\text{C}=\text{O})\text{O}_b\text{C}_3\text{-C}_8$  cycloalkyl,  
 4)  $(\text{C}=\text{O})\text{O}_b\text{aryl}$ ,  
 20 5)  $(\text{C}=\text{O})\text{O}_b\text{heterocyclyl}$ ,  
 6)  $\text{C}_1\text{-C}_{10}$  alkyl,  
 7) aryl,  
 8)  $\text{C}_2\text{-C}_{10}$  alkenyl,  
 9)  $\text{C}_2\text{-C}_{10}$  alkynyl,  
 25 10) heterocyclyl,  
 11)  $\text{C}_3\text{-C}_8$  cycloalkyl,  
 12)  $\text{SO}_2\text{R}^a$ , and  
 13)  $(\text{C}=\text{O})\text{NR}^b_2$ ,

30 said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from  $\text{R}^{11}$ , or

$\text{R}^{12}$  and  $\text{R}^{13}$  can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms

selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

R<sup>a</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl;

5

R<sup>b</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>;

10 R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;


15

R<sup>d</sup> and R<sup>d'</sup> are independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy and NR<sup>b</sup><sub>2</sub>, or

20 R<sup>d</sup> and R<sup>d'</sup> can be taken together with the phosphorous to which they are attached to form a monocyclic heterocycle with 5-7 members the ring and optionally containing, in addition to the phosphorous, one or two additional heteroatoms selected from NR<sup>e</sup>, O and S, said monocyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>; and

25 R<sup>e</sup> is selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

5. A compound of the Formula III,



or a pharmaceutically acceptable salt or stereoisomer thereof, wherein

- a is 0 or 1;  
 b is 0 or 1;  
 m is 0, 1, or 2;  
 5 r is 0 or 1;  
 s is 0 or 1;

a dashed line represents an optional double bond, provided that one and only one double bond is present in the ring;

10

R<sup>1</sup> is selected from:

- 1) (C=O)C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 2) (C=O)aryl,
  - 3) (C=O)C<sub>2</sub>-C<sub>10</sub> alkenyl,
  - 15 4) (C=O)C<sub>2</sub>-C<sub>10</sub> alkynyl,
  - 5) (C=O)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
  - 6) (C=O)heterocyclyl,
  - 7) (C=O)NR<sup>c</sup>RC<sup>c'</sup>,
  - 8) SO<sub>2</sub>NR<sup>c</sup>RC<sup>c'</sup>,
  - 20 9) SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 10) SO<sub>2</sub>-aryl,
  - 11) SO<sub>2</sub>-heterocyclyl,
  - 12) SO<sub>2</sub>-C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
  - 13) P(=O)R<sup>d</sup>R<sup>d'</sup>,
- 25 said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, heteroaryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>2</sup> is selected from:

- 1) aryl,
- 30 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 4) heterocyclyl,

said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

- 35 R<sup>3</sup>, R<sup>4</sup> and R<sup>8</sup> are independently selected from:

- 1) H,
  - 2) C<sub>1</sub>-C<sub>10</sub> alkyl,
  - 3) aryl,
  - 4) C<sub>2</sub>-C<sub>10</sub> alkenyl,
  - 5) C<sub>2</sub>-C<sub>10</sub> alkynyl,
  - 6) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
  - 7) C<sub>1</sub>-C<sub>6</sub> aralkyl,
  - 8) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
  - 9) heterocyclyl,
- 10 said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>10</sup> is independently selected from:

- 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 15 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 6) CO<sub>2</sub>H,
- 20 7) halo,
- 8) CN,
- 9) OH,
- 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,
- 11) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>12</sup>R<sup>13</sup>,
- 25 12) S(O)<sub>m</sub>R<sup>a</sup>,
- 13) S(O)<sub>2</sub>NR<sup>12</sup>R<sup>13</sup>,
- 14) oxo,
- 15) CHO,
- 16) (N=O)R<sup>12</sup>R<sup>13</sup>, or
- 30 17) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

R<sup>10'</sup> is halogen;

35

R<sup>11</sup> is selected from:

- 1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl,
- 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl,
- 3) oxo,
- 5 4) OH,
- 5) halo,
- 6) CN,
- 7) (C<sub>2</sub>-C<sub>10</sub>)alkenyl,
- 8) (C<sub>2</sub>-C<sub>10</sub>)alkynyl,
- 10 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl,
- 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl,
- 11) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-heterocyclyl,
- 12) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-N(R<sup>b</sup>)<sub>2</sub>,
- 13) C(O)R<sup>a</sup>,
- 15 14) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>R<sup>a</sup>,
- 15) C(O)H,
- 16) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>H, and
- 17) C(O)N(R<sup>b</sup>)<sub>2</sub>,
- 18) S(O)<sub>m</sub>R<sup>a</sup>, and
- 20 19) S(O)<sub>2</sub>N(R<sup>b</sup>)<sub>2</sub>;

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heterocyclyl is optionally substituted with up to three substituents selected from R<sup>b</sup>, OH, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halogen, CO<sub>2</sub>H, CN, O(C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, oxo, and N(R<sup>b</sup>)<sub>2</sub>;

25

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

- 1) H,
- 2) (C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 3) (C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 30 4) (C=O)O<sub>b</sub>aryl,
- 5) (C=O)O<sub>b</sub>heterocyclyl,
- 6) C<sub>1</sub>-C<sub>10</sub> alkyl,
- 7) aryl,
- 8) C<sub>2</sub>-C<sub>10</sub> alkenyl,

- 9) C<sub>2</sub>-C<sub>10</sub> alkynyl,  
 10) heterocyclyl,  
 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,  
 12) SO<sub>2</sub>R<sup>a</sup>, and  
 5 13) (C=O)NR<sup>b</sup><sub>2</sub>.

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R<sup>11</sup>, or

- R<sup>12</sup> and R<sup>13</sup> can be taken together with the nitrogen to which they are attached to  
 10 form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

- 15 R<sup>a</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, or heterocyclyl;

R<sup>b</sup> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>;

- 20 R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or

- R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form  
 25 a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

- R<sup>d</sup> and R<sup>d'</sup> are independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy and  
 30 NR<sup>b</sup><sub>2</sub>, or


R<sup>d</sup> and R<sup>d'</sup> can be taken together with the phosphorous to which they are attached to form a monocyclic heterocycle with 5-7 members the ring and optionally containing, in addition to the phosphorous, one or two additional heteroatoms selected from NR<sup>e</sup>,

O and S, said monocyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>; and

R<sup>e</sup> is selected from: H and (C<sub>1</sub>-C<sub>6</sub>)alkyl.

5

6. The compound according to Claim 4 of the Formula IV,



or a pharmaceutically acceptable salt or stereoisomer thereof, wherein

- 10 a is 0 or 1;  
 b is 0 or 1;  
 m is 0, 1, or 2;  
 r is 0 or 1;  
 s is 0 or 1;

15

R<sup>1</sup> is selected from:

- 1) (C=O)C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)aryl,
- 3) (C=O)C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 20 4) (C=O)heterocyclyl,
- 5) (C=O)NR<sup>c</sup>R<sup>c'</sup>,
- 6) (C=S)NR<sup>c</sup>R<sup>c'</sup>,
- 7) SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>,
- 8) SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 25 9) SO<sub>2</sub>-aryl, and
- 10) SO<sub>2</sub>-heterocyclyl,

said alkyl, aryl, cycloalkyl, and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;



R<sup>2</sup> is selected from:

- 1) aryl,
- 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 5 4) heterocyclyl,

said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>3</sup>, R<sup>4</sup> and R<sup>8</sup> are independently selected from:

- 10 1) H,
- 2) C<sub>1</sub>-C<sub>10</sub> alkyl, and
- 3) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

said alkyl, aryl, alkenyl, alkynyl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

15

R<sup>6</sup> is selected from:

- 1) aryl,
- 2) C<sub>1</sub>-C<sub>6</sub> aralkyl,
- 3) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and
- 20 4) heterocyclyl,

said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>10</sup> is independently selected from:

- 25 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl,
- 3) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 4) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl,
- 30 6) CO<sub>2</sub>H,
- 7) halo,
- 8) CN,
- 9) OH,
- 10) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl,

- 11)  $O_a(C=O)_bNR^{12}R^{13}$ ,  
 12)  $S(O)_mR^a$ ,  
 13)  $S(O)_2NR^{12}R^{13}$ ,  
 14) oxo,  
 5 15) CHO,  
 16)  $(N=O)R^{12}R^{13}$ , or  
 17)  $(C=O)_aO_bC_3-C_8$  cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from  $R^{11}$ ;

10

$R^{11}$  is selected from:

- 1)  $(C=O)_rO_s(C_1-C_{10})$ alkyl,  
 2)  $O_r(C_1-C_3)$ perfluoroalkyl,  
 3) oxo,  
 15 4) OH,  
 5) halo,  
 6) CN,  
 7)  $(C_2-C_{10})$ alkenyl,  
 8)  $(C_2-C_{10})$ alkynyl,  
 20 9)  $(C=O)_rO_s(C_3-C_6)$ cycloalkyl,  
 10)  $(C=O)_rO_s(C_0-C_6)$ alkylene-aryl,  
 11)  $(C=O)_rO_s(C_0-C_6)$ alkylene-heterocyclyl,  
 12)  $(C=O)_rO_s(C_0-C_6)$ alkylene- $N(R^b)_2$ ,  
 13)  $C(O)R^a$ ,  
 25 14)  $(C_0-C_6)$ alkylene- $CO_2R^a$ ,  
 15)  $C(O)H$ ,  
 16)  $(C_0-C_6)$ alkylene- $CO_2H$ ,  
 17)  $C(O)N(R^b)_2$ ,  
 18)  $S(O)_mR^a$ , and  
 30 19)  $S(O)_2N(R^b)_2$ ;

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heterocyclyl is optionally substituted with up to three substituents selected from  $R^b$ , OH,  $(C_1-C_6)$ alkoxy, halogen,  $CO_2H$ , CN,  $O(C=O)C_1-C_6$  alkyl, oxo, and  $N(R^b)_2$ ;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

- 1) H,
- 2) (C=O)O<sup>b</sup>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 5 3) (C=O)O<sup>b</sup>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 4) (C=O)O<sup>b</sup>aryl,
- 5) (C=O)O<sup>b</sup>heterocyclyl,
- 6) C<sub>1</sub>-C<sub>10</sub> alkyl,
- 7) aryl,
- 10 8) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 9) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 10) heterocyclyl,
- 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 12) SO<sub>2</sub>R<sup>a</sup>, and
- 15 13) (C=O)NR<sup>b</sup><sub>2</sub>,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R<sup>11</sup>, or

R<sup>12</sup> and R<sup>13</sup> can be taken together with the nitrogen to which they are attached to  
 20 form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

25 R<sup>a</sup> is independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, and heterocyclyl;

R<sup>b</sup> is independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>; and

30 R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl or

R<sup>c</sup> and R<sup>c'</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with  
5 one, two or three substituents selected from R<sup>11</sup>.

7. The compound according to Claim 6 or a pharmaceutically acceptable salt or stereoisomer thereof, wherein:

10 R<sup>1</sup> is selected from:

- 1) (C=O)NR<sup>c</sup>R<sup>c'</sup>,
- 2) SO<sub>2</sub>NR<sup>c</sup>R<sup>c'</sup>, and
- 3) SO<sub>2</sub>C<sub>1</sub>-C<sub>10</sub> alkyl,

said alkyl, is optionally substituted with one, two or three substituents selected from  
15 R<sup>10</sup>;

R<sup>2</sup> is selected from:

- 1) aryl, and
- 2) heteroaryl,

20 said aryl and heteroaryl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>3</sup>, R<sup>4</sup> and R<sup>8</sup> are independently selected from:

- 1) H, and
- 25 2) C<sub>1</sub>-C<sub>10</sub> alkyl,

said alkyl is optionally substituted with one or more substituents selected from R<sup>10</sup>;

R<sup>6</sup> is selected from:

- 1) aryl, and
- 30 2) heterocyclyl,

said alkyl, aryl and heterocyclyl is optionally substituted with one or more substituents selected from R<sup>10</sup>; and

R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, R<sup>a</sup>, R<sup>b</sup>, R<sup>c</sup> and R<sup>c'</sup> are as described immediately above.


8. The compound according to Claim 7 or a pharmaceutically acceptable salt or stereoisomer thereof, wherein:

- 5  $R^2$  and  $R^6$  are independently selected from phenyl or pyridyl, optionally substituted with one or two substituents selected from  $R^{10}$ ; and  
 $R^1$ ,  $R^3$ ,  $R^4$  and  $R^8$  are as described in Claim 7.

9. The compound according to Claim 7, or the pharmaceutically acceptable salt or stereoisomer thereof, wherein  $R^2$  is phenyl, optionally substituted with one or two substituents selected from  $R^{10}$ ; and

$R^1$ ,  $R^3$ ,  $R^4$ ,  $R^6$  and  $R^8$  are as described in Claim 7.

10. The compound according to Claim 5 of the Formula V,



wherein:

- a is 0 or 1;  
 20 b is 0 or 1;  
 m is 0, 1, or 2;  
 r is 0 or 1;  
 s is 0 or 1;
- 25  $R^1$  is selected from:
- 1)  $(C=O)C_1-C_{10}$  alkyl,
  - 2)  $(C=O)$ aryl,
  - 3)  $(C=O)C_3-C_8$  cycloalkyl,
  - 4)  $(C=O)$ heterocyclyl,

- 5)  $(C=O)NR^cR^c'$ ,  
 6)  $(C=S)NR^cR^c'$ ,  
 7)  $SO_2NR^cR^c'$ ,  
 8)  $SO_2C_1-C_{10}$  alkyl,  
 5 9)  $SO_2$ -aryl, and  
 10)  $SO_2$ -heterocyclyl,

said alkyl, aryl, cycloalkyl, and heterocyclyl is optionally substituted with one or more substituents selected from  $R^{10}$ ;

10  $R^2$  is selected from:

- 1) aryl,  
 2)  $C_1-C_6$  aralkyl,  
 3)  $C_3-C_8$  cycloalkyl, and  
 4) heterocyclyl,

15 said aryl, cycloalkyl, aralkyl and heterocyclyl is optionally substituted with one or more substituents selected from  $R^{10}$ ;

$R^3$ ,  $R^4$  and  $R^8$  are independently selected from:

- 1) H,  
 20 2)  $C_1-C_{10}$  alkyl, and  
 3)  $C_1-C_6$  perfluoroalkyl,

said alkyl is optionally substituted with one or more substituents selected from  $R^{10}$ ;

$R^{10}$  is independently selected from:

- 25 1)  $(C=O)_aO_bC_1-C_{10}$  alkyl,  
 2)  $(C=O)_aO_b$ aryl,  
 3)  $C_2-C_{10}$  alkenyl,  
 4)  $C_2-C_{10}$  alkynyl,  
 5)  $(C=O)_aO_b$  heterocyclyl,  
 30 6)  $CO_2H$ ,  
 7) halo,  
 8)  $CN$ ,  
 9)  $OH$ ,  
 10)  $O_bC_1-C_6$  perfluoroalkyl,

- 11)  $O_a(C=O)_bNR^{12}R^{13}$ ,  
 12)  $S(O)_mR^a$ ,  
 13)  $S(O)_2NR^{12}R^{13}$ ,  
 14) oxo,  
 5 15) CHO,  
 16)  $(N=O)R^{12}R^{13}$ , or  
 17)  $(C=O)_aO_bC_3-C_8$  cycloalkyl,

said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from  $R^{11}$ ;

10  $R^{10'}$  is halogen;

$R^{11}$  is selected from:

- 1)  $(C=O)_rO_s(C_1-C_{10})$ alkyl,  
 2)  $O_r(C_1-C_3)$ perfluoroalkyl,  
 15 3) oxo,  
 4) OH,  
 5) halo,  
 6) CN,  
 7)  $(C_2-C_{10})$ alkenyl,  
 20 8)  $(C_2-C_{10})$ alkynyl,  
 9)  $(C=O)_rO_s(C_3-C_6)$ cycloalkyl,  
 10)  $(C=O)_rO_s(C_0-C_6)$ alkylene-aryl,  
 11)  $(C=O)_rO_s(C_0-C_6)$ alkylene-heterocyclyl,  
 12)  $(C=O)_rO_s(C_0-C_6)$ alkylene- $N(R^b)_2$ ,  
 25 13)  $C(O)R^a$ ,  
 14)  $(C_0-C_6)$ alkylene- $CO_2R^a$ ,  
 15)  $C(O)H$ ,  
 16)  $(C_0-C_6)$ alkylene- $CO_2H$ ,  
 17)  $C(O)N(R^b)_2$ ,  
 30 18)  $S(O)_mR^a$ , and  
 19)  $S(O)_2N(R^b)_2$ ;

said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, alkylene and heterocyclyl is optionally substituted with up to three substituents selected from  $R^b$ , OH,  $(C_1-C_6)$ alkoxy, halogen,  $CO_2H$ , CN,  $O(C=O)C_1-C_6$  alkyl, oxo, and  $N(R^b)_2$ ;

R<sup>12</sup> and R<sup>13</sup> are independently selected from:

- 1) H,
- 2) (C=O)O<sup>b</sup>C<sub>1</sub>-C<sub>10</sub> alkyl,
- 5 3) (C=O)O<sup>b</sup>C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 4) (C=O)O<sup>b</sup>aryl,
- 5) (C=O)O<sup>b</sup>heterocyclyl,
- 6) C<sub>1</sub>-C<sub>10</sub> alkyl,
- 7) aryl,
- 10 8) C<sub>2</sub>-C<sub>10</sub> alkenyl,
- 9) C<sub>2</sub>-C<sub>10</sub> alkynyl,
- 10) heterocyclyl,
- 11) C<sub>3</sub>-C<sub>8</sub> cycloalkyl,
- 12) SO<sub>2</sub>R<sup>a</sup>, and
- 15 13) (C=O)NR<sup>b</sup><sub>2</sub>,

said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one, two or three substituents selected from R<sup>11</sup>, or

R<sup>12</sup> and R<sup>13</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one, two or three substituents selected from R<sup>11</sup>;

R<sup>a</sup> is independently selected from: (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, aryl, and heterocyclyl;

R<sup>b</sup> is independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>; and

R<sup>c</sup> and R<sup>c'</sup> are independently selected from: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, heterocyclyl and (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl or



RC and RC' can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 5-7 members in each ring and optionally containing, in addition to the nitrogen, one or two additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with  
 5 one, two or three substituents selected from R<sup>11</sup>.

11. A compound selected from:

- 10 4-(2-chloro-5-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (+)-4-(2,5-difluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 (-)-4-(2,5-difluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(5-chloro-2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 4-(2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2-fluoro-5-methylphenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 4-(5-bromo-2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-{{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}morpholine};
- 30 4-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}morpholine};
- N,N-dimethyl-2,4-diphenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

- 3-[2-fluoro-5-(trifluoromethyl)phenyl]-N,N-dimethyl-5-phenyl-2,3-dihydro-1H-pyrrole-1-carboxamide;
- 5 2-(3-fluorophenyl)-4-(2,5-difluorophenyl)- N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(4-fluorophenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 4-(2,5-difluorophenyl)-2-(2-fluorophenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 2-(3-bromophenyl)-4-(2,5-difluorophenyl)- N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 2-(3-aminophenyl)-4-(2,5-difluorophenyl)- N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 4-(2,5-difluorophenyl)-2-(3-methylphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(5-chloro-2-fluorophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 4-(2,5-difluorophenyl)-1-(methylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-1-(ethylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 30 4-(2,5-difluorophenyl)-2-phenyl-1-(propylsulfonyl)-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-1-(isopropylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-(5-chloro-2-fluorophenyl)-1-(methylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 35

- 4-(5-chloro-2-fluorophenyl)-1-(isopropylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-(2-fluoro-5-methylphenyl)-1-(isopropylsulfonyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 5 2-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]sulfonyl}ethanamine;
- 2-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]sulfonyl}-N,N-dimethylethanamine;
- 10 1-acetyl-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-(2-chloro-5-fluorophenyl)-1-pivaloyl-2-phenyl-2,5-dihydro-1H-pyrrole;
- 15 4-(2-chloro-5-fluorophenyl)-1-isobutyryl-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-1-(2,2-dimethylpropanoyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-ol;
- 20 1-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-ol;
- 25 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine;
- 4-(2-fluoro-5-isocyanophenyl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 4-(2,5-difluorophenyl)-2-phenyl-1-(trifluoroacetyl)-2,5-dihydro-1H-pyrrole;
- 4-(5-chloro-2-fluorophenyl)-2-phenyl-1-(trifluoroacetyl)-2,5-dihydro-1H-pyrrole;

- (1S)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-2-methylpropylamine;
- (1R)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-2-methylpropylamine;
- 5 4-(2,5-difluorophenyl)-2-phenyl-1-L-prolyl-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-2-phenyl-1-D-prolyl-2,5-dihydro-1H-pyrrole;
- 10 (4R)-4-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-1,3-thiazolidine;
- methyyl (3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutanoate;
- 15 (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanamide;
- 20 (1S)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-3-(methylthio)propylamine;
- (1S)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-3-(methylsulfonyl)propylamine;
- 25 (2S)-2-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}piperidine;
- (1S)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}pentylamine;
- 30 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(thien-2-ylmethyl)ethylamine;

- 4- {[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-1,1-dioxidotetrahydro-2H-thiopyran-4-ylamine;
- 5 (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methyl-1-oxopropan-2-amine;
- (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}propylamine;
- 10 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-phenylethanamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-phenylethanamine;
- 15 (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanamide
- 3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-amine;
- 20 (1S,2S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylbutylamine;
- (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}butylamine;
- 25 (1S)-1-cyclopropyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 30 1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopropanamine;
- 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-oxopropan-2-amine;
- 35

- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-methyl-2-oxoethylamine;
- 5 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(pyridin-2-ylmethyl)ethylamine;
- (1S)-1-cyclohexyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 10 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(4-iodobenzyl)-2-oxoethylamine;
- (1S)-1-benzyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine;
- 15 4-{(2S)-2-amino-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropyl}phenol;
- (3S)-3-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-1,2,3,4-tetrahydroisoquinoline;
- 20 (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-phenylpropylamine;
- (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-methylbutylamine;
- 25 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(pyridin-3-ylmethyl)ethylamine;
- 30 1-[(2S)-azetidin-2-ylcarbonyl]-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- (3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutanamide;
- 35

4-(2,5-difluorophenyl)-1-[(2S)-2,5-dihydro-1H-pyrrol-2-ylcarbonyl]-2-phenyl-2,5-dihydro-1H-pyrrole;

5 4-(2,5-difluorophenyl)-1-[(2-methylazetidin-2-yl)carbonyl]-2-phenyl-2,5-dihydro-1H-pyrrole;

(1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylamine;  
methyl (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanoate;  
10

4-(2,5-difluorophenyl)-2-phenyl-1-[[2S,3S)-2-phenylpyrrolidin-3-yl]carbonyl]-2,5-dihydro-1H-pyrrole;

15 4-(2,5-difluorophenyl)-2-phenyl-1-[(5-phenylpyrrolidin-3-yl)carbonyl]-2,5-dihydro-1H-pyrrole;

20 (2S)-2-amino-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-ol;

(2R,3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutan-2-ol;

(1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(methoxymethyl)-2-oxoethylamine;  
25

4-(2,5-difluorophenyl)-2-phenyl-1-(pyrrolidin-3-ylcarbonyl)-2,5-dihydro-1H-pyrrole;

4-(2,5-difluorophenyl)-2-phenyl-1-[(3-phenylpyrrolidin-3-yl)acetyl]-2,5-dihydro-1H-pyrrole;  
30

(1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3,3-difluoropropylamine;

- (1S)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxo-1-phenylpropan-1-amine;
- 4-(2,5-difluorophenyl)-2-phenyl-1-[(4S)-4-phenyl-L-prolyl]-2,5-dihydro-1H-pyrrole;
- 5 1-{2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}cyclohexanamine;
- 2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 10 4-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}piperidin-4-amine;
- (1S,3R)-3-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopentanamine;
- 15 (1R,4S)-4-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-2-en-1-amine;
- (1S,4R)-4-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-2-en-1-amine;
- 20 (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-ynylamine;
- (1R)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxo-1-phenylpropan-1-amine;
- 25 3-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-phenylpiperidine;
- 30 (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-enylamine;
- (2S)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-(methylamino)-3-oxopropan-1-ol;
- 35



- (3R,5S)-5-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}pyrrolidin-3-ol;
- 5 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(1,3-thiazol-4-ylmethyl)ethylamine;
- (1R)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}but-3-enylamine;
- 10 (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,3-dimethyl-1-oxobutan-2-amine;
- (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,4-dimethyl-1-oxopentan-2-amine;
- 15 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-[(1-methyl-1H-imidazol-4-yl)methyl]-2-oxoethylamine;
- 4-(2,5-difluorophenyl)-1-(N-6-formyl-L-lysyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 20 (2S,3S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,3-dimethyl-1-oxopentan-2-amine;
- (1S)-1-(cyclohexylmethyl)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine;
- 25 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxoethylamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(isocyanomethyl)-2-oxoethylamine;
- 30 (1S)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-3,3-dimethylbutylamine;
- 35

- 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,3-dimethyl-1-oxobutan-2-amine;
- 1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclohexanamine;
- 1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopentanamine;
- (1S)-3-(benzyloxy)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}propylamine;
- 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,3-dimethyl-1-oxobutan-2-amine;
- 1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-3-en-1-amine;
- (1S)-1-cyclopentyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 4-(2,5-difluorophenyl)-1-(2-methylpropyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 1-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine;
- (1S)-1-{{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylpropylamine;
- (1S)-2-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-cyclopropyl-2-oxoethanamine;
- (1S,2S)-1-{{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylbutylamine;

(1S)-1-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}pentylamine;

(1S)-1-{{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-  
5 3,3-dimethylbutylamine};

(1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylamine;

10 (1S)-1-([(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl)-  
2-methylpropylamine;

(1S)-1-cyclohexyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;

15  
(1S)-1-[[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]but-3-enylamine;

(1S)-1-[[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-ynylamine;

(1S)-1-benzyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine;

25 (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;

1-[1-(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine;

30 (1S)-1-{[(2S)-4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylamine;

(1S)-1-[[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]pentylamine;

- (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-methylbutylamine;
- 5 (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3,3-dimethylbutylamine;
- 1-cyclopropyl-3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-amine;
- 10 (1S)-2-[(2S)-4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-cyclopropyl-2-oxoethanamine;
- (1S)-1-{[(2S)-4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylpropylamine;
- 15 (1S,2S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylbutylamine;
- 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole;
- 20 (2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole;
- 25 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole;
- 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole;
- (2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole;
- 30 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole;

- 3-[1-[(2S)-2-amino-2-cyclopropylethanoyl]-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 5 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-isoleucyl-2,5-dihydro-1H-pyrrole;
- 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-norleucyl-2,5-dihydro-1H-pyrrole;
- 10 (2S)-4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-(3-methyl-L-valyl)-2,5-dihydro-1H-pyrrole;
- (2S)-4-(2,5-Difluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 (2S)-4-(2,5-Difluorophenyl)-N-methyl-2-phenyl-N-(piperidin-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(5-Chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-[(3R)-pyrrolidin-3-yl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 (2S)-4-(5-chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(5-chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-[(3S)-pyrrolidin-3-yl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 (2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-pyrrolidin-3-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 (2S)-4-(2,5-difluorophenyl)-N-methyl-N-[(3S)-1-methylpyrrolidin-3-yl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(2,5-difluorophenyl)-N-methyl-N-[(3R)-1-methylpyrrolidin-3-yl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 35

- (2S)-4-(2,5-difluorophenyl)-N-methyl-N-[(1-methyl-5-oxopyrrolidin-2-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 (2S)-4-(2,5-difluorophenyl)-N-(1,3-dioxolan-2-ylmethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-tetrahydrofuran-3-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 (2S)-N-(1-allylpiperidin-4-yl)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- allyl 4-[[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]piperidine-1-carboxylate;
- 15 allyl 4-[[[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]methyl]piperidine-1-carboxylate;
- (2S)-4-(2,5-difluorophenyl)-N-methyl-N-(1-methylpiperidin-4-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 4-(2,5-Difluorophenyl)-N-methyl-N-[(1-methylpiperidin-3-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(pyridin-3-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- N-benzyl-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 4-(2,5-difluorophenyl)-N-methyl-N-[(1-methyl-1H-pyrazol-4-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-[2-(dimethylamino)ethyl]-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 35

- 4-(2,5-difluorophenyl)-N-(2-hydroxyethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 4-(2,5-difluorophenyl)-N-isobutyl-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(2-pyridin-2-ylethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 4-(2,5-difluorophenyl)-N-(2-methoxyethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-(2,3-dihydroxypropyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(2-phenylethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-phenyl-N-propyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 1-Acetyl-4-(2,5-difluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole;
- (2S)-1-[4-(2,5-difluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methyl-1-oxobutan-2-amine;
- 30 (2S)-4-(2,5-difluorophenyl)-N,N,2-trimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

- (2S)-4-(5-chloro-2-fluorophenyl)-N,N,2-trimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 (2S)-4-(5-chloro-2-fluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(5-chloro-2-fluorophenyl)-N-ethyl-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 (2S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3,3-dimethyl-1-oxobutan-2-ol;
- (2S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methyl-1-oxobutan-2-ol;
- 15 (2S,3S)-1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methyl-1-oxopentan-2-ol;
- 1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-oxohexan-2-ol;
- 20 (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-oxo-3-phenylpropan-2-ol;
- 25 (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-methyl-1-oxopentan-2-ol;
- (1S)-1-cyclohexyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanol;
- 30 (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3,3-dimethyl-1-oxobutan-2-ol;
- N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-N-2,N-2-dimethylglycinamide;
- 35



- N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2-methylglycinamide;
- 5 N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)glycinamide;
- N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-methylalaninamide;
- 10 N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)glycinamide;
- N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2,N-2-dimethylglycinamide;
- 15 N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2,N-2-dimethylglycinamide, N-oxide;
- N-1-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-methylalaninamide;
- 20 N-1-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-N-2,N-2-dimethylglycinamide n-oxide;
- 25 N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-pyrrolidin-1-ylacetamide;
- 2-azetidin-1-yl-N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)acetamide;
- 30 N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-morpholin-4-ylacetamide;
- N-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)-2-piperazin-1-ylacetamide;
- 35

- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(4-methylpiperazin-1-yl)acetamide;
- 5 2-azetidin-1-yl-N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}acetamide;
- N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-pyrrolidin-1-ylacetamide;
- 10 N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-piperidin-1-ylacetamide;
- N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-morpholin-4-ylacetamide;
- 15 N-1-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N-2-(2-hydroxyethyl)glycinamide;
- N-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(4-methylpiperazin-1-yl)acetamide;
- 20 N-1-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N-2-isopropylglycinamide;
- 25 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}acetamide;
- N-1-{(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N-2-ethylglycinamide;
- 30 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-hydroxyacetamide;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}piperazine-1-carboxamide;
- 35

- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-piperidin-4-ylurea;
- 5 4-amino-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}piperidine-1-carboxamide;
- N-(2-aminoethyl)-N'-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}urea;
- 10 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-(3-morpholin-4-ylpropyl)urea;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-[2-(dimethylamino)ethyl]urea;
- 15 2-azetidin-1-yl-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}ethanesulfonamide
- 20 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(isopropylamino)ethanesulfonamide;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-pyrrolidin-1-ylethanesulfonamide;
- 25 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-morpholin-4-ylethanesulfonamide;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-piperazin-1-ylethanesulfonamide;
- 30 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(4-methylpiperazin-1-yl)ethanesulfonamide;

- N-(tert-butyl)-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetamide;
- 5 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylacetamide;
- (2S)-1-(2-azetidin-1-yl-2-oxoethyl)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 10 (2S)-4-(2,5-difluorophenyl)-1-(2-oxo-2-pyrrolidin-1-ylethyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl}morpholine;
- 15 1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl}piperazine;
- 1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl}-4-methylpiperazine;
- 20 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylbutanamide;
- 4-{2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]butanoyl}morpholine;
- 25 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethylacetamide;
- 30 N-cyclobutyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetamide;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethylpropanamide;
- 35

- N-cyclobutyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanamide;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methylpropanamide;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylpropanamide;
- 10 N-(tert-butyl)-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanamide;
- 4-{2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoyl}morpholine;
- 15 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethyl-2,2-dimethyl-4-oxobutanamide;
- (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxo-N-piperidin-4-ylbutanamide;
- 20 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid;
- (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2,2-tetramethyl-4-oxobutanamide;
- 25 (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethyl-3-oxo-3-piperazin-1-ylpropylamine;
- 30 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropyl-2,2-dimethyl-4-oxobutanamide;
- (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,2,2-trimethyl-4-oxobutanamide;
- 35

(3R)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2,2-tetramethyl-4-oxobutanamide;

5 (3R)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid;

(1R)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethyl-3-oxo-3-piperazin-1-ylpropylamine;

10 2-({(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-ethylacetamide;

2-({(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-methylacetamide;

15 2-({(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N,N-dimethylacetamide;

2-({(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-methyl-N-ethylacetamide;

20 2-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-methylacetamide;

2-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-ethylacetamide;

2-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N,N-dimethylacetamide;

30 2-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-isopropylacetamide;

2-({(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-ethyl-N-methylacetamide;

35

2-((1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)-N,N-diethylacetamide;

(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-N-(2-oxo-2-pyrrolidin-1-ylethyl)ethanamine;

(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-(2-morpholin-4-yl-2-oxoethyl)-2-oxoethanamine;

1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino]acetyl]piperidin-4-ol;

(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-[2-(4-methylpiperazin-1-yl)-2-oxoethyl]-2-oxoethanamine;

(1S)-N-(2-azetidin-1-yl-2-oxoethyl)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;

(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-[2-(1,1-dioxidothiomorpholin-4-yl)-2-oxoethyl]-2-oxoethanamine;

(1S)-N-[2-(4-acetylpiperazin-1-yl)-2-oxoethyl]-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;

(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-(2-morpholin-4-yl-2-oxoethyl)-2-oxoethanamine;

(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-N-(2-oxo-2-pyrrolidin-1-ylethyl)ethanamine;

2-((1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl)amino)-N-isopropylacetamide;

2-(dimethylamino)ethyl (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate;

- 1-methylpiperidin-4-yl (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylcarbamate;
- (2S)-4-cyclopropyl-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 (2S)-4-cyclopentyl-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 4-methylpiperazine-1-carboxylate;
- 10 1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 2-morpholin-4-ylethylcarbamate;
- N-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]oxy-carbonyl]glycine;
- 15 (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 1-methylpiperidin-4-ylcarbamate;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylmethyl(1-methylpiperidin-4-yl)carbamate;
- 20 (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 4-dimethylamino)piperidine-1-carboxylate;
- 25 tert-butyl (2S)-4-(2-chloro-5-fluoropyrimidin-4-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxylate;
- (2S)-4-(5-fluoro-2-methylpyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 (2S)-4-(2-chloro-5-fluoropyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;



- (2S)-4-(4-chloro-5-methylpyrimidin-2-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 (2S)-4-(6-chloropyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(2-chloropyrimidin-4-yl)-N,N-dimethyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 (2S)-N,N-dimethyl-4-(4-methylpyridin-3-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-N,N-dimethyl-2-phenyl-4-(1,3-thiazol-2-yl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 (2S)-N,N-dimethyl-2-phenyl-4-(1,3-thiazol-4-yl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-N,N-dimethyl-2-phenyl-4-(1,2-thiazol-5-yl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 (2S)-4-(2,5-difluorophenyl)-N-(2-hydroxyethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 N-{[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-N-methyl-beta-alanine;
- methyl N-{[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-N-methyl-beta-alaninate;
- 35

- 4-{{4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl}acetyl}morpholin-4-ium;
- 3-[(2S)-4-(2,5-difluorophenyl)-1-(2-hydroxy-2-methylpropanoyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-sulfonamide;
- 3-[4-(2,5-difluorophenyl)-1-(methylsulfonyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 3-[4-(2,5-difluorophenyl)-1-(morpholin-4-ylcarbonyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 3-[4-(2,5-difluorophenyl)-1-(2,2-dimethylpropanoyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- (2S)-4-(2,5-difluorophenyl)-1-[(methylsulfonyl)acetyl]-2-phenyl-2,5-dihydro-1H-pyrrole;
- (2S)-4-(2,5-difluorophenyl)-2-phenyl-1-[(phenylsulfonyl)acetyl]-2,5-dihydro-1H-pyrrole;
- 3-[(2S)-1-[(2S)-2-cyclopropyl-2-hydroxyethanoyl]-4-(2,5-difluorophenyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 3-{{(2S)-4-(2,5-difluorophenyl)-1-[(2S)-2-hydroxy-3,3-dimethylbutanoyl]-2,5-dihydro-1H-pyrrol-2-yl}phenol};
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanol;
- (2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-tetrahydrofuran-3-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(2-methoxyethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

- (2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-tetrahydro-2H-pyran-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 3-[(2S)-4-(2,5-difluorophenyl)-1-(piperidin-1-ylcarbonyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 4-(2,5-difluorophenyl)-N-[1-(2-fluoroethyl)piperidin-4-yl]-2-(3-hydroxyphenyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]piperidinium trifluoroacetate;
- 2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethyl 4-methylpiperazine-1-carboxylate;
- 3-{4-(2,5-difluorophenyl)-1-[(4-methylpiperazin-1-yl)carbonyl]-2,5-dihydro-1H-pyrrol-2-yl}phenol;
- 2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethyl morpholine-4-carboxylate;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(1-methyl-1H-pyrazol-4-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(isoxazol-5-ylmethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethyl dimethylaminocarboxylate;
- 2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethyl piperidine-1-carboxylate;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;

- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(5-oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(tetrahydro-2H-pyran-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-[[5-(methoxymethyl)-1H-pyrazol-3-yl]methyl]-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(4-methyl-1,2,5-oxadiazol-3-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-2-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(isoxazol-3-ylmethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-1,2,4-triazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-pyrazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(1-methyl-5-oxopyrrolidin-2-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(1-isoxazol-3-ylethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

- 4-(2,5-difluorophenyl)-N-(1,3-dioxolan-2-ylmethyl)-2-(3-hydroxyphenyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-(1,4-dioxan-2-ylmethyl)-2-(3-hydroxyphenyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(5-methyl-1,3,4-oxadiazol-2-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(methylsulfonyl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethanesulfonic acid;
- 2-hydroxyethyl (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylcarbamate;
- 3-hydroxypropyl (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylcarbamate;
- 2-hydroxyethyl {(1S)-1-isopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}carbamate;
- 2-hydroxyethyl {(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}carbamate;
- 4-hydroxybutyl (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylcarbamate;
- (2S)-4-(2,5-difluorophenyl)-1-[2-(methylsulfonyl)ethyl]-2-phenyl-2,5-dihydro-1H-pyrrole;
- (2S)-4-(2,5-difluorophenyl)-1-[2-(ethylsulfonyl)ethyl]-2-phenyl-2,5-dihydro-1H-pyrrole;
- 1-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]pentan-3-one;

- 4-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]butan-2-one;
- 5 4-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-3-methylbutan-2-one;
- 2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-*N,N*-dimethylethanesulfonamide;
- 10 3-{(2*S*)-4-(2,5-difluorophenyl)-1-[2-(methylsulfonyl)ethyl]-2,5-dihydro-1*H*-pyrrol-2-yl}phenol;
- methyl 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoate;
- 15 (2*S*)-4-(2,5-difluorophenyl)-1-[2-(ethylsulfonyl)propyl]-2-phenyl-2,5-dihydro-1*H*-pyrrole;
- 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-*N*-methylpropanamide;
- 20 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-*N,N*-dimethylpropanamide;
- 25 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-*N,N*,2-trimethylpropanamide;
- 4-{3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoyl}morpholine;
- 30 1-{3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoyl}-4-(methylsulfonyl)piperazine;
- 1-{3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoyl}piperidin-4-ol;
- 35 methyl 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoate;
- 40 2-({(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl}oxy)-*N*-ethylacetamide ;
- 4-({(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy}acetyl)morpholine;
- 45

- 2-((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy)-*N*-(2-hydroxyethyl)acetamide;
- 5 1-((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy)acetyl)-4-methylpiperazine;
- 1-((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy)acetyl)piperazine;
- 10 2-((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy)-*N*-piperidin-4-ylacetamide;
- 1-((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy)acetyl)piperidin-4-amine;
- 15 *N*-((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl)-3-morpholin-4-yl-3-oxopropan-1-amine;
- 20 *N*<sup>3</sup>-((1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl)-*N*<sup>1</sup>,*N*<sup>1</sup>-dimethyl-β-alaninamide; and
- ((1*S*)-1-((2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl)carbonyl)-2,2-dimethylpropyl(3-morpholin-4-yl-3-oxopropyl)amine;
- 25 or a pharmaceutically acceptable salt or stereoisomer thereof.

12. The compound according to Claim 11 which is selected from:

- 30 (-)-4-(2,5-difluorophenyl)-*N,N*-dimethyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide;
- 4-(5-chloro-2-fluorophenyl)-*N,N*-dimethyl-2-phenyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide;
- 35 4-(2,5-difluorophenyl)-1-(isopropylsulfonyl)-2-phenyl-2,5-dihydro-1*H*-pyrrole;
- 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-methyl-1-oxopropan-2-ol;

1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine;

(1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-  
5 2,2-dimethylpropylamine; and

(2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;

10 or a pharmaceutically acceptable salt or stereoisomer thereof.

13. The compound according to Claim 11 which is selected from:

(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-  
15 1-yl]-2-oxoethanamine;

(2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;

20 (1S)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethylpropylamine;

2-({(1S)-1-tert-butyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}amino)-N-ethylacetamide ;

25 (2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N,N-dimethyl-2,5-dihydro-1H-pyrrole-1-carboxamide;

(2S)-4-(2,5-difluorophenyl)-N-methyl-N-(1-methylpiperidin-4-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide; and  
30

(2S)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole

or a pharmaceutically acceptable salt or stereoisomer thereof.

35



14. A compound which is:

- (1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethanamine
- 5 (2*S*)-4-(2,5-difluorophenyl)-1-[(methylsulfonyl)acetyl]-2-phenyl-2,5-dihydro-1*H*-pyrrole
- 10 (2*S*)-4-(2,5-difluorophenyl)-2-phenyl-1-[(phenylsulfonyl)acetyl]-2,5-dihydro-1*H*-pyrrole
- 3-[(2*S*)-1-[(2*S*)-2-cyclopropyl-2-hydroxyethanoyl]-4-(2,5-difluorophenyl)-2,5-dihydro-1*H*-pyrrol-2-yl]phenol
- 15 3-[(2*S*)-4-(2,5-difluorophenyl)-1-[(2*S*)-2-hydroxy-3,3-dimethylbutanoyl]-2,5-dihydro-1*H*-pyrrol-2-yl]phenol
- (1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethanol
- 20 (2*S*)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-*N*-methyl-*N*-tetrahydrofuran-3-yl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- (2*S*)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-*N*-(2-methoxyethyl)-*N*-methyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 25 (2*S*)-4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-*N*-methyl-*N*-tetrahydro-2*H*-pyran-4-yl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 30 3-[(2*S*)-4-(2,5-difluorophenyl)-1-(piperidin-1-ylcarbonyl)-2,5-dihydro-1*H*-pyrrol-2-yl]phenol
- 4-(2,5-difluorophenyl)-*N*-[1-(2-fluoroethyl)piperidin-4-yl]-2-(3-hydroxyphenyl)-*N*-methyl-2,5-dihydro-1*H*-pyrrole-1-carboxamide
- 35 4-[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1*H*-pyrrol-1-yl]carbonyl](methylamino)piperidinium trifluoroacetate

- 2-[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methylamino)ethyl 4-methylpiperazine-1-carboxylate
- 3-{4-(2,5-difluorophenyl)-1-[(4-methylpiperazin-1-yl)carbonyl]-2,5-dihydro-1H-pyrrol-2-yl}phenol
- 2-[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methylamino)ethyl morpholine-4-carboxylate
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(1-methyl-1H-pyrazol-4-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(isoxazol-5-ylmethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide
- 2-[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methylamino)ethyl dimethylaminocarboxylate
- 2-[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methylamino)ethyl piperidine-1-carboxylate
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(2-oxopyrrolidin-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(5-oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(tetrahydro-2H-pyran-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-{[5-(methoxymethyl)-1H-pyrazol-3-yl]methyl}-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide

- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(4-methyl-1,2,5-oxadiazol-3-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide
- 5 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-2-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(isoxazol-3-ylmethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide
- 10 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-1,2,4-triazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-pyrazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide
- 15 2,5-dihydro-1H-pyrrole-1-carboxamide
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(1-methyl-5-oxopyrrolidin-2-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide
- 20 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-(1-isoxazol-3-ylethyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide
- 4-(2,5-difluorophenyl)-N-(1,3-dioxolan-2-ylmethyl)-2-(3-hydroxyphenyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide
- 25 4-(2,5-difluorophenyl)-N-(1,4-dioxan-2-ylmethyl)-2-(3-hydroxyphenyl)-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(5-methyl-1,3,4-oxadiazol-2-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide
- 30 2,5-dihydro-1H-pyrrole-1-carboxamide
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(methylsulfonyl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide
- 35 2-[[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethanesulfonic acid

- 2-hydroxyethyl (1S)-1-[[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-2,2-dimethylpropylcarbamate
- 5 3-hydroxypropyl (1S)-1-[[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-2,2-dimethylpropylcarbamate
- 2-hydroxyethyl {(1S)-1-isopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}carbamate
- 10 2-hydroxyethyl {(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}carbamate
- 4-hydroxybutyl (1S)-1-[[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-2,2-dimethylpropylcarbamate
- 15 (2S)-4-(2,5-difluorophenyl)-1-[2-(methylsulfonyl)ethyl]-2-phenyl-2,5-dihydro-1H-pyrrole
- 20 (2S)-4-(2,5-difluorophenyl)-1-[2-(ethylsulfonyl)ethyl]-2-phenyl-2,5-dihydro-1H-pyrrole
- 1-[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]pentan-3-one
- 4-[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]butan-2-one
- 25 4-[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methylbutan-2-one
- 2-[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-*N,N*-dimethylethanesulfonamide
- 30 3-{ (2S)-4-(2,5-difluorophenyl)-1-[2-(methylsulfonyl)ethyl]-2,5-dihydro-1H-pyrrol-2-yl}phenol
- 35 methyl 3-[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoate
- (2S)-4-(2,5-difluorophenyl)-1-[2-(ethylsulfonyl)propyl]-2-phenyl-2,5-dihydro-1H-pyrrole
- 40 3-[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-*N*-methylpropanamide
- 3-[ (2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-*N,N*-dimethylpropanamide
- 45

- 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-*N,N*,2-trimethylpropanamide
- 5 4-{3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoyl}morpholine
- 1-{3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoyl}-4-(methylsulfonyl)piperazine
- 10 1-{3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoyl}piperidin-4-ol
- methyl 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoate
- 15 2-({(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl}oxy)-*N*-ethylacetamide
- 20 4-({(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy}acetyl)morpholine
- 2-({(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy}-*N*-(2-hydroxyethyl)acetamide
- 25 1-({(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy}acetyl)-4-methylpiperazine
- 1-({(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy}acetyl)piperazine
- 30 2-({(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy}-*N*-piperidin-4-ylacetamide
- 35 1-({(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethoxy}acetyl)piperidin-4-amine
- N*-{(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl}-3-morpholin-4-yl-3-oxopropan-1-amine
- 40 *N*<sup>3</sup>-{(1*S*)-1-cyclopropyl-2-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]-2-oxoethyl}-*N*<sup>1</sup>,*N*<sup>1</sup>-dimethyl-β-alaninamide
- ((1*S*)-1-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]carbonyl)-2,2-dimethylpropyl(3-morpholin-4-yl-3-oxopropyl)amine
- 45

or a pharmaceutically acceptable salt thereof.

15. The compound according to Claim 12 which is the TFA salt of  
5 a compound selected from:

2-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]sulfonyl}-N,N-dimethylethanamine;

- 10 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-methyl-1-oxopropan-2-amine;

4-(5-chloro-2-fluorophenyl)-2-phenyl-1-(trifluoroacetyl)-2,5-dihydro-1H-pyrrole;

- 15 (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylpropylamine;

(1R)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylpropylamine;

20

4-(2,5-difluorophenyl)-2-phenyl-1-L-prolyl-2,5-dihydro-1H-pyrrole;

4-(2,5-difluorophenyl)-2-phenyl-1-D-prolyl-2,5-dihydro-1H-pyrrole;

- 25 (4R)-4-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-1,3-thiazolidine;

methyl (3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutanoate;

30

(4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanamide;

- (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-(methylthio)propylamine;
- 35

- (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-(methylsulfonyl)propylamine;
- 5 (2S)-2-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}piperidine;
- (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}pentylamine;
- 10 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(thien-2-ylmethyl)ethylamine;
- 4-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-1,1-dioxidotetrahydro-2H-thiopyran-4-ylamine;
- 15 (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methyl-1-oxopropan-2-amine;
- 20 (1S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}propylamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-phenylethanamine;
- 25 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-phenylethanamine;
- (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanamide
- 30 3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-amine;
- (1S,2S)-1-{{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylbutylamine;
- 35

- (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}butylamine;
- 5 (1S)-1-cyclopropyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopropanamine;
- 10 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-oxopropan-2-amine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-methyl-2-
- 15 oxoethylamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(pyridin-2-ylmethyl)ethylamine;
- 20 (1S)-1-cyclohexyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(4-iodobenzyl)-2-oxoethylamine;
- 25 (1S)-1-benzyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine;
- 4-{(2S)-2-amino-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropyl}phenol;
- 30 (3S)-3-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-1,2,3,4-tetrahydroisoquinoline;
- (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3-phenylpropylamine;
- 35



- (1S)-1-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-3-methylbutylamine;
- 5 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(pyridin-3-ylmethyl)ethylamine;
- 1-[(2S)-azetidin-2-ylcarbonyl]-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 10 (3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutanamide;
- 4-(2,5-difluorophenyl)-1-[(2-methylazetidin-2-yl)carbonyl]-2-phenyl-2,5-dihydro-1H-pyrrole;
- 15 (1S)-1-[[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-2,2-dimethylpropylamine;
- 20 methyl (4S)-4-amino-5-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-5-oxopentanoate;
- 4-(2,5-difluorophenyl)-2-phenyl-1-[[4-(2S,3S)-2-phenylpyrrolidin-3-yl]carbonyl]-2,5-dihydro-1H-pyrrole;
- 25 4-(2,5-difluorophenyl)-2-phenyl-1-[(5-phenylpyrrolidin-3-yl)carbonyl]-2,5-dihydro-1H-pyrrole;
- (2S)-2-amino-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-ol;
- 30 (2R,3S)-3-amino-4-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-4-oxobutan-2-ol;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(methoxymethyl)-2-oxoethylamine;
- 35

- 4-(2,5-difluorophenyl)-2-phenyl-1-(pyrrolidin-3-ylcarbonyl)-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-2-phenyl-1-[(3-phenylpyrrolidin-3-yl)acetyl]-2,5-dihydro-1H-pyrrole;
- 5 (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-3,3-difluoropropylamine;
- 10 (1S)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxo-1-phenylpropan-1-amine;
- 4-(2,5-difluorophenyl)-2-phenyl-1-[(4S)-4-phenyl-L-prolyl]-2,5-dihydro-1H-pyrrole;
- 15 1-{2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}cyclohexanamine;
- 2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 20 4-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}piperidin-4-amine;
- (1S,3R)-3-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopentanamine;
- 25 (1R,4S)-4-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-2-en-1-amine;
- (1S,4R)-4-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}cyclopent-2-en-1-amine;
- 30 (1S)-1-{[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-ynylamine;

- (1R)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxo-1-phenylpropan-1-amine;
- 3-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-2-phenylpiperidine;
- 5 (1S)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}but-3-enylamine;
- 10 (2S)-3-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-(methylamino)-3-oxopropan-1-ol;
- (3R,5S)-5-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}pyrrolidin-3-ol;
- 15 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxo-1-(1,3-thiazol-4-ylmethyl)ethylamine;
- (1R)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}but-3-enylamine;
- 20 (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,3-dimethyl-1-oxobutan-2-amine;
- 25 (2S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,4-dimethyl-1-oxopentan-2-amine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-[(1-methyl-1H-imidazol-4-yl)methyl]-2-oxoethylamine;
- 30 4-(2,5-difluorophenyl)-1-(N-6-formyl-L-lysyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- (2S,3S)-1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,3-dimethyl-1-oxopentan-2-amine;
- 35

- (1S)-1-(cyclohexylmethyl)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylamine;
- 5 (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxoethylamine;
- (1S)-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-(isocyanomethyl)-2-oxoethylamine;
- 10 (1S)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}-3,3-dimethylbutylamine;
- 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,3-dimethyl-1-oxobutan-2-amine;
- 15 1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}cyclohexanamine;
- 1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}cyclopentanamine;
- 20 (1S)-3-(benzyloxy)-1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}propylamine;
- 25 1-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,3-dimethyl-1-oxobutan-2-amine;
- 1-{{4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl}carbonyl}cyclopent-3-en-1-amine;
- 30 (1S)-1-cyclopentyl-2-[4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 4-(2,5-difluorophenyl)-1-(2-methylpropyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 35

- (1S)-2-[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-1-cyclopropyl-2-oxoethanamine;
- 5 (1S,2S)-1-{{[4-(5-chloro-2-fluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylbutylamine;
- (1S)-1-{{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-enylamine;
- 10 (1S)-1-{{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}but-3-ynylamine;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethanamine;
- 15 1-cyclopropyl-3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-oxopropan-1-amine;
- (1S,2S)-1-{{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2-methylbutylamine;
- 20 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-L-valyl-2,5-dihydro-1H-pyrrole;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-1-(2-methylalanyl)-2,5-dihydro-1H-pyrrole;
- 25 3-[1-[(2S)-2-amino-2-cyclopropylethanoyl]-4-(5-chloro-2-fluorophenyl)-2,5-dihydro-1H-pyrrol-2-yl]phenol;
- 30 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-1-L-isoleucyl-2,5-dihydro-1H-pyrrole;
- (2S)-4-(2,5-Difluorophenyl)-N-methyl-2-phenyl-N-(piperidin-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 35

- (2S)-4-(5-Chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-[(3R)-pyrrolidin-3-yl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 5 (2S)-4-(5-chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-4-(5-chloro-2-fluorophenyl)-N-methyl-2-phenyl-N-[(3S)-pyrrolidin-3-yl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 10 (2S)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-pyrrolidin-3-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- (2S)-N-(1-allylpiperidin-4-yl)-4-(2,5-difluorophenyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 4-(2,5-Difluorophenyl)-N-methyl-N-[(1-methylpiperidin-3-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(pyridin-3-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 4-(2,5-difluorophenyl)-N-methyl-N-[(1-methyl-1H-pyrazol-4-yl)methyl]-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 4-(2,5-difluorophenyl)-N-[2-(dimethylamino)ethyl]-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-N-methyl-2-phenyl-N-(2-pyridin-2-ylethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 30 4-(5-chloro-2-fluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 35

- (2S)-1-[4-(2,5-difluorophenyl)-2-methyl-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methyl-1-oxobutan-2-amine;
- 5 N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-N-2,N-2-dimethylglycinamide;
- N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-N-2-methylglycinamide;
- 10 N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]glycinamide;
- N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-2-methylalaninamide;
- 15 N-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-2-pyrrolidin-1-ylacetamide;
- 20 2-azetidin-1-yl-N-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]acetamide;
- N-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-2-morpholin-4-ylacetamide;
- 25 N-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-2-piperazin-1-ylacetamide;
- N-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-2-(4-methylpiperazin-1-yl)acetamide;
- 30 N-1-[(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl]-N-2-isopropylglycinamide;

- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}piperazine-1-carboxamide;
- 5 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-piperidin-4-ylurea;
- 4-amino-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}piperidine-1-carboxamide;
- 10 N-(2-aminoethyl)-N'-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}urea;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-(3-morpholin-4-ylpropyl)urea;
- 15 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-N'-[2-(dimethylamino)ethyl]urea;
- 2-azetidin-1-yl-N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}ethanesulfonamide
- 20 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(isopropylamino)ethanesulfonamide;
- 25 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-pyrrolidin-1-ylethanesulfonamide;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-morpholin-4-ylethanesulfonamide;
- 30 N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-piperazin-1-ylethanesulfonamide;
- N-{(1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl}-2-(4-methylpiperazin-1-yl)ethanesulfonamide;
- 35



- N-(tert-butyl)-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetamide;
- 5 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylacetamide;
- (2S)-1-(2-azetidin-1-yl-2-oxoethyl)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 10 (2S)-4-(2,5-difluorophenyl)-1-(2-oxo-2-pyrrolidin-1-ylethyl)-2-phenyl-2,5-dihydro-1H-pyrrole;
- 4-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl}morpholine;
- 15 1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl}piperazine;
- 20 1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetyl}-4-methylpiperazine;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylbutanamide;
- 25 4-{2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]butanoyl}morpholine;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethylacetamide;
- 30 N-cyclobutyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]acetamide;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethylpropanamide;
- 35

- N-cyclobutyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanamide;
- 5 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methylpropanamide;
- 2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropylpropanamide;
- 10 N-(tert-butyl)-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanamide;
- 4-{2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoyl}morpholine;
- 15 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-ethyl-2,2-dimethyl-4-oxobutanamide;
- (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxo-N-piperidin-4-ylbutanamide;
- 20 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid;
- 25 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2,2-tetramethyl-4-oxobutanamide;
- (1S)-1-[[[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl]-2,2-dimethyl-3-oxo-3-piperazin-1-yl]propylamine;
- 30 (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-isopropyl-2,2-dimethyl-4-oxobutanamide;

- (3S)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,2,2-trimethyl-4-oxobutanamide;
- (3R)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2,2-tetramethyl-4-oxobutanamide;
- (3R)-3-amino-4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2,2-dimethyl-4-oxobutanoic acid;
- (1R)-1-{[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]carbonyl}-2,2-dimethyl-3-oxo-3-piperazin-1-ylpropylamine
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 4-methylpiperazine-1-carboxylate;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl 1-methylpiperidin-4-ylcarbamate;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethylmethyl(1-methylpiperidin-4-yl)carbamate;
- (1S)-1-cyclopropyl-2-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-2-oxoethyl-4-dimethylamino)piperidine-1-carboxylate;
- (2S)-N,N-dimethyl-4-(4-methylpyridin-3-yl)-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-{[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]acetyl}morpholin-4-ium;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-piperidin-4-yl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 2-[[4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-2,5-dihydro-1H-pyrrol-1-yl]carbonyl](methyl)amino]ethyl-4-methylpiperazine-1-carboxylate;

- 3-{4-(2,5-difluorophenyl)-1-[(4-methylpiperazin-1-yl)carbonyl]-2,5-dihydro-1H-pyrrol-2-yl}phenol;
- 5 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(1-methyl-1H-pyrazol-4-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluoro-phenyl)-2,5-dihydro-2-(3-hydroxyphenyl)-N-methyl-N--[2-(4-methyl-1-piperazinyl)-2-oxoethyl]-H-pyrrole-1-carboxamide;
- 10 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[(5-oxo-4,5-dihydro-1H-1,2,4-triazol-3-yl)methyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 15 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-{[5-(methoxymethyl)-1H-pyrazol-3-yl]methyl}-N-methyl-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-4-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 20 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-(1,3-thiazol-2-ylmethyl)-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-1,2,4-triazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 25 4-(2,5-difluorophenyl)-2-(3-hydroxyphenyl)-N-methyl-N-[2-(1H-pyrazol-1-yl)ethyl]-2,5-dihydro-1H-pyrrole-1-carboxamide;
- 4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]butan-2-one;
- 30 4-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-3-methylbutan-2-one;
- 3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N-methylpropanamide;
- 35 3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N-dimethylpropanamide;
- 40 3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]-N,N,2-trimethylpropanamide;
- 4-{3-[(2S)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1H-pyrrol-1-yl]propanoyl}morpholine;
- 45

1-{3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoyl}-4-(methylsulfonyl)piperazine;

5 1-{3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoyl}piperidin-4-ol; and

methyl 3-[(2*S*)-4-(2,5-difluorophenyl)-2-phenyl-2,5-dihydro-1*H*-pyrrol-1-yl]propanoate.

10 16. A pharmaceutical composition that is comprised of a compound in accordance with Claim 1 and a pharmaceutically acceptable carrier.

15 17. A method of treating or preventing cancer in a mammal in need of such treatment that is comprised of administering to said mammal a therapeutically effective amount of a compound of Claim 1.

20 18. A method of treating cancer or preventing cancer in accordance with Claim 17 wherein the cancer is selected from cancers of the brain, genitourinary tract, lymphatic system, stomach, larynx and lung.

25 19. A method of treating or preventing cancer in accordance with Claim 17 wherein the cancer is selected from histiocytic lymphoma, lung adenocarcinoma, small cell lung cancers, pancreatic cancer, glioblastomas and breast carcinoma.

20. A process for making a pharmaceutical composition which comprises combining a compound of Claim 1 with a pharmaceutically acceptable carrier.

30 21. The composition of Claim 16 further comprising a second compound selected from:

- 35
- 1) an estrogen receptor modulator,
  - 2) an androgen receptor modulator,
  - 3) a retinoid receptor modulator,
  - 4) a cytotoxic/cytostatic agent,
  - 5) an antiproliferative agent,

- 5
- 6) a prenyl-protein transferase inhibitor,
  - 7) an HMG-CoA reductase inhibitor,
  - 8) an HIV protease inhibitor,
  - 9) a reverse transcriptase inhibitor,
  - 10) an angiogenesis inhibitor, and
  - 11) a PPAR- $\gamma$  agonist,
  - 12) a PPAR- $\delta$  agonists;
  - 13) an inhibitor of cell proliferation and survival signaling, and
  - 14) an agent that interferes with a cell cycle checkpoint.

10

22. The composition of Claim 21, wherein the second compound is an angiogenesis inhibitor selected from the group consisting of a tyrosine kinase inhibitor, an inhibitor of epidermal-derived growth factor, an inhibitor of fibroblast-derived growth factor, an inhibitor of platelet derived growth factor, an MMP  
15 inhibitor, an integrin blocker, interferon- $\alpha$ , interleukin-12, pentosan polysulfate, a cyclooxygenase inhibitor, carboxyamidotriazole, combretastatin A-4, squalamine, 6-O-(chloroacetyl-carbonyl)-fumagillol, thalidomide, angiostatin, troponin-1, and an antibody to VEGF.

20

23. The composition according to Claim 16 further comprising a proteasome inhibitor.

25

24. The composition according to Claim 16 further comprising a aurora kinase inhibitor.

25. The composition according to Claim 16 further comprising a Raf kinase inhibitor.

30

26. The composition according to Claim 16 further comprising a serine/threonine kinase inhibitor.

27. The composition according to Claim 16 further comprising an inhibitor of another mitotic kinesin which is not KSP.

28. The composition of Claim 22, wherein the second compound is an estrogen receptor modulator selected from tamoxifen and raloxifene.

5 29. A method of treating cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with radiation therapy.

30. A method of treating or preventing cancer that comprises administering a therapeutically effective amount of a compound of Claim 1 in  
10 combination with a compound selected from:

- 1) an estrogen receptor modulator,
- 2) an androgen receptor modulator,
- 3) a retinoid receptor modulator,
- 4) a cytotoxic/cytostatic agent,
- 15 5) an antiproliferative agent,
- 6) a prenyl-protein transferase inhibitor,
- 7) an HMG-CoA reductase inhibitor,
- 8) an HIV protease inhibitor,
- 9) a reverse transcriptase inhibitor,
- 20 10) an angiogenesis inhibitor,
- 11) PPAR- $\gamma$  agonists,
- 12) PPAR- $\delta$  agonists,
- 13) an inhibitor of inherent multidrug resistance,
- 14) an anti-emetic agent,
- 25 15) an agent useful in the treatment of anemia,
- 16) an agent useful in the treatment of neutropenia,
- 17) an immunologic-enhancing drug,
- 18) an inhibitor of cell proliferation and survival signaling, and
- 19) an agent that interferes with a cell cycle checkpoint.

30

31. A method of treating cancer that comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with radiation therapy and a compound selected from:

- 1) an estrogen receptor modulator,
- 35 2) an androgen receptor modulator,

- 5                   3)     a retinoid receptor modulator,  
                  4)     a cytotoxic/cytostatic agent,  
                  5)     an antiproliferative agent,  
                  6)     a prenyl-protein transferase inhibitor,  
                  7)     an HMG-CoA reductase inhibitor,  
                  8)     an HIV protease inhibitor,  
                  9)     a reverse transcriptase inhibitor,  
                  10)    an angiogenesis inhibitor,  
                  11)    PPAR- $\gamma$  agonists,  
10               12)    PPAR- $\delta$  agonists,  
                  13)    an inhibitor of inherent multidrug resistance,  
                  14)    an anti-emetic agent,  
                  15)    an agent useful in the treatment of anemia,  
                  16)    an agent useful in the treatment of neutropenia,  
15               17)    an immunologic-enhancing drug,  
                  18)    an inhibitor of cell proliferation and survival signaling, and  
                  19)    an agent that interferes with a cell cycle checkpoint.

20               32.     A method of treating or preventing cancer which comprises  
administering a therapeutically effective amount of a compound of Claim 1 and  
paclitaxel or trastuzumab.

25               33.     A method of treating or preventing cancer which comprises  
administering a therapeutically effective amount of a compound of Claim 1 and a  
GPIIb/IIIa antagonist.

                  34.     The method of Claim 33 wherein the GPIIb/IIIa antagonist is  
tirofiban.

30               35.     A method of treating or preventing cancer which comprises  
administering a therapeutically effective amount of a compound of Claim 1 in  
combination with a COX-2 inhibitor.



36. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a proteasome inhibitor.
- 5 37. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with an aurora kinase inhibitor.
- 10 38. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a Raf kinase inhibitor.
- 15 39. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with a serine/threonine kinase inhibitor.
- 20 40. A method of treating or preventing cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with an inhibitor of a mitotic kinesin that is not KSP.
41. A method of modulating mitotic spindle formation which comprises administering a therapeutically effective amount of a compound of Claim 1.
- 25 42. A method of inhibiting the mitotic kinesin KSP which comprises administering a therapeutically effective amount of a compound of Claim 1.

## SEQUENCE LISTING

<110> Merck & Co., Inc.  
Arrington, Kenneth L.  
Coleman, Paul J.  
Cox, Christopher D.  
Fraley, Mark E.  
Garbaccio, Robert M.  
Hartman, George D.  
Hoffman, William F.  
Tasber, Edward S.

<120> MITOTIC KINESIN INHIBITORS

<130> 21114Y

<150> 60/388,621

<151> 2002-06-14

<150> 60/403,830

<151> 2002-08-15

<150> 60/426,940

<151> 2002-11-15

<150> 60/458,318

<151> 2003-03-28

<160> 2

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 42

<212> DNA

<213> Artificial Sequence

<220>

<223> Completely Synthetic Oligonucleotide

<400> 1

gcaacgatta atatggcgtc gcagccaaat tcgtctgcga ag

42

<210> 2

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> Completely Synthetic Oligonucleotide

<400> 2

gcaacgctcg agtcagtgat gatggtggtg atgctgattc acttcaggct tattcaatat 60

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/18482

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/535, 31/40; C07D 265/30, 207/02; A61P 43/00

US CL : 514/237.5, 423; 544/162; 548/538

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/237.5, 423; 544/162; 548/538

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
EAST, WEST

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
STN-CAS online

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3,644,414 (HELSEY et al) 22 February 1972 (22.02.1972), see entire reference.	11, 12 and 16-42

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

14 November 2003 (14.11.2003)

Date of mailing of the international search report

01 DEC 2003

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US  
Commissioner for Patents  
P.O. Box 1450  
Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Andrea D Small

Telephone No. (703) 308-1234

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/18482

### Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claim Nos.: 1-10, 13-15 and parts of 11 and parts of 16-42  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
Please See Continuation Sheet
3. ☐ Claim Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐  
☐

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

PCT/US03/18482

### Continuation of Box I Reason 2:

In these claims, the numerous variables (e.g. R1, R2, R3, R4, R5, R6, etc.) and their voluminous meanings and their seemingly endless permutations and combinations and their lengthy list of divergent compounds in claims 11, 13, 14 and 15, make it virtually impossible to determine the full scope and complete meaning of the claimed subject matter. As presented, the claimed subject matter cannot be regarded as being a clear and concise description for which protection is sought and as such the listed claims do not comply with the requirements of PCT, Article 6. Thus it is impossible to carry out a meaningful search on same. A search will be made on the first discernable inventions, namely the first 18 species named in claim 11, claim 12 in its entirety and parts of claims 16-42 that read on these named compounds.

## 203. Synthesen in der Emetin-Reihe

11. Mitteilung [1]<sup>1)</sup>2-Arylsubstituierte Hydrobenzo[a]chinolizine<sup>2)</sup>von H. Bruderer, M. Baumann, M. Uskoković und A. Brossi<sup>3)</sup>

(13. VIII. 64)

Verschiedene Benzo[a]chinolizine besitzen interessante pharmakologische Eigenschaften [2]. Zu dieser Gruppe kann man auch die beiden von GOOTJES & NAUTA [3] synthetisierten isomeren 2-*p*-Chlorphenyl-substituierten 9,10-Dimethoxy-1,2,3,4,6,7-hexahydro-benzo[a]chinolizine XI und XIII zählen. Nach den Angaben dieser Autoren sollen im analgetisch und antiphlogistisch wirksamen Isomer XIII die Ringe B/C *cis*-verknüpft sein<sup>5)</sup>.

Da bis dahin *cis*-verknüpfte Hydrobenzo[a]chinolizine nicht beschrieben wurden und die erwähnten Autoren keine näheren Angaben über die relative Konfiguration des 2-ständigen *p*-Chlorphenylrestes zum 11b-ständigen Wasserstoffatom gegeben haben, schien es von Interesse, die Struktur der erwähnten Verbindungen abzuklären. Über diese Untersuchungen wurde bereits an anderer Stelle ausführlich berichtet [4].

In der vorliegenden Mitteilung beschreiben wir die von uns zur Darstellung dieser Verbindungen entwickelte neue Synthese, die auch leicht zur Gewinnung von Strukturanalogen verwendet werden kann [5].

Bei der Umsetzung der beiden Ketone I [6] und II [7] mit einem Überschuss von *p*-Chlorphenylmagnesiumbromid entstehen in beiden Fällen Gemische von je zwei isomeren tert. Carbinolen (III, V und IV, VI)<sup>4)</sup>. Alle vier tert. Carbinole zeigen in ihren IR.-Spektren die BOHLMANN-Bande und können demzufolge als *trans*-verknüpfte Chinolizidine betrachtet werden. Die in ihren Formeln festgehaltene relative Konfiguration lässt sich aus ihren 60-MHz-Spektren ableiten (Tab. I).

GARBISCH & PATTERSON [8] haben kürzlich berichtet, dass im Fall der stereoisomeren 4-*t*-Butyl-1-phenyl-cyclohexanole das Hydroxyproton des Isomeren mit äquatorialer Hydroxylgruppe im Kernresonanzspektrum eine kleinere chemische Verschiebung aufweist als das Isomere mit axialer Hydroxylgruppe. Dieser Unterschied ist auf stärkere Wasserstoffbrückenbindung der äquatorialen Hydroxylgruppe zurückzuführen. Dementsprechend sollten die Verbindungen V und VI, in welchen die Hydroxylgruppen eine kleinere chemische Verschiebung aufweisen als in den damit

<sup>1)</sup> Die Zahlen in eckigen Klammern verweisen auf das Literaturverzeichnis, S. 1860.

<sup>2)</sup> Auszugsweise berichtet von einem von uns (M. U.) am 147. Meeting der Amer. chem. Ges. in der Sektion für Med. Chemie, in Philadelphia im April 1964. Vgl. Abstr. des Meetings, Div. of Medicinal Chemistry, S. 7M (1964).

<sup>3)</sup> Gegenwärtige Adresse: HOFFMANN-LA ROCHE INC., Nutley, New Jersey.

<sup>4)</sup> Die in dieser Arbeit beschriebenen Verbindungen stellen Racemate dar; die angegebenen Strukturformeln entsprechen daher jeweils absolut nur einem der beiden optischen Antipoden des Racemats.

<sup>5)</sup> *cis*-Verknüpfung: Das freie Elektronenpaar auf dem tert. Stickstoffatom ist *cis* ständig zum Wasserstoffatom am benachbarten tert. Kohlenstoffatom (11b) gelegen.